

- PHF6 mutation also found in the original Borjeson-Forssman-Lehmann syndrome family. *Eur J Hum Genet* 2004;12:787-9.
- 4 **Turner G**, Lower KM, White SM, Delatycki M, Lampe AK, Wright M, Smith JC, Kerr B, Schelley S, Hoyme HE, De Vries BB, Kleefstra T, Grømpe M, Cox B, Gecz J, Partington M. The clinical picture of the Borjeson-Forssman-Lehmann syndrome in males and heterozygous females with PHF6 mutations. *Clin Genet* 2004;65:226-32.
 - 5 **Plenge RM**, Stevenson RA, Lubs HA, Schwartz CE, Willard HF. Skewed X-chromosome inactivation is a common feature of X-linked mental retardation disorders. *Am J Hum Genet* 2002;71:168-73.
 - 6 **Allen RC**, Zoghbi HY, Moseley AB, Rosenblatt HM, Belmont JW. Methylation of HpaII and HhaI sites near the polymorphic CAG repeat in the human androgen-receptor gene correlates with X chromosome inactivation. *Am J Hum Genet* 1992;51:1229-39.
 - 7 **Carrel L**, Willard HF. An assay for X inactivation based on differential methylation at the fragile X locus, FMR1. *Am J Med Genet* 1996;64:27-30.
 - 8 **van Kamp H**, Jansen R, Willemze R, Fibbe WE, Landegent JE. Studies on clonality by PCR analysis of the PGK-1 gene. *Nucleic Acids Res* 1991;19:2794.
 - 9 **Thouin MM**, Giron JM, Hoffman EP. Molecular analysis of genetic identity. In: Dracopoli NC, Haines JL, Korf BR, et al, editors. *Current protocols in human genetics*, vol 2. New York: John Wiley and Sons, 2002:9.7.1-6.
 - 10 **Baumstark A**, Lower KM, Sinkus A, Andriuskeviciute I, Jurkeniene L, Gecz J, Just W. Novel PHF6 mutation p.D333del causes Borjeson-Forssman-Lehmann syndrome. *J Med Genet* 2003;40:e50.
 - 11 **Vallee D**, Chevrier E, Graham GE, Lazzaro MA, Lavigne PA, Hunter AG, Picketts DJ. A novel PHF6 mutation results in enhanced exon skipping and mild Borjeson-Forssman-Lehmann syndrome. *J Med Genet* 2004;41:778-83.
 - 12 **Paganini F**, Baralle FE. Genomic variants in exons and introns: identifying the splicing spoilers. *Nat Rev Genet* 2004;5:389-96.
 - 13 **Cooper DN**, Youssoufian H. The CpG dinucleotide and human genetic disease. *Hum Genet* 1988;78:151-5.
 - 14 **Visootsak J**, Rosner B, Dykens E, Schwartz C, Hahn K, White SM, Szeftel R, Graham JM. Clinical and behavioral features of patients with Borjeson-Forssman-Lehmann syndrome with mutations in PHF6. *J Pediatr* 2004;145:819-25.
 - 15 **Braunschweig D**, Simcox T, Samaco RC, LaSalle JM. X-Chromosome inactivation ratios affect wild-type MeCP2 expression within mosaic Rett syndrome and MeCP2-/- mouse brain. *Hum Mol Genet* 2004;13:1275-86.
 - 16 **Weaving LS**, Ellaway CJ, Gecz J, Christodoulou J. Rett syndrome: clinical review and genetic update. *J Med Genet* 2005;42:1-7.
 - 17 **Matsuo K**, Murano I, Kajii T. Borjeson-Forssman-Lehmann syndrome in a girl. *Jinruidengaku Zasshi* 1984;29:121-6.
 - 18 **Petridou M**, Kimiskidis V, Deligiannis K, Kazis A. Borjeson-Forssman-Lehmann syndrome: two severely handicapped females in a family. *Clin Neurol Neurosurg*, 1997;99:148-50.
 - 19 **Kubota T**, Oga S, Ohashi H, Iwamoto Y, Fukushima Y. Borjeson-Forssman-Lehmann syndrome in a woman with skewed X-chromosome inactivation. *Am J Med Genet* 1999;87:258-61.

CORRECTION

doi: 10.1136/jmg.2005.031369corr1

In the Original article titled, Oculocerebrocutaneous syndrome: the brain malformation defines a core phenotype (*J Med Genet* 2005;42:913-921) the supplementary figures were missing from the paper. The supplementary figures are available on the JMG website at <http://www.jmedgenet.com/supplemental>.