



Figure 2 MLPA copy number analysis. The normalised probe intensities for (A) the proband with the *SDHB* gene deletion and (B) a healthy person without the deletion are shown.

phaeochromocytoma and neuroblastoma. *Br J Cancer* 2004;**91**:1835–41.

3. **Grau E**, Oltra S, Orellana C, Hernandez-Marti M, Castel V, Martinez F. There is no evidence that the *SDHB* gene is involved in neuroblastoma development. *Oncol Res* 2005;**15**:393–8.

4. **Tischler AS**, DeLellis RA, Biales B, Nunnemacher G, Carabba V, Wolfe HJ. Nerve growth factor-induced neurite outgrowth from normal human chromaffin cells. *Lab Invest* 1980;**43**:399–409.

5. **Schlisio S**, Kenchappa RS, Vredevelde LC, George RE, Stewart R, Greulich H, Shahriari K, Nguyen NV, Piggy P, Dahia PL, Pomeroy SL, Maris JM, Look AT, Meyerson M, Peeper DS, Carter BD, Kaelin WG Jr. The kinesin KIF1B{beta} acts downstream from EglN3 to induce apoptosis and is a potential 1p36 tumor suppressor. *Genes Dev* 2008;**22**:884–93.

CORRECTION

doi:10.1136/jmg.2008.059568corr1

In the mutation report by Russell-Swetek *et al* that was published in the September 2008 issue of the journal (Russell-Swetek A, West AN, Minturn JE, Jenkins J, Rodriguez-Galindo C, Ribeiro R, Zambetti GP. Identification of a novel *TP53* germline mutation E285V in a rare case of paediatric adrenocortical carcinoma and choroid plexus carcinoma. *J Med Genet* 2008;**45**:603–6), the spelling of one of the authors' names was incorrect: J E Mintern should be J E Minturn. The journal apologises for this error.