



OPEN ACCESS

ORIGINAL ARTICLE

Intermediate C9orf72 alleles in neurological disorders: does size really matter?

Adeline S L Ng,¹ Eng-King Tan^{2,3}

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/jmedgenet-2017-104752>).

¹Department of Neurology, National Neuroscience Institute, Tan Tock Seng Hospital, Singapore

²Department of Neurology, National Neuroscience Institute, Singapore General Hospital, Singapore

³Neuroscience and Behavioural Disorders, Duke-NUS Graduate Medical School, Singapore

Correspondence to

Dr Eng-King Tan, Department of Neurology, Singapore General Hospital, Block 1 Level 3, Outram Road, Singapore 169108, Singapore; gnrtek@sg.h.com.sg

Received 21 April 2017

Revised 25 May 2017

Accepted 26 May 2017

Published Online First

8 July 2017

ABSTRACT

C9orf72 repeat expansions is a major cause of familial frontotemporal dementia (FTD) and amyotrophic lateral sclerosis (ALS) worldwide. Sizes of <20 hexanucleotide repeats are observed in controls, while up to thousands associate with disease. Intermediate C9orf72 repeat lengths, however, remain uncertain. We systematically reviewed the role of intermediate C9orf72 alleles in C9orf72-related neurological disorders. We identified 49 studies with adequate available data on normal or intermediate C9orf72 repeat length, involving subjects with FTD, ALS, Parkinson's disease (PD), atypical parkinsonism, Alzheimer's disease (AD) and other aetiologies. We found that, overall, normal or intermediate C9orf72 repeat lengths are not associated with higher disease risk across these disorders, but intermediate allele sizes appear to associate more frequently with neuropsychiatric phenotypes. Intermediate sizes were detected in subjects with personal or family history of FTD and/or psychiatric illness, parkinsonism complicated by psychosis and rarely in psychiatric cohorts. Length of the hexanucleotide repeat may be influenced by ethnicity (with Asian controls displaying shorter normal repeat lengths compared with Caucasians) and underlying haplotype, with more patients and controls carrying the 'risk' haplotype rs3849942 displaying intermediate alleles. There is some evidence that intermediate alleles display increased methylation levels and affect normal transcriptional activity of the C9orf72 promoter, but the 'critical' repeat size required for initiation of neurodegeneration remains unknown and requires further study. In common neurological diseases, intermediate C9orf72 repeats do not influence disease risk but may associate with higher frequency of neuropsychiatric symptoms. This has important clinical relevance as intermediate carriers pose a challenge for genetic counselling.

INTRODUCTION

The hexanucleotide repeat GGGGCC expansion within the first intron of the C9orf72 gene on chromosome 9p21 has been associated with amyotrophic lateral sclerosis (ALS) and frontotemporal dementia (FTD),^{1,2} accounting for ~40% of familial ALS, 30% of familial FTD and ~8% of sporadic ALS cases in predominantly Caucasian populations.³ Most healthy individuals carry up to 25 repeats, with more than half carrying only two repeats, while pathogenic expansions of hundreds² to several thousands have been reported in FTD and ALS.^{1,4} Other repeat expansion disorders have

explored the potential effect intermediate alleles have on conferring disease: in Huntington's disease (HD), intermediate CAG lengths overlapped with patients with HD on behavioural measures⁵ and pathological evidence of HD was found in patients with intermediate (range 27–35) CAG repeats.⁵ Spinocerebellar ataxia 2 patients with apparently sporadic disease have shown intermediate CAG repeats of 22–35 units in the ATXN2 locus.⁶

In C9orf72-related disease, a repeat length of >30 has typically been defined as pathogenic,² mainly due to technical limitations of the repeat-primed PCR (rp-PCR) technique. While cell and Drosophila models indicate that expression of >30 repeats are sufficient to cause neurodegeneration,⁷ it remains unknown how many repeats are truly needed to cause disease. Small repeat sizes of 20–22 were reportedly pathogenic in FTD,⁸ and the clinical phenotype of ALS cases with 20–30 repeats appear similar to those with >30 repeats.⁹ Given the conflicting evidence concerning pathogenicity of intermediate repeat numbers, we sought to summarise the current literature and provide insights into the potential role intermediate C9orf72 repeats may play (if any) in modulating clinical disease.

Review of the current literature

We conducted a PubMed and Google Scholar search for all relevant full-text studies and articles using search terms including 'C9orf72', 'intermediate alleles' and 'repeat size' in neurological diseases. Only English articles with available data (including online supplementary material) on C9orf72 allele lengths in normal and intermediate repeat ranges were included. Methods used by each study for repeat-length analysis (Southern blotting, rp-PCR and fluorescent fragment length analysis) are listed in the online supplementary table files.

Intermediate C9orf72 repeats do not appear to influence disease risk or clinical characteristics in non-expansion cases and controls

We identified 49 studies on intermediate alleles in neurodegenerative diseases, including 13 studies with FTD-spectrum cases, and 16 studies with ALS cases. Eight studies found no significant difference in mean repeat lengths between expansion-negative FTD cases and controls.^{10–17} Sporadic FTD cases appear to show similar allele frequency distribution as controls, with the two-unit repeat being the most common, followed by the 5-repeat, 6-repeat, 7-repeat or 8-repeat unit.^{12–19} Repeat range was higher in a Dutch study of 363 FTD patients where



CrossMark

To cite: Ng ASL, Tan E-K. *J Med Genet* 2017;**54**:591–597.

10% had an FTD-ALS syndrome, with FTD cases showing mean (SD) size of 13.9 ± 14.0 repeats (range 1–64) compared with 9.1 ± 6.8 (range 2–35) in controls. Notably, a proband in an FTD-ALS family carried 26 repeats, with affected family members carrying a repeat range of 8–29.²⁰ Pathologically confirmed corticobasal degeneration (n=18) and progressive supranuclear palsy (PSP) (n=177) cases carried repeat lengths ranging from 2 to 22, with one clinically diagnosed ‘typical’ PSP case with family history of dementia and parkinsonism carrying 27 repeats.²¹ In sporadic ALS, repeat length and allele frequency distribution appear to be similar between cases and controls.^{12 13 22–28} This was similarly seen in familial ALS,^{26 28 29} but a Japanese study reported slightly longer repeat lengths in familial ALS non-expansion samples (5.90 ± 3.91 repeats) carrying the risk ‘A’-allele compared with those without (3.36 ± 2.13 repeats).²⁷ In sporadic ALS, no correlation between lengths of the shortest, longest and sum of both alleles without the repeat expansion with age of onset, survival or region of disease onset was seen.²⁴

Nineteen studies of PD and atypical parkinsonism cases were identified. Nine PD studies reported range of repeat lengths similar between PD cases and controls,^{14 16 17 30–35} with no correlation between repeat length and family history or age of disease onset.³⁵ No evidence of association between repeat length and age-at-onset or risk of PD, essential tremor (ET) or restless legs syndrome after correction for multiple testing ($p < 0.0028$) was seen,³¹ and the length of the longest non-expanded repeat allele did not correlate with age at onset ($p = 0.11$) in PD.³² None of 29 LRRK2 mutation carriers screened for C9orf72 had an expanded or intermediate allele (<11 repeats).¹⁶ Twenty-three cases of classical PD harbouring larger-than-normal alleles (21–38 repeats) have been reported,^{14 16 34 36} including two with severe dementia¹⁶ and one young-onset ‘typical’ PD,¹⁴ suggesting that C9orf72 intermediate repeats may contribute to risk for PD and ET plus parkinsonism.³⁴ However, intermediate allele frequency appeared to be similar to that of controls (2.2% of PD vs 2.9% in controls).³⁰ Using a lower cut-off of <20 for intermediate repeat length, Nuytemans *et al* reported more intermediate alleles in PD (2%) compared with controls (0.3%), with intermediate repeats significantly associating with increased PD risk ($p = 0.008$) at an OR of 9.6, but with an extremely wide CI due to rarity of the intermediate repeat. Thirteen intermediate repeat patients demonstrated a classical PD phenotype responsive to levodopa, with no suggestion of ALS or family history of ALS.³⁴ Overall, the small number of cases reported and the occurrence of intermediate repeats in some controls suggests that C9orf72 intermediate repeats are most likely a susceptibility factor or arise from phenotypic heterogeneity, rather than causal for PD.³⁴ This was confirmed when no intermediate or expanded repeats was found in an autopsy-confirmed PD data set.³⁷ Overall, only a small number of large expansions (>30 repeats) have been found in PD, suggesting that >30 repeats are not a common cause of PD (0.2% of 3500 tested).³⁷ Finally, a meta-analysis by the Genetic Epidemiology of Parkinson’s disease (GEO-PD) consortium cohort of 7494 patients with PD revealed a small increase in PD risk with an increasing number of C9orf72 repeats, but no robust association was detected. The cohort size suggested an association for the 10-unit repeat and for pooled alleles of ≥ 17 repeats in increasing PD risk, but this did not reach significance after correction for multiple testing.³⁸

In nine atypical parkinsonian disease studies, including pathologically confirmed multiple system atrophy (MSA)^{21 39} and diffuse Lewy body dementia (DLB)^{40 41} cases, pathogenic expansions were rare, and repeat range of non-expansion alleles were similar to that of controls.^{14 17 21 30 33 39 42} Two pathologically diagnosed DLB samples

carried 32 repeats and one with 33 repeats, although no details were provided regarding presence or absence of TDP43 pathology in these patients.⁴¹ Intermediate repeats (20–29 units) were significantly more frequent in atypical parkinsonism cases than controls ($p < 0.034$) and were seen in four (4.3%) female patients (with 20, 22, 23 and 28 repeats), three of whom had non-classical atypical parkinsonism.⁴² PSP patients with intermediate repeat lengths of 26 and 30 were reported, but it is unclear if the repeat expansions and clinical symptoms are related.³³ Five studies of Alzheimer’s disease (AD), including one study with 80% of subjects under age 65,⁴³ were included.^{16 32 43–45} Within the normal and intermediate repeat range, higher repeat numbers did not increase risk for AD,^{16 32} or interact with apolipoprotein E (APOE) genotype.⁴⁴ No significant difference in the frequency of short (<7) and intermediate (>7) repeats was seen between AD and mild cognitive impairment (MCI) subjects and controls; MCI cases with intermediate repeats were not at greater risk for AD,⁴⁵ and intermediate allele sizes had no influence on cerebrospinal fluid core AD biomarkers.⁴⁵

Intermediate C9orf72 alleles may be associated with a higher frequency of psychiatric symptoms

Four FTD families were reported to carry 20-repeat or 22-repeat alleles associated with the surrogate founder haplotype, segregating consistently in all affected siblings, with unaffected siblings carrying wild-type alleles (two to nine repeats).⁸ Most of the nine carriers with >20 to >30 repeats had extended periods of psychiatric symptoms and subjective cognitive complaints before clear neurological deterioration. In other studies, a 29-repeat allele was found in a patient from a Dutch family with C9orf72-related FTD-ALS,²⁰ and four patients with limb-onset ALS with intermediate repeats (range 20–22) had lower mean (SD) age of diagnosis compared with patients with <20 repeats (47.6 ± 15.9 vs 62.8 ± 11.2 years).⁹ Both patients in the latter study with 22 repeats exhibited cognitive and behavioural impairment similar to larger expansions and had family history of FTD, while two patients with 20 and 21 repeats had family history of dementia and psychiatric illness. In atypical PD or PD with psychosis, intermediate expansions (20–29) were detected in three female cases with atypical parkinsonian syndromes (severe rigid akinetic parkinsonism) and neuropsychiatric symptoms including schizoaffective psychosis and an FTD-like dementia.⁴² Unfortunately, these cases did not have pathological diagnosis. In schizophrenia, the pathogenic repeat expansion was detected in only two patients (0.67%), but the estimated number of repeats in controls appeared lower (2–5 units) than in schizophrenic patients without the expansion (2–30 units).⁴⁶ Only 4/130 (3%) patients with psychosis from the Northern Finland Birth Cohort had intermediate repeat sizes (17–26 units).⁴⁷ An Irish psychosis cohort found seven samples with >22 repeats including two schizophrenia cases (with 28 and 27 repeats) and five controls (with 23–26 repeats), but with no overall evidence of association between repeat length and schizophrenia.⁴⁸ Unfortunately, detailed phenotypic description of these intermediate carriers remains of variable quality across studies, both for motor and non-motor signs. Overall, it remains possible that the pathological cut-off for C9orf72 expansions is disease dependent, modulated by unknown or individual genetic factors, with intermediate allele sizes potentially predisposing to neuropsychiatric sequelae.

Repeat lengths in the normal and intermediate range may be haplotype dependent

The full background 82-SNP ‘founder’ haplotype (r) from which the C9orf72 expansion arose is present in almost all




Figure 1 Possible mechanistic pathways involving intermediate C9orf72 alleles.

populations studied in the ‘1000 Genome’ database (~15% of European controls).⁴⁹ Nearly 50% of individuals with non-(r) haplotypes carry up to two copies compared with only 5% of those with the (r) haplotype, who carry eight repeats or more. In Irish psychosis patients, 78% of samples with >6 repeats carried the founder haplotype compared with only 3% with <6 repeats.⁴⁸ A meta-analysis of genome-wide association study data from five European populations reduced the ‘r’ haplotype to a common 20-SNP ‘risk’ haplotype (rs3849942, allele ‘A’) in significant association with FTD and ALS,⁵⁰ and controls have shown significantly longer median repeat length (eight units) on the ‘risk’ haplotype compared with the wild-type haplotype (rs3849942 allele ‘G’; 2 units).¹ In PD, the rs3849942 allele was confirmed in 100% of all intermediate carriers, with the tagging (T) allele seen in 95% of all individuals with ≥ 8 repeats, versus 10% in those with <8 repeats.³⁴ Pathologically confirmed PD cases identified the T-allele in 40% of patients, and >90% of T-allele carriers carried >8 repeats.³⁷ Japanese ALS patients carrying at least one rs3849942 allele had significantly longer repeats than those without.²⁷

Possible mechanistic pathways involving intermediate alleles

Intermediate alleles are ‘predisposing’ alleles that lead to large expansions over many subsequent generations

C9orf72 alleles at <20 repeats appear stable between generations, while small expansions (20–150 repeats) are susceptible to unfaithful inheritance⁴ or somatic instability (figure 1).²⁰ Intermediate repeats may be prone to gaining and losing copies as opposed to longer repeats whose length mostly increases.³⁴ In French control samples, the largest repeat (22 units) changed size twice in the same family with all intergenerational changes occurring from a starting length of >10 repeats on the same rs3849942A ‘risk’ haplotype, suggesting that either repeat length

and/or risk haplotype confer instability.⁴ Increased frequency of short indels in the GC-rich low complexity sequence adjacent to the expanded repeat in C9orf72 carriers suggests that subsequent pathological expansion may be due to replication slippage.⁵¹ However, the younger generation in 20 families of index patients with intermediate alleles of up to 22 units did not show longer intermediate repeat lengths or pathological expansions, suggesting that intermediate length might be ‘predisposing’ rather than ‘pre-mutation’ alleles for further stepwise expansion over many generations.⁵¹ Large-scale multigenerational cohorts will be required to determine how many repeats constitute such an allele.

Intermediate alleles may display increased methylation but at lower levels than large expansions

Hypermethylation of the CpG island 5’ to the C9orf72 GGGGCC repeat has been associated with expansion in ALS, but methylation changes have not been detected in either normal or intermediate alleles (up to 43 repeats), which questioned the cut-off of 30 repeats for pathogenicity.⁵² Comparison of methylation levels of homozygous normal intermediate repeat carriers (I/I) (7–24 units) with homozygous normal short repeat carriers (S/S) (2–6 units) in ALS and FTD-ALS demonstrated a slight but significant increase in methylation from I/I samples compared with S/S samples in both patient and control groups, with higher methylation levels in I/I carriers derived mostly from carriers with at least one allele of >8 units. Intermediate wild-type alleles of expansion carriers were also significantly more methylated than normal short wild-type alleles ($p < 0.0001$). Notably, 5’ CpG and GGGGCC methylation was significantly higher in brain than in the blood of patients with normal repeats ($p < 0.0001$).⁵³ Given that the degree of methylation remained very low (<5%) in samples without long expansions versus ~10% in expansion

Table 1 Unanswered questions pertaining to intermediate C9orf72 alleles

Minimum C9orf72 repeat length required for neurodegeneration remains unknown	<ul style="list-style-type: none"> ▶ Repeat lengths from ≤15 to 27 in the blood have not shown somatic instability in limited studies.^{56,57} ▶ TDP43 formation may be 'age-related' findings; pathological studies found no expansion or intermediate alleles (20–29 repeats) among LRRK2 G2019S carriers and AD cases with concomitant TDP43-positive inclusions.¹⁶
Variation of large-normal and intermediate C9orf72 repeat lengths by ethnicity needs to be determined	<ul style="list-style-type: none"> ▶ European haplotype exists in Asian cohorts,^{25,27,49} but at a lower frequency (<2% in Chinese controls compared with 9% in European controls).²⁵ ▶ European founder haplotype seen in 15% of Europeans compared with only 0.5% in Asians.⁴⁹ ▶ Asian controls appear to carry smaller normal repeat lengths (7–14) compared with Caucasian controls (0–32).³⁸ ▶ Chinese samples with >12 repeats are rarely observed.²⁵
Effect of homozygosity versus heterozygosity of normal-length and intermediate-length C9orf72 alleles on disease susceptibility	<ul style="list-style-type: none"> ▶ Higher methylation levels seen in homozygous intermediate allele carriers compared with homozygous short allele carriers.⁵³ ▶ No variability in regions flanking the GGGGCC repeat found in pathogenic expansion carriers with second allele repeat size within normal range (2–11).¹⁶ ▶ Homozygous intermediate repeat carriers reported more frequently in FTD compared with controls (6.1% vs 4.6%).⁵¹

Unanswered questions pertaining to intermediate C9orf72 alleles.

FTD, frontotemporal dementia; LRRK2, leucine-rich repeat kinase; TDP43, TAR DNA-binding protein 43.

carriers, further work needs to be done to elucidate any epigenetic modulation in intermediate C9orf72 allele carriers and the minimum repeat cut-off required for this to occur.

Intermediate alleles affect normal transcriptional activity of the C9orf72 promoter

In vitro reporter gene expression studies have shown significantly decreased transcriptional activity of the C9orf72 promoter with increasing number of normal and intermediate repeats, with a 24-unit containing promoter showing >50% reduction in transcription compared with 2 units on the wild-type allele.⁵¹ In human kidney and neuroblastoma cell lines (HEK293T cells), intermediate repeats (7–24) have shown significantly decreased C9orf72 promoter activity compared with normal short repeats (2–6 units),⁵³ with lower transcriptional activity appearing more prominent in the presence of small deletions flanking the GGGGCC unit. Overall, these findings suggest that increasing repeat length can lead to decreased promoter activity via increased methylation of CpG sequences in larger repeats, with transcriptional silencing of the promoter.⁵³

UNANSWERED QUESTIONS

Minimum repeat length in blood corresponding with large expansions in CNS tissue

The minimum repeat length required for disease is not known due to somatic instability, with variation in length between tissues, even within the same individual, complicating genotype-phenotype studies (table 1).⁵⁴ Subtle differences in repeat length within the same patient and between monozygotic twins strongly suggest that random, stochastic expansion events may occur during cell division, resulting in somatic and/or germline mosaicism contributing to intraindividual and interindividual repeat variation.⁵⁵ Limited data show that repeat lengths from ≤15 to 27 in the blood do not show somatic instability.^{56,57} Postmortem studies involving a wide range of CNS tissue examination in cases with large-normal and intermediate repeats lengths in peripheral blood are needed.

Variation of large-normal and intermediate repeat lengths by ethnicity

In the GEO-PD consortium, Asian controls appeared to carry smaller normal repeat lengths (7–14) compared with Caucasian controls (0–32).³⁸ Caucasian alleles frequently contain up to 43 repeats,^{4,52} but in Chinese samples, >12 repeats are rarely observed.²⁵ The European haplotype does exist in Asian cohorts^{25,27,49} but at a lower frequency (<2% in Chinese controls

compared with 9% in European controls).²⁵ Data from the 1000-genomes project estimate the frequency of the European founder haplotype in 15% of Europeans compared with only 0.5% in Asians.⁴⁹ Two Han Chinese ALS expansion carriers had methylation of the upstream CpG island but did not carry the founder haplotype,²⁵ implying that C9orf72 alleles may undergo dynamic mutation independent of the European haplotype. This 'Chinese haplotype' was also associated with ≥8 repeats (20% vs 8% in other haplotypes), implying that both the Scandinavian and Chinese haplotypes may confer repeat length instability.²⁵ It remains possible that the haplotype may be genetic background for expansion carriers and not necessarily associated with the expansion, as none of 53 FTD patients recruited in the USA with diverse ethnic backgrounds shared the exact same Scandinavian risk haplotype.¹¹ Multicentre collaborative efforts pooling Asian samples are needed to determine if the lower incidence of pathogenic C9orf72 expansions in multiple Asian cohorts is due to shorter Asian repeat lengths that may be at lower risk for instability.

'Critical' repeat size required for initiation of neurodegeneration remains unknown

Toxic gain-of-function by dipeptide repeat proteins (DPRs) and/or RNA foci generated by unconventional repeat-associated non-ATG translation and TDP43 inclusions remains a possible mechanism for C9orf72 pathogenesis.³ Cognitively normal cases from the Queen Square Brain Bank screened for C9orf72 expansions found one 84 year-old subject with intermediate repeats (20–40) showing TDP43-positive lesions, sense and antisense RNA foci in the frontal cortex and p62-positive inclusions containing all five DPRs, but more sparsely compared with FTD cases with large expansions.⁵⁸ DPR inclusions and RNA foci were, however, notably absent in two pathologically normal cases with 20 repeats, suggesting it may not be long enough to trigger the entire disease cascade necessary for TDP43 pathology. TDP43 formation may also represent 'age-related' phenomena, and pathological studies have found no expansion or intermediate alleles (20–29 repeats) among LRRK2 G2019S carriers and AD cases who had concomitant TDP43-positive inclusions.¹⁶

Effect of homozygosity versus heterozygosity of normal and intermediate alleles

In pathogenic expansion carriers, repeat size of the second allele appear to be within the normal range (2–11) and unlikely to contribute to disease given that no variability in the regions flanking the hexanucleotide repeat was found.¹⁶ Homozygous

Table 2 Priorities for research into C9ORF72 intermediate alleles

Address methodological differences	Repeat-primed PCR should be combined with Southern blot for estimation of repeat size and not be used in isolation. Sequencing should be used for smaller repeat sizes.
Deciphering role of DNA composition in expanded repeats	The 'critical repeat size' may not be key, but rather the exact composition within the expanded repeats, including the possibility of repeat interruptions.
Developing novel genotyping methods	Developing the ability to conduct a fast comprehensive systematic screen for repeat expansions on a genome-wide level.
Adopting a multigeneration intrafamilial approach	Phenotype–genotype correlations within large multigeneration families can minimise the effect of heterogeneity in genetic background and confounders.
Detailed clinical phenotyping	Detailed clinical and neuropsychological profiling in symptomatic and asymptomatic carriers and healthy controls for accurate genotype–phenotype correlations.
Developing better cellular models of C9orf72	Patient-derived induced pluripotent stem cell neurons harbouring different repeat sizes and transgenic non-human primates can provide additional insights.
Longitudinal cohort study	A longitudinal cohort including subjects with intermediate alleles with long-term follow-up supported by detailed clinical, neuroimaging and biochemical evaluation.
Multicentre study approach for optimising sample size	A global multicentre approach with standardised protocols and a centralised genotyping laboratory will aid in identifying rare associations, elucidate variation according to ethnicity and minimise diagnostic errors.

intermediate repeat carriers have been reported more frequently in FTD subjects compared with controls (6.1% vs 4.6%),⁵¹ but in PD cases with >20 repeats, all but two of 407 intermediate repeat carriers were heterozygous for the intermediate allele and a low copy number allele, with only one sample *homozygous* for intermediate repeats. This proband carried three different maximum intermediate repeat alleles in two offspring and herself, developing ET at age 20 with mild parkinsonism at 97 years old.³⁴ As described earlier, higher methylation levels were seen in homozygous intermediate allele compared with homozygous short allele carriers.⁵³

FUTURE DIRECTIONS AND CONCLUSIONS

Further steps for elucidating the uncertain role(s) of intermediate C9orf72 hexanucleotide repeats include (table 2): (1) common centralised genotyping methodology for comparing allele size accurately. The common rp-PCR method cannot reliably distinguish between repeat sizes >30–60 units² and is inadequate when used without confirmation by Southern blot, as confirmed by a masked study at 14 laboratories showing lack of accuracy when rp-PCR was used in isolation.⁵⁹ Southern blot analysis, however, is affected by age and somatic heterogeneity, hence for smaller range repeats, sequencing to determine the actual size may be more useful and will also provide the actual DNA composition of the repeat; additionally, Southern blot results should be interpreted in conjunction with recent improved rp-PCR techniques⁶⁰; (2) deciphering DNA composition within expanded repeats will be important, as the 'critical repeat size' may not be key, but rather the exact composition within the expanded repeats, including the presence or absence of stabilising interruptions; (3) developing novel genotyping methods for elucidating the role of other GGGGCC loci and repeat expansions in the human genome as well as other pathogenic genes in modulating the risk of C9orf72-linked disorders; (4) developing a fast comprehensive systematic screen for repeat expansions on a genome-wide level; (5) development of novel cellular and in vivo models given that current ones do not fully recapitulate clinical disease. The wide range of pathogenic GGGGCC repeat sizes and the large size of the repeat region, combined with the fact that it is entirely GC makes the generation of cell models technically difficult. Patient-derived induced pluripotent stem cell neurons harbouring different repeat sizes and transgenic non-human primates can provide additional insights.

Many studies involved retrospectively collected samples or were based on limited clinical information, rendering clinical

phenotyping of intermediate allele carriers less accurate. Efforts to ensure detailed clinical and neuropsychological profiling in symptomatic and asymptomatic carriers and healthy controls remains vital; accurate phenotype–genotype correlations within large multigeneration families can also aid in minimising the effect of confounders and genetic background heterogeneity. Finally, in genetic association studies, sample size remains critical. Thus, a global multicentre approach with standardised clinical and neuroimaging protocols and a centralised genotyping laboratory will aid in identifying rare associations and minimise diagnostic errors, as well as investigate differences according to ethnicity, potentially unravelling new clues concerning the role of intermediate C9orf72 alleles.

Contributors A SL Ng conducted the literature search and drafting of the manuscript. EK Tan revised the manuscript for intellectual content and provided the outline for the draft. A SL Ng and EK Tan drafted and approved the manuscript.

Funding A SL Ng is supported by the Singapore Ministry of Health's National Medical Research Council (NIG grant) and the SingHealth Foundation Award (PRISM grant). EK Tan is supported by the Singapore Ministry of Health's National Medical Research Council (STaR and Parkinson's Disease TCR grants).

Competing interests None declared.

Provenance and peer review Commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- DeJesus-Hernandez M, Mackenzie IR, Boeve BF, Boxer AL, Baker M, Rutherford NJ, Nicholson AM, Finch NA, Flynn H, Adamson J, Kouri N, Wojtas A, Sengdy P, Hsiung GY, Karydas A, Seeley WW, Josephs KA, Coppola G, Geschwind DH, Wszolek ZK, Feldman H, Knopman DS, Petersen RC, Miller BL, Dickson DW, Boylan KB, Graff-Radford NR, Rademakers R. Expanded GGGGCC hexanucleotide repeat in noncoding region of C9ORF72 causes chromosome 9p-linked FTD and ALS. *Neuron* 2011;72:245–56.
- Renton AE, Majounie E, Waite A, Simón-Sánchez J, Rollinson S, Gibbs JR, Schymick JC, Laaksoviirta H, van Swieten JC, Myllykangas L, Kalimo H, Paetau A, Abramzon Y, Remes AM, Kaganovich A, Scholz SW, Duckworth J, Ding J, Harmer DW, Hernandez DG, Johnson JO, Mok K, Rytten M, Trabzuni D, Guerreiro RJ, Orrell RW, Neal J, Murray A, Pearson J, Jansen IE, Sondervan D, Seelaar H, Blake D, Young K, Halliwell N, Callister JB, Toulson G, Richardson A, Gerhard A, Snowden J, Mann D, Neary D, Nalls MA, Peuralinna T, Jansson L, Isoviita VM, Kaivorinne AL, Hölttä-Vuori M, Ikonen E, Sulkava R, Benatar M, Wu J, Chiò A, Restagno G, Borghero G, Sabatelli M, Heckerman D, Rogava E, Zinman L, Rothstein JD, Sendtner M, Drepper C, Eichler EE,

- Alkan C, Abdullaev Z, Pack SD, Dutra A, Pak E, Hardy J, Singleton A, Williams NM, Heutink P, Pickering-Brown S, Morris HR, Tienari PJ, Traynor BJ. ITALSGEN Consortium. A hexanucleotide repeat expansion in C9orf72 is the cause of chromosome 9p21-linked ALS-FTD. *Neuron* 2011;72:257–68.
- 3 Ng AS, Rademakers R, Miller BL. Frontotemporal dementia: a bridge between dementia and neuromuscular disease. *Ann NY Acad Sci* 2015;1338:71–93.
- 4 Beck J, Poulter M, Hensman D, Rohrer JD, Mahoney CJ, Adamson G, Campbell T, Uphill J, Borg A, Fratta P, Orrell RW, Malaspina A, Rowe J, Brown J, Hodges J, Sidle K, Polke JM, Houlden H, Schott JM, Fox NC, Rossor MN, Tabrizi SJ, Isaacs AM, Hardy J, Warren JD, Collinge J, Mead S. Large C9orf72 hexanucleotide repeat expansions are seen in multiple neurodegenerative syndromes and are more frequent than expected in the UK population. *Am J Hum Genet* 2013;92:345–53.
- 5 Ha AD, Jankovic J. Exploring the correlates of intermediate CAG repeats in Huntington disease. *Postgrad Med* 2011;123:116–21.
- 6 Kim JM, Hong S, Kim GP, Choi YJ, Kim YK, Park SS, Kim SE, Jeon BS. Importance of low-range CAG expansion and CAA interruption in SCA2 Parkinsonism. *Arch Neurol* 2007;64:1510–8.
- 7 Xu Z, Poidevin M, Li X, Li Y, Shu L, Nelson DL, Li H, Hales CM, Gearing M, Wingo TS, Jin P. Expanded GGGGCC repeat RNA associated with amyotrophic lateral sclerosis and frontotemporal dementia causes neurodegeneration. *Proc Natl Acad Sci U S A* 2013;110:7778–83.
- 8 Gómez-Tortosa E, Gallego J, Guerrero-López R, Marcos A, Gil-Neciga E, Sainz MJ, Díaz A, Franco-Macias E, Trujillo-Tiebas MJ, Ayuso C, Pérez-Pérez J. C9orf72 hexanucleotide expansions of 20–22 repeats are associated with frontotemporal deterioration. *Neurology* 2013;80:366–70.
- 9 Byrne S, Heverin M, Elamin M, Walsh C, Hardiman O. Intermediate repeat expansion length in C9orf72 may be pathological in amyotrophic lateral sclerosis. *Amyotroph Lateral Scler Frontotemporal Degener* 2014;15:148–50.
- 10 Dobson-Stone C, Hallupp M, Bartley L, Shepherd CE, Halliday GM, Schofield PR, Hodges JR, Kwok JB. C9orf72 repeat expansion in clinical and neuropathologic frontotemporal dementia cohorts. *Neurology* 2012;79:995–1001.
- 11 Ferrari R, Mok K, Moreno JH, Cosentino S, Goldman J, Pietrini P, Mayeux R, Tierney MC, Kapogiannis D, Jicha GA, Murrell JR, Ghetti B, Wassermann EM, Grafman J, Hardy J, Huey ED, Momeni P. Screening for C9orf72 repeat expansion in FTL. *Neurobiol Aging* 2012;33:1850.e1–1850.e11.
- 12 Itzcovich T, Xi Z, Martinetto H, Chrem-Méndez P, Russo MJ, de Ambrosi B, Uchitel OD, Nogués M, Silva E, Rojas G, Bagnatti P, Amengual A, Campos J, Rogava E, St George-Hyslop P, Allegri R, Seveler G, Surace EI. Analysis of C9orf72 in patients with frontotemporal dementia and amyotrophic lateral sclerosis from Argentina. *Neurobiol Aging* 2016;40:192.e13–192.e15.
- 13 Jiao B, Tang B, Liu X, Yan X, Zhou L, Yang Y, Wang J, Xia K, Shen L. Identification of C9orf72 repeat expansions in patients with amyotrophic lateral sclerosis and frontotemporal dementia in mainland China. *Neurobiol Aging* 2014;35:936.e19–936.e22.
- 14 Lin CH, Chen TF, Chiu MJ, Lin HI, Wu RM. Lack of c9orf72 repeat expansion in Taiwanese patients with mixed neurodegenerative disorders. *Front Neurol* 2014;5:59.
- 15 Rutherford NJ, Heckman MG, DeJesus-Hernandez M, Baker MC, Soto-Ortolaza AI, Rayaprolu S, Stewart H, Finger E, Volkening K, Seeley WW, Hatanpaa KJ, Lomen-Hoerth C, Kertesz A, Bigio EH, Lippa C, Knopman DS, Kretzschmar HA, Neumann M, Caselli RJ, White CL, Mackenzie IR, Petersen RC, Strong MJ, Miller BL, Boeve BF, Uitti RJ, Boylan KB, Wszolek ZK, Graff-Radford NR, Dickson DW, Ross OA, Rademakers R. Length of normal alleles of C9orf72 GGGGCC repeat do not influence disease phenotype. *Neurobiol Aging* 2012;33:2950.e5–2950.e7.
- 16 Xi Z, Zinman L, Grinberg Y, Moreno D, Sato C, Bilbao JM, Ghani M, Hernández I, Ruiz A, Boada M, Morón FJ, Lang AE, Marras C, Bruni A, Colao R, Maletta RG, Puccio G, Rainero I, Pinessi L, Galimberti D, Morrison KE, Moorby C, Stockton JD, Masellis M, Black SE, Hazrati LN, Liang Y, van Haersma de With J, Fornazzari L, Villagra R, Rojas-García R, Clarimón J, Mayeux R, Robertson J, St George-Hyslop P, Rogava E. Investigation of c9orf72 in 4 neurodegenerative disorders. *Arch Neurol* 2012;69:1583–90.
- 17 Yeh TH, Lai SC, Weng YH, Kuo HC, Wu-Chou YH, Huang CL, Chen RS, Chang HC, Traynor B, Lu CS. Screening for C9orf72 repeat expansions in parkinsonian syndromes. *Neurobiol Aging* 2013;34:1311.e3–1311.e4.
- 18 Ogaki K, Li Y, Takanashi M, Ishikawa K, Kobayashi T, Nonaka T, Hasegawa M, Kishi M, Yoshino H, Funayama M, Tsukamoto T, Shioya K, Yokochi M, Imai H, Sasaki R, Kokubo Y, Kuzuhara S, Motoi Y, Tomiyama H, Hattori N. Analyses of the MAPT, PGRN, and C9orf72 mutations in Japanese patients with FTL, PSP, and CBS. *Parkinsonism Relat Disord* 2013;19:15–20.
- 19 Tang M, Gu X, Wei J, Jiao B, Zhou L, Zhou Y, Weng L, Yan X, Tang B, Xu J, Shen L. Analyses MAPT, GRN, and C9orf72 mutations in Chinese patients with frontotemporal dementia. *Neurobiol Aging* 2016;46:235.e11–235.e15.
- 20 Simón-Sánchez J, Doppler EG, Cohn-Hokke PE, Hukema RK, Nicolaou N, Seelaar H, de Graaf JR, de Koning I, van Schoor NM, Deeg DJ, Smits M, Raaphorst J, van den Berg LH, Schelhaas HJ, De Die-Smulders CE, Majoor-Krakauer D, Rozemuller AJ, Willemsen R, Pijnenburg YA, Heutink P, van Swieten JC. The clinical and pathological phenotype of C9orf72 hexanucleotide repeat expansions. *Brain* 2012;135:723–35.
- 21 Schottlaender LV, Polke JM, Ling H, MacDoanld ND, Tucci A, Nanji T, Pittman A, de Silva R, Holton JL, Revesz T, Sweeney MG, Singleton AB, Lees AJ, Bhatia KP, Houlden H. The analysis of C9orf72 repeat expansions in a large series of clinically and pathologically diagnosed cases with atypical parkinsonism. *Neurobiol Aging* 2015;36:1221.e1–1221.e6.
- 22 Abramychewa NY, Lysogorskaia EV, Stepanova MS, Zakharova MN, Kovrazhkina EA, Razinskaya OD, Smirnov AP, Maltsev AV, Ustyugov AA, Kukharsky MS, Khritankova IV, Bachurin SO, Cooper-Knock J, Buchman VL, Ilarioshkin SN, Skvortsova VI, Ninkina N. C9orf72 hexanucleotide repeat expansion in ALS patients from the Central European Russia population. *Neurobiol Aging* 2015;36:2908.e5–2908.e9.
- 23 Byrne S, Elamin M, Bede P, Shatunov A, Walsh C, Corr B, Heverin M, Jordan N, Kenna K, Lynch C, McLaughlin RL, Iyer PM, O'Brien C, Phukan J, Wynne B, Bokde AL, Bradley DG, Pender N, Al-Chalabi A, Hardiman O. Cognitive and clinical characteristics of patients with amyotrophic lateral sclerosis carrying a C9orf72 repeat expansion: a population-based cohort study. *Lancet Neurol* 2012;11:232–40.
- 24 Debray S, Race V, Crabbé V, Herdewyn S, Matthijs G, Goris A, Dubois B, Thijs V, Robberecht W, Van Damme P. Frequency of C9orf72 repeat expansions in amyotrophic lateral sclerosis: a Belgian cohort study. *Neurobiol Aging* 2013;34:2890.e7–2890.e12.
- 25 He J, Tang L, Benyamin B, Shah S, Hemani G, Liu R, Ye S, Liu X, Ma Y, Zhang H, Cremin K, Leo P, Wray NR, Visscher PM, Xu H, Brown MA, Bartlett PF, Mangelsdorf M, Fan D. C9orf72 hexanucleotide repeat expansions in Chinese sporadic amyotrophic lateral sclerosis. *Neurobiol Aging* 2015;36:2660.e1–2660.e8.
- 26 Jang JH, Kwon MJ, Choi WJ, Oh KW, Koh SH, Ki CS, Kim SH. Analysis of the C9orf72 hexanucleotide repeat expansion in Korean patients with familial and sporadic amyotrophic lateral sclerosis. *Neurobiol Aging* 2013;34:1311.e7–1311.e9.
- 27 Konno T, Shiga A, Tsujino A, Sugai A, Kato T, Kanai K, Yokoseki A, Eguchi H, Kuwabara S, Nishizawa M, Takahashi H, Onodera O. Japanese amyotrophic lateral sclerosis patients with GGGGCC hexanucleotide repeat expansion in C9orf72. *J Neurol Neurosurg Psychiatry* 2013;84:398–401.
- 28 Mok KY, Koutsis G, Schottlaender LV, Polke J, Panas M, Houlden H. High frequency of the expanded C9orf72 hexanucleotide repeat in familial and sporadic Greek ALS patients. *Neurobiol Aging* 2012;33:1851.e1–1851.e5.
- 29 Ogaki K, Li Y, Atsuta N, Tomiyama H, Funayama M, Watanabe H, Nakamura R, Yoshino H, Yato S, Tamura A, Naito Y, Taniguchi A, Fujita K, Izumi Y, Kaji R, Hattori N, Sobue G. Japanese Consortium for Amyotrophic Lateral Sclerosis research (JaCALS). Analysis of C9orf72 repeat expansion in 563 Japanese patients with amyotrophic lateral sclerosis. *Neurobiol Aging* 2012;33:2527.e11–16.
- 30 Akimoto C, Forsgren L, Linder J, Birve A, Backlund I, Andersson J, Nilsson AC, Alstermark H, Andersen PM. No GGGGCC-hexanucleotide repeat expansion in C9orf72 in parkinsonism patients in Sweden. *Amyotroph Lateral Scler Frontotemporal Degener* 2013;14:26–9.
- 31 DeJesus-Hernandez M, Rayaprolu S, Soto-Ortolaza AI, Rutherford NJ, Heckman MG, Traynor S, Strongosky A, Graff-Radford N, Van Gerpen J, Uitti RJ, Shih JJ, Lin SC, Wszolek ZK, Rademakers R, Ross OA. Analysis of the C9orf72 repeat in Parkinson's disease, essential tremor and restless legs syndrome. *Parkinsonism Relat Disord* 2013;19:198–201.
- 32 Harms MB, Neumann D, Benitez BA, Cooper B, Carrell D, Racette BA, Perlmutter JS, Goate A, Cruchaga C. Parkinson disease is not associated with C9orf72 repeat expansions. *Neurobiol Aging* 2013;34:1519.e1–1519.e2.
- 33 Lesage S, Le Ber I, Condroyer C, Broussolle E, Gabelle A, Thobois S, Pasquier F, Mondon K, Dion PA, Rochefort D, Rouleau GA, Dürr A, Brice A. French Parkinson's Disease Genetics Study Group. C9orf72 repeat expansions are a rare genetic cause of parkinsonism. *Brain* 2013;136:385–91.
- 34 Nuytemans K, Bademci G, Kohli MM, Beecham GW, Wang L, Young JI, Nahab F, Martin ER, Gilbert JR, Benatar M, Haines JL, Scott WK, Züchner S, Pericak-Vance MA, Vance JM. C9orf72 intermediate repeat copies are a significant risk factor for Parkinson disease. *Ann Hum Genet* 2013;77:351–63.
- 35 Majoune E, Renton AE, Mok K, Doppler EG, Waite A, Rollinson S, Chiò A, Restagno G, Nicolaou N, Simon-Sanchez J, van Swieten JC, Abramzon Y, Johnson JO, Sendtner M, Pampallett R, Orrell RW, Mead S, Sidle KC, Houlden H, Rohrer JD, Morrison KE, Pall H, Talbot K, Ansoorge O, Hernandez DG, Arepalli S, Sabatelli M, Mora G, Corbo M, Giannini F, Calvo A, Englund E, Borghero G, Remes AM, Remes AM, Laaksovirta H, Van Derlin VM, Trojanowski JQ, Van Derlin VM, Lu CS, Nalls MA, Drory VE, Lu CS, Yeh TH, Le Ber I, Takahashi Y, Tsujii S, Le Ber I, Brice A, Drepper C, Williams N, Kirby J, Shaw P, Hardy J, Tienari PJ, Heutink P, Morris HR, Pickering-Brown S, Traynor BJ. Frequency of the C9orf72 hexanucleotide repeat expansion in patients with amyotrophic lateral sclerosis and frontotemporal dementia: a cross-sectional study. *Lancet Neurol* 2012;11:323–30.
- 36 Daoud H, Noreau A, Rochefort D, Paquin-Lanthier G, Gauthier MT, Provencher P, Pourcher E, Dupré N, Chouinard S, Jodoin N, Soland V, Fon EA, Dion PA, Rouleau GA. Investigation of C9orf72 repeat expansions in Parkinson's disease. *Neurobiol Aging* 2013;34:1710.e7–1710.e9.
- 37 Nuytemans K, Inchausti V, Beecham GW, Wang L, Dickson DW, Trojanowski JQ, Lee VM, Mash DC, Frosch MP, Foroud TM, Honig LS, Montine TJ, Dawson TM, Martin ER, Scott WK, Vance JM. Absence of C9orf72 expanded or intermediate repeats in autopsy-confirmed Parkinson's disease. *Mov Disord* 2014;29:827–30.
- 38 Theuns J, Verstraeten A, Sleegers K, Wauters E, Gijselink I, Smolders S, Crosiers D, Corsmit E, Elinck E, Sharma M, Krüger R, Lesage S, Brice A, Chung SJ, Kim MJ, Kim YJ, Ross OA, Wszolek ZK, Rogava E, Xi Z, Lang AE, Klein C, Weissbach A, Mellick GD,

- Silburn PA, Hadjigeorgiou GM, Dardiotis E, Hattori N, Ogaki K, Tan EK, Zhao Y, Aasly J, Valente EM, Petrucci S, Annesi G, Quattrone A, Ferrarese C, Brighina L, Deutschländer A, Puschmann A, Nilsson C, Garraux G, DeDoux MS, Pfeiffer RF, Boczarzka-Jedynak M, Opala G, Maraganore DM, Engelborghs S, De Deyn PP, Cras P, Cruts M, Van Broeckhoven C. GEO-PD Consortium. Global investigation and meta-analysis of the C9orf72 (G4C2)_n repeat in Parkinson disease. *Neurology* 2014;83:1906–13.
- 39 Scholz SW, Majounie E, Revesz T, Holton JL, Okun MS, Houlden H, Singleton AB. Multiple system atrophy is not caused by C9orf72 hexanucleotide repeat expansions. *Neurobiol Aging* 2015;36:1223.e1–1223.e2.
- 40 Geiger JT, Arthur KC, Dawson TM, Rosenthal LS, Pantelyat A, Albert M, Hillis AE, Crain B, Pletnikova O, Troncoso JC, Scholz SW. C9orf72 Hexanucleotide Repeat Analysis in Cases with Pathologically Confirmed Dementia with Lewy Bodies. *Neurodegener Dis* 2016;16:370–2.
- 41 Kun-Rodrigues C, Ross OA, Orme T, Shepherd C, Parkkinen L, Darwent L, Hernandez D, Ansoore O, Clark LN, Honig LS, Marder K, Lemstra A, Scheltens P, van der Flier W, Louwersheimer E, Holstege H, Rogava E, St George-Hyslop P, Londos E, Zetterberg H, Barber I, Braae A, Brown K, Morgan K, Maetzler W, Berg D, Troakes C, Al-Sarraj S, Lashley T, Holton J, Compta Y, Van Deerlin V, Trojanowski JQ, Serrano GE, Beach TG, Clarimon J, Lleó A, Morenas-Rodriguez E, Lesage S, Galasko D, Masliah E, Santana I, Diez M, Pastor P, Tienari PJ, Myllykangas L, Oinas M, Revesz T, Lees A, Boeve BF, Petersen RC, Ferman TJ, Escott-Price V, Graff-Radford N, Cairns NJ, Morris JC, Stone DJ, Pickering-Brown S, Mann D, Dickson DW, Halliday GM, Singleton A, Guerreiro R, Bras J. Analysis of C9orf72 repeat expansions in a large international cohort of dementia with Lewy bodies. *Neurobiol Aging* 2017;49:214.e13–214.e15.
- 42 Cannas A, Solla P, Borghero G, Floris GL, Chio A, Mascia MM, Modugno N, Muroli A, Orofino G, Di Stefano F, Calvo A, Moglia C, Restagno G, Meloni M, Farris R, Ciaccio D, Puddu R, Vacca MI, Melis R, Murrur MR, Tranquilli S, Corongiu D, Rolesu M, Cuccu S, Marrosu MG, Marrosu F. C9ORF72 intermediate repeat expansion in patients affected by atypical parkinsonian syndromes or Parkinson's disease complicated by psychosis or dementia in a Sardinian population. *J Neurol* 2015;262:2498–503.
- 43 Rollinson S, Halliwell N, Young K, Callister JB, Toulson G, Gibbons L, Davidson YS, Robinson AC, Gerhard A, Richardson A, Neary D, Snowden J, Mann DMA, Pickering-Brown SM. Analysis of the hexanucleotide repeat in C9ORF72 in Alzheimer's disease. *Neurobiol Aging* 2012;33:1846.e5–1846.e6.
- 44 Kohli MA, John-Williams K, Rajbhandary R, Naj A, Whitehead P, Hamilton K, Carney RM, Wright C, Crocco E, Gwirtzman HE, Lang R, Beecham G, Martin ER, Gilbert J, Benatar M, Small GW, Mash D, Byrd G, Haines JL, Pericak-Vance MA, Züchner S. Repeat expansions in the C9ORF72 gene contribute to Alzheimer's disease in Caucasians. *Neurobiol Aging* 2013;34:1519.e5–1519.e12.
- 45 Cacace R, Van Cauwenbergh C, Bettens K, Gijssels I, van der Zee J, Engelborghs S, Vandenbulcke M, Van Dongen J, Bäumer V, Dillen L, Mattheijssens M, Peeters K, Cruts M, Vandenbergh R, De Deyn PP, Van Broeckhoven C, Sleegers K. C9orf72 G4C2 repeat expansions in Alzheimer's disease and mild cognitive impairment. *Neurobiol Aging* 2013;34:1712.e1–1712.e7.
- 46 Galimberti D, Reif A, Dell'osso B, Kittel-Schneider S, Leonhard C, Herr A, Palazzo C, Villa C, Fenoglio C, Serpente M, Cioffi SMG, Prunas C, Paoli RA, Altamura AC. Scarpini E. C9ORF72 hexanucleotide repeat expansion is a rare cause of schizophrenia. *Neurobiol Aging* 2014.
- 47 Solje E, Miettunen J, Marttila R, Helisalmi S, Laitinen M, Koivumaa-Honkanen H, Isohanni M, Hiltunen M, Jääskeläinen E, Remes AM. The C9ORF72 expansion sizes in patients with psychosis: a population-based study on the Northern Finland Birth Cohort 1966. *Psychiatr Genet* 2016;26:92–4.
- 48 Fahey C, Byrne S, McLaughlin R, Kenna K, Shatunov A, Donohoe G, Gill M, Al-Chalabi A, Bradley DG, Hardiman O, Corvin AP, Morris DW. Analysis of the hexanucleotide repeat expansion and founder haplotype at C9ORF72 in an Irish psychosis case-control sample. *Neurobiol Aging* 2014;35:1510.e1–1510.e5.
- 49 Smith BN, Newhouse S, Shatunov A, Vance C, Topp S, Johnson L, Miller J, Lee Y, Troakes C, Scott KM, Jones A, Gray I, Wright J, Hortobágyi T, Al-Sarraj S, Rogelj B, Powell J, Lupton M, Lovestone S, Sapp PC, Weber M, Nestor PJ, Schelhaas HJ, Asbroek AA, Silani V, Gellera C, Taroni F, Ticozzi N, Van den Berg L, Veldink J, Van Damme P, Robberecht W, Shaw PJ, Kirby J, Pall H, Morrison KE, Morris A, de Belleroche J, Vianney de Jong JM, Baas F, Andersen PM, Landers J, Brown RH, Weale ME, Al-Chalabi A, Shaw CE. The C9ORF72 expansion mutation is a common cause of ALS+/-FTD in Europe and has a single founder. *Eur J Hum Genet* 2013;21:102–8.
- 50 Mok K, Traynor BJ, Schymick J, Tienari PJ, Laaksovirta H, Peuralinna T, Myllykangas L, Chio A, Shatunov A, Boeve BF, Boxer AL, DeJesus-Hernandez M, Mackenzie IR, Waite A, Williams N, Morris HR, Simón-Sánchez J, van Swieten JC, Heutink P, Restagno G, Mora G, Morrison KE, Shaw PJ, Rollinson PS, Al-Chalabi A, Rademakers R, Pickering-Brown S, Orrell RW, Nalls MA, Hardy J. Chromosome 9 ALS and FTD locus is probably derived from a single founder. *Neurobiol Aging* 2012;33:209.e3–209.e8.
- 51 van der Zee J, Gijssels I, Dillen L, Van Langenhove T, Theuns J, Engelborghs S, Philtjens S, Vandenbulcke M, Sleegers K, Sieben A, Bäumer V, Maes G, Corsmit E, Borroni B, Padovani A, Archetti S, Pernecky R, Diehl-Schmid J, de Mendonça A, Miltenberger-Miltenyi G, Pereira S, Pimentel J, Nacmias B, Bagnoli S, Sorbi S, Graff C, Chiang HH, Westerlund M, Sanchez-Valle R, Llado A, Gelpi E, Santana I, Almeida MR, Santiago B, Frisoni G, Zanetti O, Bonvicini C, Synofzik M, Maetzler W, Vom Hagen JM, Schöls L, Heneka MT, Jessen F, Matej R, Parobkova E, Kovacs GG, Ströbel T, Sarafov S, Tourné I, Jordanova A, Danek A, Arzberger T, Fabrizi GM, Testi S, Salmon E, Santens P, Martin JJ, Cras P, Vandenbergh R, De Deyn PP, Cruts M, Van Broeckhoven C, van der Zee J, Gijssels I, Dillen L, Van Langenhove T, Theuns J, Sleegers K, Bäumer V, Maes G, Cruts M, Van Broeckhoven C, Engelborghs S, De Deyn PP, Cras P, Engelborghs S, De Deyn PP, Vandenbulcke M, Vandenbulcke M, Borroni B, Padovani A, Archetti S, Pernecky R, Diehl-Schmid J, Synofzik M, Maetzler W, Müller Vom Hagen J, Schöls L, Synofzik M, Maetzler W, Müller Vom Hagen J, Schöls L, Heneka MT, Jessen F, Ramirez A, Kurzweily D, Sachtleben C, Mairer W, de Mendonça A, Miltenberger-Miltenyi G, Pereira S, Firmo J, Pimentel J, Sanchez-Valle R, Llado A, Antonelli A, Molinuevo J, Gelpi E, Graff C, Chiang HH, Westerlund M, Graff C, Kihluth Ståhlbom A, Thonberg H, Nennesmo I, Börjesson-Hanson A, Nacmias B, Bagnoli S, Sorbi S, Bessi V, Piaceri I, Santana I, Santiago B, Santana I, Helena Ribeiro M, Rosário Almeida M, Oliveira C, Massano G, Garet C, Pires P, Frisoni G, Zanetti O, Bonvicini C, Sarafov S, Tourné I, Jordanova A, Tourné I, Kovacs GG, Ströbel T, Heneka MT, Jessen F, Ramirez A, Kurzweily D, Sachtleben C, Mairer W, Jessen F, Matej R, Parobkova E, Danel A, Arzberger T, Maria Fabrizi G, Testi S, Ferrari S, Cavallaro T, Salmon E, Santens P, Cras P. European Early-Onset Dementia Consortium. A pan-European study of the C9orf72 repeat associated with FTD: geographic prevalence, genomic instability, and intermediate repeats. *Hum Mutat* 2013;34:363–73.
- 52 Xi Z, Zinman L, Moreno D, Schymick J, Liang Y, Sato C, Zheng Y, Ghani M, Dib S, Keith J, Robertson J, Rogava E. Hypermethylation of the CpG island near the G4C2 repeat in ALS with a C9orf72 expansion. *Am J Hum Genet* 2013;92:981–9.
- 53 Gijssels I, Van Mossevelde S, van der Zee J, Sieben A, Engelborghs S, De Bleeck J, Ivanou A, Deryck O, Edbauer D, Zhang M, Heeman B, Bäumer V, Van den Broeck M, Mattheijssens M, Peeters K, Rogava E, De Jonghe P, Cras P, Martin JJ, de Deyn PP, Cruts M, Van Broeckhoven C. The C9orf72 repeat size correlates with onset age of disease, DNA methylation and transcriptional downregulation of the promoter. *Mol Psychiatry* 2016;21:1112–24.
- 54 van Blitterswijk M, DeJesus-Hernandez M, Niemantsverdriet E, Murray ME, Heckman MG, Diehl NN, Brown PH, Baker MC, Finch NA, Bauer PO, Serrano G, Beach TG, Josephs KA, Knopman DS, Petersen RC, Boeve BF, Graff-Radford NR, Boylan KB, Petrucelli L, Dickson DW, Rademakers R. Association between repeat sizes and clinical and pathological characteristics in carriers of C9ORF72 repeat expansions (Xpansize-72): a cross-sectional cohort study. *Lancet Neurol* 2013;12:978–88.
- 55 Dols-Icardo O, García-Redondo A, Rojas-García R, Sánchez-Valle R, Noguera A, Gómez-Tortosa E, Pastor P, Hernández I, Esteban-Pérez J, Suárez-Calvet M, Antón-Aguirre S, Amer G, Ortega-Cubero S, Blesa R, Fortea J, Alcolea D, Capdevila A, Antonell A, Lladó A, Muñoz-Blanco JL, Mora JS, Galán-Dávila L, Rodríguez De Rivera FJ, Lleó A, Clarimón J. Characterization of the repeat expansion size in C9orf72 in amyotrophic lateral sclerosis and frontotemporal dementia. *Hum Mol Genet* 2014;23:749–54.
- 56 Fratta P, Polke JM, Newcombe J, Mizielinska S, Lashley T, Poulter M, Beck J, Preza E, Devoy A, Sidle K, Howard R, Malaspina A, Orrell RW, Clarke J, Lu CH, Mok K, Collins T, Shoaib M, Nanji T, Wray S, Adamson G, Pittman A, Renton AE, Traynor BJ, Sweeney MG, Revesz T, Houlden H, Mead S, Isaacs AM, Fisher EM. Screening a UK amyotrophic lateral sclerosis cohort provides evidence of multiple origins of the C9orf72 expansion. *Neurobiol Aging* 2015;36:546.e1–546.e7.
- 57 Nordin A, Akimoto C, Wuolikainen A, Alstermark H, Jonsson P, Birve A, Marklund SL, Graffmo KS, Forsberg K, Brännström T, Andersen PM. Extensive size variability of the GGGGCC expansion in C9orf72 in both neuronal and non-neuronal tissues in 18 patients with ALS or FTD. *Hum Mol Genet* 2015;24:3133–42.
- 58 Gami P, Murray C, Schottlaender L, Bettencourt Conceição, De Pablo Fernandez E, Mudanohwo E, Mizielinska S, Polke JM, Holton JL, Isaacs AM, Houlden H, Revesz T, Lashley T. A 30-unit hexanucleotide repeat expansion in C9orf72 induces pathological lesions with dipeptide-repeat proteins and RNA foci, but not TDP-43 inclusions and clinical disease. *Acta Neuropathol* 2015;130:599–601.
- 59 Akimoto C, Volk AE, van Blitterswijk M, Van den Broeck M, Leblond CS, Lumbroso S, Camu W, Neitzel B, Onodera O, van Rheenen W, Pinto S, Weber M, Smith B, Proven M, Talbot K, Keagle P, Chesi A, Ratti A, van der Zee J, Alstermark H, Birve A, Calini D, Nordin A, Tradowsky DC, Just W, Daoud H, Angerbauer S, DeJesus-Hernandez M, Konno T, Lloyd-Jani A, de Carvalho M, Mouzat K, Landers JE, Veldink JH, Silani V, Gitler AD, Shaw CE, Rouleau GA, van den Berg LH, Van Broeckhoven C, Rademakers R, Andersen PM, Kubisch C. A blinded international study on the reliability of genetic testing for GGGGCC-repeat expansions in C9orf72 reveals marked differences in results among 14 laboratories. *J Med Genet* 2014;51:419–24.
- 60 Cleary EM, Pal S, Azam T, Moore DJ, Swingler R, Gorrie G, Stephenson L, Colville S, Chandran S, Porteous M, Warner JP. Improved PCR based methods for detecting C9orf72 hexanucleotide repeat expansions. *Mol Cell Probes* 2016;30:218–24.



Intermediate C9orf72 alleles in neurological disorders: does size really matter?

Adeline S L Ng and Eng-King Tan

J Med Genet 2017 54: 591-597 originally published online July 8, 2017
doi: 10.1136/jmedgenet-2017-104752

Updated information and services can be found at:
<http://jmg.bmj.com/content/54/9/591>

These include:

References

This article cites 59 articles, 3 of which you can access for free at:
<http://jmg.bmj.com/content/54/9/591#BIBL>

Open Access

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Clinical genetics](#) (257)
[Open access](#) (191)

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>