Phenocopies

W. LENZ

Institut für Humangenetik, University of Münster, Federal Republic of Germany

Phenocopies by Heat Treatment

Goldschmidt (1935) proposed the concept of phenocopy for what he thought to be 21 different types of abnormal wings and 10 different types of abnormalities of bristles, eyes, and other organs produced by heat treatment of *Drosophila* larvae. He expressly stated: 'The enumeration shows already that the phenotypes of the great majority of mutants known in *Drosophila* can be produced in these experiments as non-genetic modifications. I do not doubt that in fact any phenotype of any mutant may be copied in this way if we succeed in creating still more possibilities of influence by further modifications of experimental conditions.'

Even earlier, Zeleny (1923), having demonstrated the effects of temperature differences on the phenotype of the bar eye mutants of *Drosophila* clearly stated 'The gene ultrabar has the same type of reaction as a temperature difference'. In F. Fischer's earlier experiments (1895) on butterflies, heat produced wing patterns identical to that of a known geographic variety. Fischer referred to this effect as 'transmutation'. One reason why similar results have not been attained in mammals is probably the smaller range of temperatures compatible with survival in warm-blooded animals than in insects.

The similarity between genetic phenotypes in *Drosophila* and their heat-induced phenocopies may in part be explained by heat inactivation of enzymes.

Heat inactivation of genetic enzyme variants at temperatures compatible with survival is well-known from many microorganisms, particularly from *E. coli*, but also from *Neurospora*, *Primula sinensis*, rabbits, and Siamese cats. In *E. coli* there are also temperature sensitive mutants of the *lac*-operon regulator gene and of a gene controlling initiation of

chromosomal replication (for references see Bresch and Hausmann, 1970).

Heat instability is also a feature of several genetic haemoglobin variants in man (Genova, Torino, Bibba, Hammersmith, Sydney) usually associated with inclusion body anaemia. The heat effect in these cases, however, is important for diagnosis *in vitro* rather than for clinical manifestations. Furthermore, heat sensitivity is found in several variants of glucose-6-phosphate dehydrogenase (Chicago, Ohio, Oklahoma, Duarte, Albuquerque, Eyssen), associated with spontaneous haemolytic anaemia.

High temperature does not appear to induce clearcut phenocopies in mammals. No association between fever during the first trimester and malformation of the infant has been found. Virus infections with a rise in temperature to at least 38° C have been found in 2.7% of 342 pregnancies resulting in malformed children and in 2.5% of 8533 pregnancies resulting in non-malformed children (Villumsen, 1970). The incidence of bacterial infections in the first trimester in the same material was 1.8% for malformed children and 2.9% for non-malformed children.

Hypoxia and Abnormal Implantation

Büchner (1958), produced a variety of malformations in various amphibian and avian species by hypoxia. Similar to Goldschmidt's assumption (1935) for *Drosophila* he felt that in man 'each genetic malformation can be copied by one caused exogenously'; thus, Büchner continues, 'exogenous malformations are in principle phenocopies of malformations which may also be genetic'. This idea has not received any firm support from observations in man. Various attempts to link poor implantation, bleeding, or general anaesthesia in early pregnancy with hypoxia and consecutive malformations have been unsuccessful (see Villumsen, 1970).

There is no statistical evidence nor any evidence in selected cases that hypoxia induces the same types of malformations in human embryos which, as a rule, have a genetic basis. Grebe (1954) in a review on 'Genes and phenocopies in the aetiology of human malformations', has discussed phenocopy in relation to mongolism, anencephaly, cleft lip and palate, and limb malformations. None of these examples of presumed phenocopies is convincing.

Abnormal implantation has been thought to be a cause of hypoxia and malformations. The evidence is, at best, indirect. The most clearcut type of abnormal implantation, ie, extra-uterine pregnancy, is commonly associated with gross deformities of clearly exogenous mechanical origin quite different from any genetic syndrome or from true developmental malformations (von Winckel, 1902; Lelling, 1938). There is no evidence that extra-uterine pregnancy might produce phenocopies, and the same is probably true for other types of abnormal implantation.

Induction of Phenocopies by Chemicals

Rapoport (1939) has produced morphological abnormalities in *Drosophila* similar to mutant phenotypes by adding various chemicals, especially salts of metals, some of them known to be enzyme inhibitors, to the food. He tested only one stock of *Drosophila* and found specific effects associated with specific chemicals. More detailed studies comparing chemical phenocopies in *Drosophila* with known mutants have been published by Gloor (1947), Bodenstein and Abdel-Malek (1949), Sang and McDonald (1954), and Goldschmidt and Piter-

nick (1957). When Goldschmidt and Piternick (1957) studied the genetic background of borate-induced phenocopies in *Drosophila* they were inclined to attribute the conspicuous genetic differences observed between stocks to common subthreshold mutants brought out by borate. A similar argument may be applied to heat-induced phenocopies which also vary with the genotype of the experimental animals. Heat sensitive enzyme variants may be conceived as products of genes which are subthreshold under ordinary conditions.

Landauer (1957 and 1958) has been able to produce the rumpless phenotype in chickens by insulin, boric acid, or pilocarpin. The results varied with the genotype. Landauer thinks the term phenocopy is inappropriate for cases in which the recessive gene for rumplessness becomes apparent through chemical teratogens. He prefers to reserve the term 'phenocopy' for experimentally produced conditions identical in phenotype and, at least in part, in developmental pathway to a genetic condition. Rubella embryopathy, according to Landauer (1959), should not be called a phenocopy. In Ancel's monograph (1950) on chemical induction of malformations in vertebrates the word phenocopy appears neither on the five contents pages, nor on the four pages of the index. Thus confusion is avoided.

Defects of the caudal spine similar to rumplessness in chickens are known also in man. The more extensive defects which include not only the sacrum but several lumbar and sometimes even thoracic vertebrae are sometimes associated with maternal diabetes (Ghiaï, 1972; see Figs. 1, 2a, and 2b).

FIG. 1. Agenesis of sacrum in the child of a diabetic mother. (By courtesy of Professor von Harnack, Universitäts-Kinderklinik, Düsseldorf.)

FIGS. 2a and 2b. Agenesis of the lumbar spine and sacrum in a child born to a diabetic mother (onset of diabetes at 29 yr, parturition at 34 yr). (Paediatric Clinic, Münster University.)

Partial and often asymmetric sacral defects are sometimes found associated with thalidomide-induced anotia. In both instances, however, phenocopy would be a misleading term, as no genetic phenotype which exactly corresponds is known in man.

Between 1935 and 1957 Goldschmidt's use of the term phenocopy underwent considerable change, which has not been generally realized, for he often failed to clarify to what he was referring. He applied the term phenocopy rather indiscriminately to: (1) phenotypes produced by environmental factors in a genotype otherwise not associated with the same phenotype, if the same phenotype also occurs on a genetic basis without the environmental factor; (2) phenotypes produced by environmental factors, but so far unknown in other genotypes; (3) phenotypical effects of specific genes, when these are only apparent under certain environmental conditions and identical with the usual phenotypical effects of other possibly is allelic genes.

Many medical authors have followed this vague and conflicting nomenclature (eg, Büchner, 1958; Degenhardt, 1958).

Henke, von Finck, and Ma (1941), studying heat modifications in *Drosophila* made a distinction between *true* phenocopies, in which the environmental agent affects the same developmental phase as the gene (the effect of which is copied) and *false* phenocopies, in which only the end result is identical.

A similar point of view is expressed by Hadorn (1955), who defined phenocopy on the basis of two strict criteria: (1) identity in all details of the pattern of phenotypical manifestation with a locus specific genetic pattern; (2) identity of the developmental path leading to the trait.

Nachtsheim (1957 and 1961) defended the rather vague application of the concept of phenocopy. His main argument was that it is not usually possible to trace the origin of a genetic trait in man, so the question whether a phenocopy is a true or a false one has to be left open.

With increasing precision in description and diagnosis of genetic entities, a wider use of the term phenocopy in man has become dubious, if not obsolete. Nachtsheim (1961) discussed epilepsy, haemophilia A and B, afibrinogenaemia, and deficiencies of clotting factors V, VII, and X as genetic conditions which may also be phenocopied. Obviously, supposed phenocopies of these conditions are identified by single symptoms rather than by the total pattern of the disease. Many more symptoms have either a genetic or an environmental cause or combined causes; to use the term 'phenocopy' to refer to this well-known medical fact no longer serves any useful purpose.

In the third edition of his catalogues of Mendelian inheritance in man, McKusick (1971) lists 943 autosomal dominant, 783 autosomal recessive, and

150 X-linked phenotypes. It is quite obvious to anyone acquainted with any of the fields of medical genetics that, at best, true phenocopies are established in an almost negligibly small minority of these conditions, and that there are no specific reasons to suspect phenocopies for almost every well-defined monogenic entity. This is not unexpected for monogenic phenotypes are basically molecular diseases or polymorphic variants. Environmental influences rarely affect only one gene product and even when they do, there is still the difference between the permanent genetic block and the temporary and variable block brought about by inhibitor substances or heat inactivation.

At first glance, vitamin deficiencies would appear to be good candidates for phenocopies. One might expect that environmental lack of a vitamin would produce the same effect as a genetic block of its resorption, metabolism, or attachment to the protein apoenzyme. The similarity between vitamin D deficiency rickets and pseudo-vitamin D deficiency rickets is striking. However, this is a rather special case; deficiency of one vitamin does not usually occur without deficiencies of other vitamins or other dietary constituents and is not usually permanent, as is the genetic block.

Phenocopies may also be expected to occur in endocrinological disorders. Most endocrine functions are limited to one or a few glands which, with the exception of the gonads, have no known function other than that of producing hormones. Hormone production may be defective either because one of the genetic biochemical links is missing, or because some external agent may have damaged a particular gland. Two examples are listed in Table I.

TABLE I

Disease	Genetic Prototype	Phenocopy
Panpituitary dwarfism	Autosomal recessive (Rimoin <i>et al</i> , 1968)	Birth trauma Bierich (1965)
	X-linked recessive (Rimoin and Schimke, 1971)	
Pendred's syndrome	Autosomal recessive-defective organization of iodide with deafness (Harnack <i>et al</i> , 1961)	Endemic cretinism due to environmental iodine deficiency (König, 1968)

The similarity between the two genetic types of panpituitary dwarfism and that caused by birth trauma to the pituitary body (Bierich, 1965) is striking. A somewhat similar phenotype may also

be the consequence of emotional and nutritional deprivation (Silver and Finkelstein, 1967). As in genetic and traumatic pituitary dwarfism these cases show retardation of growth in height, bone age, and sexual maturation. There may be some metabolic features indicative of pituitary failure, but there are distinctive marks of deprivation dwarfism such as voracious appetite and an increase in weight and height, shortly after the psychological situation has been improved.

Pendred's syndrome and endemic cretinism share two main symptoms—deafness and goitre. Mental deficiency and low stature, prominent in endemic cretinism, are absent or rare in Pendred's syndrome. Still, it is tempting to speculate that the association of deafness and goitre in both diseases is due to a similar mechanism, ie, the lack of iodine in the inner ear and in the thyroid.

In their comprehensive book, 'Genetic disorders of the endocrine glands', Rimoin and Schimke (1971) are very open-minded towards the possibilities of phenocopies. Table II cites some examples from their article.

TABLE II

Genetic Condition	'Phenocopy'
Autosomal recessive: Schmidt's syndrome (polyendocrine deficiency syndrome)	Virus infection? autoantibody formation?
X-linked dominant, autosomal dominant: decrease of thyroxine-binding globulin	Cirrhosis, nephrotic syndrome, malabsorption states
Autosomal recessive or X-linked recessive: 18-dehydrogenase deficiency (hypoaldosteronism)	Drug induced (heparin)
Autosomal recessive: Bartter syndrome	Sodium depletion by chronic laxative abuse

The external causes of the polyendocrine deficiency syndrome are not well established. The other examples are obviously copies of single signs rather than of disease patterns. Generally, phenocopies are rare, and play a minor role in practical medicine and genetic counselling. They are, however, most important as models which might help to elucidate the pathways of gene action.

Genetic Limb Malformations and Thalidomide Phenocopies

No phenocopies are known for most types of monogenic limb malformations found in man, such as split hand and split foot; the various types of brachydactyly, polydactyly, and syndactyly; the

genetic types of peripheral defects, and many other complex syndromes. There are, however, several specific genetic types of bone defects and of triphalangy of the thumb which are sufficiently similar to some thalidomide cases to make a discussion of the phenocopy concept meaningful. The following seven types will be selected for discussion.

1. Autosomal dominant aplasia of the radius (Holt-Oram syndrome).
2. Radius aplasia-thrombocytopenia syndrome.
3. Fanconi's panmyelopathy.
4. Roberts' syndrome (tetraphocomelia with cleft lip and palate).
5. Pseudothalidomide syndrome.
6. Tibia aplasia with triphalangy of the thumb.
7. Triphalangy of the thumb.

Types 1, 6, and 7 are autosomal dominant, the other types are autosomal recessive. The morphology of the individual bone defects as well as of their associations in types 1, 3, 6, and 7 is very much the same as in many thalidomide cases. Not all thalidomide cases, however, are phenocopies. Complete absence of the arms, some forms of bony syndactyly of the fingers, proximal femoral defects, total or subtotal defects of all of the long bones of the legs, aplasia of the first toes, etc, do occur in some thalidomide cases, though not in any of the seven types listed above nor in any other monogenic phenotypes in man. Cardiac malformations may be associated with types 1, 2, and 3 and renal malformations with type 3. However, some more severe cardiac defects and other internal malformations—comprising duodenal stenosis, aplasia of the gallbladder, aplasia of the appendix, uterus bicornis, etc—do occur in many severe thalidomide cases, but not in most of the seven genetic prototypes. The association of thalidomide limb deformities with anotia, abducens and facial paralysis, microphthalmos and coloboma has no counterpart in any known genetic syndrome. The midfacial haemangioma, on the other hand, may be seen in types 1, 2, and 4.

The spectrum of each of the seven genetic prototypes is much more restricted than that of thalidomide embryopathy. Thalidomide embryopathy is not a single syndrome, definable on the basis of its component symptoms, but a teratogenic series of interconnected syndromes. Cases of thalidomide embryopathy with exactly the same time of intake constitute well-defined syndromes: such as a syndrome of anotia, facial and abducens paralysis, dysplasia of the sacrum, and ectopic kidney following thalidomide intake around the 35th to 37th postmenstrual day, or a syndrome of anorectal stenosis, inguinal hernia, and triphalangy of the

thumbs following ingestion of the drug around the 48th to 50th postmenstrual day. There is no overlap between these two phenotypes following very early and very late intake. If, however, cases with intake at various intervals in between, or cases with both early and late intake are studied, a clear pattern of the interconnected series of syndromes is seen.

The similarities and differences between thalidomide embryopathy and each of the seven genetic prototypes will now be discussed in more detail.

Prototype 1

Dominant Radius Aplasia

The main symptoms common in both dominant radius defects and thalidomide embryopathy are listed in Table III. There are no clear-cut signs present only in the dominant radius aplasia but lacking in thalidomide cases, though clinodactyly of the fifth fingers and mild funnel chest appear to be common in the genetic prototype but not in thalidomide cases. The difference between prototype 1 and its phenocopy may be in part explained in terms of the time sequence of organogenesis. The dominant gene appears to affect development only between the 42nd and 50th postmenstrual day. This may explain why ear defects and renal malformations are absent. But difference in time cannot explain everything. The defects of the lower limbs in thalidomide cases originate at the same time—around the 44th to 46th day—when thalidomide causes arm defects identical to those seen in the dominant radius defect syndrome. This may be explained by assuming that the same normal

TABLE III

	Dominant Radius Defect	Thalidomide Embryopathy
Duane syndrome	+	+
Proximal defect of humerus	+	+
Radius hypoplasia	+	+
Radius aplasia	+	+
Radioulnar synostosis	+	+
Thumb hypoplasia	+	+
Thumb aplasia	+	+
Thumb triphalangy	+	+
Heart (atrial septal defect)	+	+
Heart (ventricular septal defect)	+	+
Midface haemangioma	+	+
Anotia	—	+
Amelia of arms	—	+
Duodenal atresia	—	+
Complex cardiovascular malformations	—	+
Kidney malformations	—	+
Femur defects	—	+
Tibia defects	—	+

embryological process is blocked to a lesser degree by the gene than it is in some instances by thalidomide. If thalidomide malformations are studied in relation to the amount of the drug ingested, it is apparent that leg malformations do not usually occur at doses less than 100 mg/day taken for more than one day. It is a well-known fact from experimental teratogenesis in mammals, that the hind legs are more resistant to teratogenic agents than the arms (von Kreybig, 1968).

One striking fact about the dominant prototype 1 is the variability of the phenotype. A mother with triphalangy or hypoplastic thumbs only and with no cardiac defect may have a child with severe three-finger phocomelia with distal humerus rudiments. A similar difference in thalidomide cases can be attributed to a time difference of about one week—50th versus 43rd postmenstrual day. It is difficult to understand how one and the same gene may have different times of activity in different patients.

The Duane syndrome has been reported in a few patients with prototype 1 (Mennerich, 1923; Gifford, 1926; Ferrell, Jones, and Lucas, 1966). The relative scarcity of reported cases of Holt–Oram syndrome showing this association may be misleading. The Duane syndrome may easily escape the attention of a physician mainly interested in the more conspicuous orthopaedic or cardiologic problems of his patients. Judging from the thalidomide experience the Duane syndrome has a still earlier origin at around the 35th to 37th postmenstrual day. The association of two such rare and not directly related conditions as radius defects and the Duane syndrome both in the genetic prototype 1 and in thalidomide embryopathy suggests a common biochemical pathway. The Duane syndrome can also occur alone as a dominant condition. The Duane syndrome within the framework of thalidomide intake is not a true phenocopy, as it is almost invariably associated with ear defects; an association apparently unknown as a genetic entity.

Both abducens and facial paralysis in thalidomide cases are usually interpreted as showing that thalidomide does not only affect mesodermal structures, such as muscles and bones, but also ectodermal structures (Partsch, 1969). I doubt whether this is justified, although thalidomide might interfere primarily with the development of the facial and eye muscles. The close proximity of the premyoblast cells from which the facial muscles originate to the origin of the external ear in the 4.2–6.5 mm embryo (Gasser, 1967) might explain the association of anotia and so-called cranial nerve defects in thalidomide embryopathy.

Prototype 2

Radius Aplasia–Thrombocytopenia Syndrome

Superficially, this syndrome with radius aplasia or phocomelia of the arms is strikingly similar to many cases of thalidomide embryopathy. The similarity extends to midfacial haemangioma, cardiovascular defects, and dislocation of the hip. There are, however, several clearly distinctive signs which have been found in every case of prototype 2 and in no case of thalidomide embryopathy. These are thrombocytopenia with deficiency of megakaryocytes in the bone marrow; neonatal leucocytosis; essentially normal thumbs, but with the radii invariably absent (see Fig. 3); and brachymesophalangy of the fifth fingers.

Ear defects, paralysis of cranial nerves, renal malformations, bone defects of the legs, and triphalangy of the thumbs do not occur in the radius aplasia–thrombocytopenia syndrome. The critical period of organogenesis which is affected in this syndrome would appear to be narrowly restricted to about the 44th to 45th postmenstrual day. The difference between the genetic prototype 2 and thalidomide embryopathy, however, cannot be explained by differences in time and/or by the degree of inter-

Fig. 3. X-ray typical of radius aplasia–thrombocytopenia syndrome in one of two affected sibs. (Orthopaedic Clinic, Hüffer-Stiftung, Münster University.)

ference with normal organogenesis. There is an obvious qualitative difference: in prototype 2 a small misshaped rudiment may be all that is left of the three long bones of the arms, and yet the hand has five metacarpals and five fingers, whereas in thalidomide cases long bone defects of the arm of a similar degree are invariably accompanied by aplasia of other fingers as well. Differential diagnosis is unambiguous in every single cases of the radius aplasia-thrombocytopenia syndrome. Thus, even if the platelet deficiency is not considered, thalidomide does not produce a true phenocopy of prototype 2.

The skeletal phenotype of prototype 2 alone is sufficiently distinct to make the diagnosis virtually certain. Cases in which the platelets have not been studied but with the same pattern of bilateral radius aplasia and five fingers present probably should be included in the same category (Vrolik, 1849; Gruber, 1865; Rodenstein, 1875/76; Blencke, 1904; Homi, 1920; Hill, 1937; Birch-Jensen, 1949 [p. 104; Figs. 58 and 427]; Barsky, 1958; Gerold, 1959; Matzen und Fleissner, 1969 [Fig. 589]; G. Farkas, personal communication). These cases have never been found with dominant transmission. The distinction from dominant radius aplasia is clear cut both by phenotypical and genetic criteria. Two or more cases within one sibship have been reported by Blencke (1904), by Barsky (1958), and by Gerold (1959).

The genetics of the radius aplasia-thrombocytopenia syndrome is not quite clear. The incidence of the condition for autosomal recessive inheritance in sibs of probands agrees fairly well with expectation. Consanguinity of parents has not been noted so far. Formally, this absence of parental consanguinity in a rare familial condition, may be explained by postulating a compound heterozygote of two alleles t_1 and t_2 of the normal allele t_0 . If homozygotes for t_2 were lethal in early embryonic life, the possibility of detecting them would be minimal. But what about the homozygotes for t_1 ? They might represent Fanconi's panmyelopathy, which has thrombocytopenia, radial defects, and sometimes cardiac defects in common with the radius defect-thrombocytopenia syndrome. Alternatively, both homozygotes, t_1t_1 and t_2t_2 , might be lethal and the compound, t_1t_2 , might show imperfect intragenic complementation.

Prototype 3

Fanconi's Panmyelopathy

Fanconi's panmyelopathy can be associated with a surprising variety of malformations, most of which

TABLE IV
MALFORMATIONS FOUND IN FANCONI'S
PANMYELOPATHY (Bátke, 1964)

Radius aplasia	Persistent ductus arteriosus
Radius hypoplasia	Duodenal stenosis
Radioulnar synostosis	Rudimentary appendix
Thumb aplasia	Anal atresia
Thumb hypoplasia	Right lung—two lobes
Thumb triphalangy (rare)	Inguinal hernia
Thumb duplication	Microphthalmos
Pelvic dystopia of kidney	Strabism
Hydronephrosis	Facial paralysis
Renal aplasia	Epicanthal folds
Atresia of auditory canal	Dislocation of hips
Aplasia of internal ear	Coccygeal dimple
Malformed auricles	Choanal atresia
Deafness	

TABLE V
OCCURRENCE OF SOME MALFORMATIONS IN
133 CASES OF FANCONI'S PANMYELOPATHY
(Bátke, 1964)

	Bilateral	Unilateral
<i>Radius</i>		
Aplasia	6 cases	5 cases
Hypoplasia	1	5
Radioulnar synostosis	1	—
<i>Thumb</i>		
Aplasia	10	13
Hypoplasia	15	19
Unspecified defects	4	1
Duplication	—	7

also occur in thalidomide cases (see Tables IV and V). The incidence of internal malformations in cases of Fanconi's anaemia reported in the literature is heavily influenced by the fact that surviving cases are much more likely to be published. If the index cases are left aside, the very high incidence of gross malformations in stillborn sibs tends to give much higher estimates.

Most of these malformations, especially deafness and renal defects, correspond to the effect of the intake of small doses of thalidomide during the first half of the sensitive period of thalidomide embryopathy, ie, between the 35th and 42nd day; malformations phenocopied later, like triphalangy of the thumbs, are rare. While the pattern of organ malformations is very similar in Fanconi's panmyelopathy and in thalidomide embryopathy, both conditions have additional distinctive symptoms, so that serious diagnostic doubts do not occur. Thrombocytopenia, leucopenia, and anaemia as well as pigment spots, dwarfism, and chromosome breaks typical of Fanconi's panmyelopathy are not seen in thalidomide embryopathy. It does not necessarily follow that the basic mechanism is different in both conditions. The same biochemical mechanism might be permanently but incompletely blocked in the genetic prototype and intermittently in the

pharmacological phenocopy. Permanent high intake of thalidomide during the whole sensitive period seems to be invariably lethal in human as well as in some other primate embryos.

Thrombocytopenia has been a rather common side effect of thalidomide taken by adults. It has been attributed to drug allergy, but this has never been proved. The mechanism may be the same as in Fanconi's anaemia.

Prototypes 4 and 5

There is some doubt about the identity (J. Opitz, personal communication) or non-identity (McKusick, 1971) of type 4 (Roberts' syndrome [1919] of tetraphocomelia) and type 5 (the SC-pseudothalidomide syndrome). In Roberts' syndrome the limb defects are more severe, there is bilateral cleft lip and palate, and the face is definitely malformed with a flat nose and prominent eyes. There is no close similarity in any details to thalidomide cases. Bilateral cleft lip and palate do occur in thalidomide cases, though rarely, and then only in association with ear malformations.

In the pseudothalidomide syndrome only the arms may be malformed, while the legs remain normal, or the fibulae may be missing with the tibiae present and fused with the femora; this type of defect is unknown in thalidomide cases. Severe hypoplasia of the thumbs and the fifth fingers, presence of all fingers even with aplasia of radius and ulna, and synostosis of the fourth and fifth metacarpals are further characteristics of the pseudothalidomide syndrome. Thalidomide cases are not phenocopies of either syndrome, the morphological differences suggest different developmental mechanisms. In Roberts' syndrome (1919) and in the pseudothalidomide syndrome, however, the same basic mechanism may well be affected. If the differences do not represent variable manifestations of the same gene, they may be due to isallelic mutations. Cases with intermediate morphology, such as the one reported by Ströer (1939)—cleft lip and palate, well developed humeri, aplasia of radius, ulna and fibula, synostosis of the fourth and fifth metacarpals, aplasia of thumbs—suggests that both conditions may only be the extreme manifestations of one entity. Alternatively, such intermediate cases may represent independent isallelic mutations. For instance, in Ströer's case (1939) a compound heterozygote of the two isallelic mutations underlying Roberts' syndrome (1943) and the pseudothalidomide syndrome is virtually excluded, because the parents were related. Ströer's assumption (1939), that cleft lip and palate without limb deformities when present in a few members of the

same family, is a manifestation of the heterozygote state, receives at best weak support from his published pedigree and none from other reports.

Prototype 6

Tibia Aplasia with Triphalangy of Thumbs

Bilateral aplasia or severe hypoplasia of the tibia in association with excessive preaxial polydactyly of the feet and triphalangeal thumbs has been observed with a clearly autosomal dominant pattern of transmission (Eaton and McKusick, 1969). In some members only triphalangy of the thumbs with preaxial polydactyly of the hands (Lenz, Theopold, and Thomas, 1964, one unpublished observation) or triphalangy of the thumbs only (Reber, 1967/68) is seen.

Reber also found a case of postaxial hexadactyly of both hands in a distant relative, related to the patient through seven unaffected persons. There is no good reason for his assumption that this case has the same genetic basis as preaxial polydactyly in father and son.

Tibia aplasia with duplication of the big toes and triphalangeal thumbs also occurs following thalidomide intake between the 46th and 50th postmenstrual day (see Figs. 4a and 4b). In the genetic prototype the triphalangeal thumbs are rather well developed, preaxial polydactyly of the hands is more common, and the polydactyly of the feet may be excessive—eight toes on either foot. Polydactylism with postminimus of the hands is only seen in some cases of the genetic syndrome.

The analogy between genetic prototype and phenocopy is so close that it is difficult to escape the conclusion that the same mechanism is affected. The existing differences may be due to differences in time and degree of the block. The tibia aplasia-triphalangy syndrome seems to represent a block acting during a shorter period, than in the Holt-Oram syndrome.

Prototype 7

Triphalangeal Thumbs with and without Preaxial Polydactyly

Triphalangeal thumbs which sometimes look very much like index fingers can occur in the Holt-Oram syndrome, in the tibia aplasia-triphalangy syndrome, and, though rarely, in Fanconi's panmyelopathy. Triphalangy of the thumbs with congenital anaemia and with a ventricular septal defect in one case, as seen by Aase and Smith (1969) in two brothers, may be a separate entity. In families with typical Holt-Oram syndrome some members may show triphalangeal thumbs as the only abnormality. It is

FIGS. 4a and 4b. Thalidomide embryopathy. Irregular intake of thalidomide before and during early pregnancy.

FIGS. 5a and 5b. Thalidomide embryopathy: triphalangy and polydactyly. The first of several prescriptions of thalidomide was probably taken on the eighth postmenstrual day if the average duration of pregnancy is assumed.

FIG. 6. Thalidomide embryopathy. Thalidomide was prescribed during the second month of pregnancy. The right hand is an almost exact mirror image of the left hand.

FIGS. 7a and b. Dominant triphalangeal thumbs with preaxial polydactyly.

difficult, therefore, to decide whether in a given family with few affected members triphalangy of the thumbs is a phenotypical variety of the Holt–Oram syndrome, or an independent entity. In some families, triphalangy of the thumbs with or without concomitant cardiac malformations is seen in some members, whereas aplasia of the thumbs and radio-ulnar synostosis is seen in others (Haas, 1939; Ferber, 1953). These cases probably belong to the Holt–Oram syndrome, and they may show the mesobrachyphalangy of the fifth fingers and funnel chest often found in that condition. In other families triphalangy is associated with duplication and even triplication of the thumbs. Nylander (1931) observed duplication of the thumbs and higher degrees of preaxial polydactylism, up to five thumbs, usually combined with triphalangy in 41 members of a large Swedish kinship. Müller (1936) reported a family with 11 members in three generations affected by triphalangeal thumbs and preaxial polydactyly.

In some families, triphalangeal thumbs are associ-

ated with preaxial polydactyly of the hands and feet (Atwood and Pond, 1917; Manoiloff, 1931; Haas, 1939; Hefner, 1940; Barsky, 1951; Komai, Ozaki, and Inokuma, 1953; Hopf, 1959; Demarinis and Wildervanck, 1960; Kersting, 1966 [case 8]; Temtamy, 1966 [Fig. 7]). Another dominant type is triphalangy with doubling of the terminal phalanges of the thumbs (Lapidus, Guidotti, and Coletti, 1943; Ecke, 1962). Swanson and Brown (1962) observed duplication of the first phalanx in three out of 30 members of a family with bilateral triphalangeal thumbs. Two out of four members affected with triphalangy in a family reported by Müller (1937) had duplicated thumbs as well. Wildervanck (1954) saw additional trapezoid shaped middle phalanges of the thumbs in six members in three generations of a family. Unilateral duplication of the terminal phalanx was present in one case. In a family reported by Ströer (1936) 16 patients in five generations had triphalangeal thumbs, preaxial polydactyly and complete syndactyly of the third to fifth fingers or fourth and fifth fingers bilaterally.

FIG. 8a. Triphalangeal thumbs. Autosomal dominant type. There are six cases in two generations of this family.

FIG. 8b. The same anomaly in a second cousin of the patient shown in Fig. 8a. (Institute of Human Genetics, Münster University.)

In still other families triphalangeal thumbs are inherited as an isolated autosomal dominant trait with a more regular manifestation than is usually found in Holt–Oram families (Salzer, 1898; Roberts, 1943; Sallam, 1955; Klemm, 1956; Barsky, 1958; Abramowitz, 1959; Frere, 1960; Kersting, 1966 [case 5; see Fig. 8]). Rudimentary postaxial polydactyly is occasionally seen in members of families with this later type of application, in which preaxial polydactyly is rare (Rieder, 1900; Cotte, 1923).

Triphalangy of the thumbs in thalidomide cases is variable. It is sometimes associated with polydactylism of the hands or feet (see Figs. 6 and 9). But judging from the thalidomide experience the dominant types of triphalangeal thumbs represent a genetic block strictly limited in time to the end of the sensitive period of organ development.

Varus angulation of the femur, similar to that seen in some late-intake thalidomide cases in association with triphalangeal thumbs and occasionally in the radius aplasia—thrombocytopenia

syndrome, has been described by Polinelli (1962) in one affected member of a family with six cases of triphalangy of the thumbs in four generations—where one case was associated with preaxial polydactyly of the hands.

One might, of course, attempt to split triphalangy of the thumbs into still more types. It is not quite clear, however, how they should be delineated. While in a given family the morphological type tends to be generally the same and different from that seen in many other families with triphalangy, there is some intrafamilial variability and some overlap between types.

One might, furthermore, enlarge the list of prototypes of thalidomide embryopathy by adding: (1) the autosomal recessive poikiloderma–radius aplasia syndrome; (2) isolated preaxial polydactyly of the feet, or (3) the dominant isolated radioulnar synostosis, which is very much like the same abnormality seen in thalidomide cases. In thalidomide embryopathy, however, skin changes are absent, preaxial polydactyly of the feet does not

FIG. 9. Thalidomide embryopathy. Two tablets of thalidomide were taken on the 45th postmenstrual day.

occur without hand malformations, and radioulnar synostosis is always associated with aplasia or hypoplasia of the radial ray.

Summary of Comparisons of Thalidomide Phenocopy with Genetic Prototypes

Thalidomide embryopathy comprises a great variety of skeletal and other malformations, many of which are also seen in genetic disorders, sometimes in the same characteristic patterns. Many cases of thalidomide embryopathy closely mimic genetic malformations. They are probably true phenocopies. However, the concept 'phenocopy' does not adequately describe the total phenotypical spectrum of thalidomide embryopathy, which comprises many complex malformation patterns for which a genetic prototype is unknown. The monogenic prototypes differ from thalidomide embryopathy in several ways.

1. There is less variability in degree.
2. There is less variability in the timing of the block interfering with embryonic development.
3. Some genetic prototypes show symptoms unknown in thalidomide embryopathy. This difference probably reflects either different basic mechanisms (probably for types 4 and 5), or the consequences of a permanent partial block in the genetic prototype, as compared to an intermittent block in thalidomide embryopathy.
4. Thalidomide cases do show some malformations not seen in the genetic prototypes. This may be due to temporary complete blocking of a metabolic pathway, the permanent genetic blocking of which would be lethal.
5. Some genetic prototypes predominantly show malformations corresponding to a block in the first half of the sensitive period (prototype 3), others predominantly to a block in the middle period (prototypes 1 and 2), still others to a late block (prototypes 6 and 7).

Speculations on the Nature of the Thalidomide Block

Nudleman and Travill (1971) have produced defects of the preaxial structures of the upper limbs in rabbits by feeding thalidomide to the females on days 7–11 of pregnancy. The fetuses were removed on day 21. The most striking difference between experimental and control animals was seen in the glycogen content of hypertrophic chondrocytes and calcified chondrocytes, which were depleted of glycogen in the controls and filled in the thalidomide-treated animals. A similar interference with glycogen utilization was seen in the

hypertrophic chondrocytes of the sternebrae of thalidomide-treated rats (Globus and Gibson, 1968). These experimental findings fit in with the speculation that thalidomide might interfere with mobilization of embryonic glycogen, based on the correspondence between the distribution of embryonic glycogen in the embryo and the pattern of thalidomide malformations (Lenz, 1963). Dyban (1962) has studied the glycogen content of different tissues in human embryos and collated the data with similar studies from other mammals. Accumulation typically precedes morphological differentiation of the blastemas, from which the heart and the gut, the auditory canal, the muscles, and the skeleton originate. This is not the case, or only to a lesser extent in the precursor cells of the liver, the spleen, the gonads, and the central nervous system. Embryonic glycogen appears to be different from adult glycogen. It is more resistant to histochemical procedures. If the teratogenic action of thalidomide is explained by interference with glycogen mobilization, it becomes necessary to postulate different metabolic pathways for embryonic and adult glycogen, because the well-known postnatal glycogenoses are not associated with gross morphological malformations.

The genetic prototypes of thalidomide embryopathy lend themselves to speculations along the same lines. Thalidomide may be conceived of as an inhibitor of an enzyme engaged in the mobilization of embryonic glycogen. The genetic prototypes would then possibly affect the same metabolic chain at the same or at adjacent links.

REFERENCES

- Aase, J. M. and Smith, D. W. (1969). Congenital anemia and triphalangeal thumbs. A new syndrome. *Journal of Pediatrics*, **74**, 471–474.
- Abramowitz, I. (1959). Triphalangeal thumb in a Bantu family. *Journal of Bone and Joint Surgery*, **41B**, 766–771.
- Ancel, P. (1950). *La Chimiotératogénèse chez les Vertébrés. Réalisation des Monstruosités par des Substances chimiques.* (Encyclopédie scientifique. Bibliothèque de Biologie Générale.) Gaston Doin et C^{ie}, Paris.
- Atwood, E. and Pond, C. P. (1917). A polydactylous family. *Journal of Heredity*, **8**, 96–98.
- Bätke, W. (1964). *Missbildungen bei der Fanconi-Panmyelopathie.* Inaugural dissertation, Hamburg.
- Barsky, A. J. (1951). Congenital anomalies of the hand. *Journal of Bone and Joint Surgery*, **33A**, 35–64.
- Barsky, A. (1958). *Congenital Anomalies of the Hand and their Surgical Treatment.* Thomas, Springfield, Illinois.
- Bierich, J. R. (1965). Ätiopathogenese und klinisches Bild hypotalamischer und hypophysärer Wachstumsstörungen. *Monatsschrift für Kinderheilkunde*, **113**, 269–277.
- Birch-Jensen, A. (1949). *Congenital Deformities of the Upper Extremities.* Andelsbogtrykkeriet i Odense.
- Blencke, D. (1904). Ein weiterer Beitrag zur sogenannten Klump-hand. *Zeitschrift für orthopädische Chirurgie*, **13**, 654–657.
- Bodenstein, D. and Abdel-Malek, A. (1949). The induction of aristopedia by nitrogen mustard in *Drosophila virilis*. *Journal of Experimental Zoology*, **111**, 95–115.

- Bresch, C. and Hausmann, R. (1970). *Klassische und molekulare Genetik*, 2nd ed. Springer, Berlin, Heidelberg, and New York.
- Büchner, F. (1958). Die Bedeutung peristatischer Faktoren für die Entstehung der Missbildungen und Missbildungsstörungen. *Verhandlungen der Deutschen Gesellschaft für innere Medizin*, **64**, 13–33.
- Cotte, G. (1923). Pouce bot varus congénital. *Revue d'Orthopédie*, **10**, 411–413.
- Degenhardt, K.-H. (1958). Aufgaben der Phänokopie-Forschung. *Deutsche Medizinische Wochenschrift*, **83**, 1085–1088.
- Demarinis, F. and Wildervanck, L. S. (1960). Pre-axiale polydactylie (verdubbelde duim) en trifalangie. *Nederlandsch Tijdschrift voor Geneeskunde*, **104**, 2169–2171.
- Dyban, A. P. (1962). *Grundriss der pathologischen Embryologie des Menschen*. G. Fischer, Jena.
- Eaton, G. O. and McKusick, V. A. (1969). A seemingly unique polydactyly-syndactyly syndrome in four persons in three generations. *Birth Defects: Original Article Series*, **5**, pt. 1, pp. 221–225. National Foundation—March of Dimes, New York.
- Ecke, H. (1962). Beitrag zu den Doppelmisbildungen im Bereich der Finger. *Brun's Beiträge zur klinischen Chirurgie*, **205**, 463–468.
- Ferber, C. (1953). Ein Beitrag zur Dreigliedrigkeit des 'Daumens'. *Zeitschrift für Orthopädie und ihre Grenzgebiete*, **83**, 55–64.
- Ferrell, R. L., Jones, B., and Lucas, R. V., Jr. (1966). Simultaneous occurrence of the Holt-Oram and the Duane Syndromes. *Journal of Pediatrics*, **69**, 630–634.
- Fischer, E. (1895). *Transmutation der Schmetterlinge infolge Temperaturänderungen*. R. Friedländer, Berlin.
- Frere, J. M. (1960). A case having thumbs with three phalanges simulating fingers. *Southern Medical Journal*, **23**, 536.
- Gasser, R. F. (1967). The development of the facial muscles in man. *American Naturalist*, **120**, 357–376.
- Gerold, M. (1959). Frakturheilung bei einem seltenen Fall kongenitaler Anomalie der oberen Gliedmassen. *Zentralblatt für Chirurgie*, **84**, 831–834.
- Ghiari, M. (1972). *Kaudales Regressions-Syndrom bei Kindern diabetischer Mütter*. Inaugural dissertation, Münster.
- Gifford, H. (1926). Congenital defects of abduction and other ocular movements and their relation to birth injuries. *American Journal of Ophthalmology*, **9**, 3–22.
- Giobus, M. and Gibson, M. A. (1968). A histological and histochemical study of the development of the sternum in thalidomide-treated rats. *Teratology*, **1**, 235–256.
- Gloor, H. (1947). Phänokopie-Versuche mit Aether an Drosophila. *Revue Suisse de Zoologie*, **54**, 637–712.
- Goldschmidt, R. (1935). Gen und Aussencharakter. III. *Biologisches Zentralblatt*, **55**, 535–554.
- Goldschmidt, R. and Pitternick, L. K. (1957). The genetic background of chemically induced phenocopies in Drosophila. *Journal of Experimental Zoology*, **135**, 127–202.
- Grebe, H. (1954). Gene und Phänotypen in der Ätiologie menschlicher Missbildungen. *Acta Geneticae Medicae et Gemellologiae*, **3**, 197–209.
- Gruber, W. (1865). Ueber congenitalen Radiusmangel. *Archiv für Pathologie, Anatomie und Physiologie*, **32**, 211–228.
- Haas, S. L. (1939). Three-phalangeal thumbs. *American Journal of Roentgenology Radium Therapy*, **42**, 677–682.
- Hadorn, E. (1955). *Letalfaktoren in ihrer Bedeutung für Erbpathologie und Genphysiologie der Entwicklung*. Thieme, Stuttgart.
- Harnack, G. A. von, Horst, W., and Lenz, W. (1961). Das erbliche Syndrom: Innenohrschwerhörigkeit und Jodfehlverwertung mit Kropf. *Deutsche medizinische Wochenschrift*, **86**, 2421–2428.
- Hefner, R. A. (1940). Hereditary polydactyly associated with extra phalanges in the thumbs. *Journal of Heredity*, **31**, 25–27.
- Henke, K., Finck, E. von, and Ma, S.-Y. (1941). Über sensible Perioden für die Auslösung von Hitzemodifikationen bei Drosophila und die Beziehungen zwischen Modifikationen und Mutationen. *Zeitschrift für Vererbungslehre*, **79**, 267–316.
- Hill, L. (1937). Congenital abnormalities—phocomelus and congenital absence of radius. *Surgery, Gynecology and Obstetrics*, **65**, 475–479.
- Homi, C. (1920). A case of phocomelus. *Lancet*, **2**, 128.
- Hopf, A. (1959). Die angeborenen Veränderungen des Unterarmes und der Hand. In *Handbuch der Orthopädie*, vol. 3, pp. 419–506. Thieme, Stuttgart.
- Kersting, D. (1966). *Eine klinisch-erbbiologische Untersuchung der Hyperphalangie*. Inaugural dissertation, Münster.
- Klemm, F. W. (1956). Dolichophalangie mit echter Hyperphalangie. *Fortschritte der Röntgenstrahlen*, **85**, 256–257.
- Komai, T., Ozaki, Y., and Inokuma, W. (1953). A Japanese kindred of hyperphalangism of thumbs and duplication of thumbs and big-toes. *Folia Hereditaria et Pathologica*, **2**, 307–312.
- König, M. P. (1968). *Die kongenitale Hypothyreose und der endemische Kretinismus*. Springer, Berlin.
- Kreybig, T. von (1968). *Experimentelle Praenataltoxikologie*. Cantor, Aulendorf/Württemberg.
- Landauer, W. (1957). Phenocopies and genotype, with special reference to sporadically-occurring developmental variants. (With a discussion by R. B. Goldschmidt.) *American Naturalist*, **91**, 79–94.
- Landauer, W. (1958). On phenocopies, their developmental physiology and genetic meaning. *American Naturalist*, **92**, 201–213.
- Landauer, W. (1959). The phenocopy concept: Illusion or reality? *Experientia*, **15**, 409–412.
- Lapidos, P. W., Guidotti, F. P., and Coletti, C. J. (1943). Triphalangeal thumb. *Surgery, Gynecology and Obstetrics*, **77**, 178–186.
- Lelling, E. (1938). *Zum Thema der Verbindung der Extrauterin-gravidität mit Missbildungen*. Inaugural dissertation, Halle-Wittenberg.
- Lenz, W. (1963). Das Thalidomid-Syndrom. *Fortschritte der Medizin*, **81**, 148–155.
- Lenz, W., Theopold, W., and Thomas, J. (1964). Triphalangie des Daumens als Folge von Thalidomidschädigung. *Münchener Medizinische Wochenschrift*, **106**, 2033–2041.
- McKusick, V. A. (1971). *Mendelian Inheritance in Man. Catalogs of Autosomal Dominant, Autosomal Recessive, and X-linked Phenotypes*, 3rd edition. The Johns Hopkins Press, Baltimore.
- Manoiloff, E. O. (1931). A rare case of hereditary hexadactylism. *American Journal of Physical Anthropology*, **15**, 503–508.
- Matzen, P.-F. and Fleissner, H. K. (1969). *Orthopädischer Röntgenatlas*. G. Thieme, Stuttgart.
- Mennerich, P. (1923). Ein Fall von Retraktionsbewegungen der Augen bei angeborenen Anomalien der äusseren Augenmuskeln. *Zeitschrift für Augenheilkunde*, **50**, 173–180.
- Müller, W. (1936). Beiträge zur Kenntnis des dreigliedrigen Daumens. *Archiv für klinische Chirurgie*, **185**, 377–386.
- Müller, W. (1937). Die angeborenen Fehlbildungen der menschlichen Hand. In *Erb- und Konstitutionsbiologie der Hand*. G. Thieme, Leipzig.
- Nachtsheim, H. (1957). Mutation und Phänokopie bei Säugetier und Mensch. Ihre theoretische und praktische Bedeutung für Genetik und Eugenik. *Experientia*, **13**, 57–96.
- Nachtsheim, H. (1961). Kritische Betrachtungen zu einigen modernen Begriffen der Genetik: Phänokopie und Genokopie, Embryopathie und Genopathie. *Archiv der Julius-Stiftung*, **36**, 70–89.
- Nudleman, K. L. and Travill, A. A. (1971). A morphological and histochemical study of thalidomide-induced upper limb malformations in rabbit fetuses. *Teratology*, **4**, 409–426.
- Nylander, E. S. (1931). Präaxiale Polydaktylie in fünf Generationen einer schwedischen Sippe. *Uppsala Läkareförenings Förhandlingar*, new series, **36**, 275–292.
- Partsch, C.-J. (1969). Angeborene Ohrmissbildung als Ursache der Taubheit. *Annales Universitatis Saraviensis*, **16**, 228–349.
- Polinelli, U. (1962). Un caso di iperfalangia familiare dei pollici. *Minerva Nipologica*, **12**, 373–375.
- Rapoport, J. A. (1939). Specific morphosis in Drosophila induced by chemical compounds. *Bulletin of Biology of the U.S.S.R.*, **7**, 414–417.
- Reber, M. (1967/68). Un syndrome osseux peu commun associant une heptadactylie et une aplasie des tibias. *Journal de Génétique Humaine*, **16**, 15–39.
- Rieder, H. (1900). Eine Familie mit dreigliedrigen Daumen. *Zeitschrift für Morphologie und Anthropologie*, **2**, 117–197.
- Rimoin, D. L., Merimee, T. J., Rabinowitz, D., and McKusick, V. A. (1968). Genetic aspects of clinical endocrinology. *Recent Progress in Hormone Research*, **24**, 365–437.
- Rimoin, D. L. and Schimke, R. N. (1971). *Genetic Disorders of the Endocrine Glands*. C. V. Mosby, St Louis.
- Roberts, J. B. (1919). A child with double cleft lip and palate protrusion of the intermaxillary portion of the upper jaw and imperfect development of the bones of the four extremities. *Annals of Surgery*, **70**, 252–253.

- Roberts, E. (1943). Hereditary hyperphalangism of the thumb. *Journal of Heredity*, **34**, 290–292.
- Rodenstein, L. A. (1875/76). Case of arrest of development of both upper extremities. *American Journal of Obstetrics*, **8**, 663–664.
- Sallam, A. M. (1955). Triphalangeal thumbs. *Archives of Surgery*, **71**, 257–259.
- Salzer, H. (1898). Zwei Fälle von dreigliedrigem Daumen. *Anatomischer Anzeiger*, **14**, 124–131.
- Sang, J. H. and McDonald, J. M. (1954). Production of phenocopies in *Drosophila* using salts, particularly sodium metaborate. *Journal of Genetics*, **52**, 392–412.
- Silver, H. K. and Finkelstein, M. (1967). Deprivation dwarfism. *Journal of Pediatrics*, **70**, 317–324.
- Ströer, W. F. H. (1936). Familiäres Auftreten von Reihen erblicher Hand- und Fussabweichungen. *Der Erbarzt*, **3**, 22–26.
- Ströer, W. F. H. (1939). Über das Zusammentreffen von Hasenscharte mit ernstesten Extremitätenmissbildungen. *Der Erbarzt*, **7**, 101–104.
- Swanson, A. B. and Brown, K. S. (1962). Hereditary triphalangeal thumb. *Journal of Heredity*, **53**, 259–265.
- Temtamy, S. (1966). *Genetic Factors in Hand Malformations*. PhD Thesis, The Johns Hopkins University.
- Villumsen, A. L. (1970). *Environmental Factors in congenital Malformations. A Prospective Study of 9,006 Human Pregnancies*. F.A.D.L.s Forlag, Copenhagen, Aarhus and Odense.
- Vrolik, W. (1849). *Tabulae ad illustrandum embryogenesis hominis et mammalium*. G. M. P. Londonck, Amsterdam.
- Wildervanck, L. S. (1954). Drie erfelijke afwijkingen van de bovenste extremiteiten. *Nederlandsche Tijdschrift voor Geneeskunde*, **98**, 1681–1685.
- Winckel, F. von (1902). *Über die Missbildungen von ektopisch entwickelten Früchten und deren Ursachen*. J. F. Bergmann, Wiesbaden.
- Zeleny, C. (1923). The temperature coefficient of a heterozygote. *Biological Bulletin*, **44**, 105–112.