Haemoglobin D Punjab (D Los Angeles)

F. VELLA* and H. LEHMANN†

Summary. A search for haemoglobin variants undertaken in Canada revealed 21 unrelated instances of Hb D Punjab amongst 207,300 specimens tested. Of these, eight came from East Indian immigrants and the rest from Canadians of United Kingdom origin. No instances of Hb D Punjab were found in 14,500 specimens from Canadian Indians that were tested. The geographical origins of 27 instances of Hb D Punjab characterized at the MRC Abnormal Haemoglobin Unit, Cambridge are presented. Of these five were natives of the British Isles. The results of surveys undertaken in the United Kingdom are summarized. The global distribution of Hb D Punjab is discussed.

Haemoglobin D (Hb D) was the third abnormal haemoglobin to be reported (Itano, 1951; Sturgeon, Itano, and Bergren, 1955). It was discovered in a family of mixed British and American-Indian origin from the Los Angeles area, of which some members had a haemolytic anaemia with a tendency for the erythrocytes to sickle slowly on reduction. The characteristics of Hb D were: (1) an electrophoretic mobility identical to that of Hb S at alkaline pH, (2) a normal solubility when in the reduced state, and (3) absence of sickling of erythrocytes which contain it in association with Hb A. The Hb D in this family has been characterized as $\alpha_2\beta_2^{121 \text{ Glu} \rightarrow \text{Gln}}$ and called D Los Angeles (Babin, Jones, and Schroeder, 1964).

Between the discovery of Hb D and the elucidation of its molecular structure, abnormal haemoglobins with the same properties were found to occur in many ethnic groups. Thus, they were reported in white persons of European extraction (Dacie, 1954; White and Beaven, 1954; Chernoff, 1958) as well as in English (Stewart and MacIver, 1956; Smith and Conley, 1959), Portuguese (Arends, Layrisse, and Rincon, 1959), Italian (Bowman and Ingram, 1961), German (Martin *et al.*, 1960), Greek (Gouttas *et al.*, 1960), Turkish (Aksoy and Lehmann, 1956; Gammack *et al.*, 1961), Persian (Hynes and Lehmann, 1956), Indian (Bird and Lehmann, 1956; Jacob, Lehmann, and Raper, 1956), as well as

American negro subjects (Chernoff, 1956; McCurdy, 1959; Marder and Conley, 1959). Five specimens of Hb D of widely differing ethnic origin—namely Hb D Chicago (Bowman and Ingram, 1961); Hb D North Carolina (Smith and Conley, 1959); Hbs D Punjab, D Portugal, and D Cyprus supplied by Dr H. Lehmann—were all shown to have the composition $\alpha_2\beta_2^{121 \text{ Glu} \rightarrow \text{Gln}}$ (Baglioni, 1962). These findings demonstrated that this variant has a wide distribution. It was soon demonstrated that the electrophoretic and solubility properties of Hb D were by no means specific for Hb D Punjab by the discovery of many instances of Hb D which have a different molecular structure from Hb D Punjab. For example, D Baltimore (Marder and Conley, 1959), D Washington (McCurdy *et al.*, 1961), D St Louis (Minnich *et al.*, 1962), G Azuakoli (Lehmann and Nwokolo, 1959), G Bristol (Raper *et al.*, 1960), G Philadelphia (Atawater, Schwartz, and Tocantins, 1960) all of which have the structure $\alpha_2\beta_2^{68 \text{ Asn} \rightarrow \text{Lys}}$ (Baglioni and Ingram, 1961; Weatherall, Sigler, and Baglioni, 1962; Dance, Huehns, and Shooter, 1964) as do also $\text{G}_{\text{St}-1}$ (Bowman *et al.*, 1966) and G Knoxville (Chernoff and Pettit, 1965); while G Galveston is $\beta^{43 \text{ Glu} \rightarrow \text{Ala}}$ (Bowman *et al.*, 1964); D Ibadan is $\beta^{87 \text{ Thr} \rightarrow \text{Lys}}$ (Watson-Williams *et al.*, 1965); D Bushman is $\beta^{16 \text{ Gly} \rightarrow \text{Arg}}$ (Wade, Jenkins, and Huehns, 1967).

Some confusion arose from the different behaviour on starch gel electrophoresis and paper electrophoresis at pH 8.6 of some variants. Hb D Punjab moves in the same manner as Hb S on both techniques. Certain haemoglobins, however, move slightly faster than Hb S on paper electrophoresis

Received 14 February 1974.

* Department of Biochemistry, University of Saskatchewan, Saskatoon, Sask, Canada. On sabbatical leave at †.

† MRC Abnormal Haemoglobin Unit, University Department of Biochemistry, Cambridge, United Kingdom.

particularly if barbitone buffer is used and these were named Hb G, but on starch gel electrophoresis they behave like Hb D. Depending on which medium research workers used for electrophoresis they named the haemoglobin variant D or G.

This paper describes the occurrence of Hb D Punjab in Canada and in the United Kingdom and reviews its global distribution.

Hb D Punjab in Canada

Between July 1965 and April 1973, some 222,000 blood samples obtained from blood donors or from hospital patients in various parts of the four Canadian provinces of Ontario, Manitoba, Saskatchewan, and Alberta were screened during a deliberate search for haemoglobin variants that have a different net charge from that of normal haemoglobin. For this purpose, a filter-paper electrophoresis technique was used (TEB—barbiturate discontinuous buffer system, pH 8.9; Vella, 1967). Twenty-nine of these specimens were found to contain an abnormal haemoglobin fraction which had the following properties: (1) it accounted for 30–40% of the total pigment, (2) it had the electrophoretic mobility of Hb S at pH 8.9, (3) it did not separate from Hb A in citrate-agar gel electrophoresis at pH 6.1 (Robinson *et al*, 1957), (4) it was associated with a single haemoglobin A₂ band of normal concentration, and with normal amounts of alkali-resistant haemoglobin (0.5–1.5% by the one minute alkali denaturation method; Singer, Chernoff, and Singer, 1951), (5) it showed a normal solubility when reduced in 2.24 M phosphate buffer (Itano, 1953), and (6) it was not associated with sickling of erythrocytes when these were treated with metabisulphite. All these subjects were considered to be heterozygotes for an Hb D gene. In 21 of them, the abnormal haemoglobin was isolated, purified, and characterized in Saskatoon on the basis of fingerprint patterns of tryptic digests by methods that have already been described (Vella *et al*, 1967b). Confirmatory results were obtained on three of these subjects by fingerprinting and by amino-acid analysis of isolated abnormal peptides undertaken at Cambridge. In all cases, the fingerprints were indistinguishable from those of Hb D Punjab while the amino-acid analysis of aberrant peptides confirmed this characterization.

The incidence of Hb D Punjab in the four provinces is summarized in Table I. Eight of these subjects were of East Indian origin, while the remainder were white Canadians of British stock. The overall frequency appears to be 1/10,000 of the

TABLE I
HAEMOGLOBIN D PUNJAB IN CANADA*

Province	No. of Samples	D Punjab	
		Total	No. in East Indians
Ontario	14,700	3	1
Manitoba	84,500	5	1
Saskatchewan	81,400	9	3
Alberta	26,700	4	3
Total	207,300	21	8

* No instance of this variant was encountered in 14,500 blood samples from persons of Canadian Indian origin.

population tested. If only the instances in white subjects are considered, the overall frequency becomes 1/16,000 (ranging between 1/7,000 in Ontario and 1/27,000 in Alberta).

Hb D Punjab in the United Kingdom

The first instances of Hb D in English persons were reported by Stewart and MacIver (1956) from Jamaica, and Smith and Conley (1959) from North Carolina. The haemoglobin of the family reported by the latter was characterized as $\beta 121 \text{ Glu} \rightarrow \text{Gln}$ by Baglioni (1962). Other instances of Hb D Punjab in English persons abroad have been reported from Ghana (Ringelmann *et al*, 1967) and Australia (Ungar *et al*, 1973).

Several extensive surveys have been undertaken in recent years to determine the incidence of abnormal haemoglobin variants in the United Kingdom. The frequency of Hb D Punjab revealed by these is summarized in Table II. The first instance to be detected was that in an East Anglian family (Huntsman *et al*, 1963; Konigsberg *et al*, 1965). Black (1969) found four instances amongst some 8,000 newborn English infants that he tested. However, other surveys have given only negative results and the overall frequency in British persons appears to be about 1 in 7,500. This is in good agreement with the frequency suggested by the Canadian results. The incidence is much higher amongst recent immigrants to Britain, especially amongst those of Indian origin. A recent survey amongst some 7000 immigrant school children in Birmingham showed Hb D to be present in 0.54% (Stuart *et al*, 1973).

During the last nine years, 27 instances of Hb D Punjab have been characterized at the MRC Abnormal Haemoglobin Unit in Cambridge. The geographical origin of these is shown in Table III.

TABLE II
INCIDENCE OF Hb D PUNJAB IN SURVEYS IN
THE UNITED KINGDOM

Source	No.		References
	Examined	D Punjab	
British servicemen in Singapore and Malaysia	4,387	0	Vella (1961/1962)
Norfolk	1,000	1	Huntsman <i>et al</i> (1963) Konigsberg <i>et al</i> (1965)
Oxford and Peterborough	1,971	0	Liddell <i>et al</i> (1964)
Cambridge	3,000	0	Sick <i>et al</i> (1967)
London and other English cities*	10,000	6†	Black (1969)
London‡	10,000	0	F. E. Boulton and R. G. Huntsman (personal communication)
Scotland	3,968	0	Cook and Lehmann (1973)
Ireland	2,500	0	D. Tills (personal communication)
Total	36,826	7	

* Cord blood survey.

† Includes two instances among East Indian immigrants.

‡ Myoglobin survey. (This survey did detect some instances of Hb S and C.)

TABLE III
HAEMOGLOBIN D PUNJAB
CHARACTERIZED AT
CAMBRIDGE
(unpublished cases only)

Source of Specimens	D Punjab	
	Total	No. in East Indians
England	11	6
Canada	5	2
Australia	1	—
South Africa	2	1
Lebanon	2	—
Italy	1	—
Switzerland	2*	—
France	1	—
Portugal	1	—
Venezuela	1	—

* Both Italians.

Global incidence and distribution

The distribution and incidence of Hb D Punjab is summarized in Table IV. While single instances of this variant have been reported in persons originating from various parts of Europe, the incidence increases appreciably from the Eastern Mediterranean to the western part of the Indian sub-continent, and especially in Iran, the Punjab, and the western half of India where between 1–2% may have the variant in their blood. It has been reported once from Thailand (Wasi *et al*, 1968) and has been found in a Batak Indonesian from Medan (L. E. Lie-Injo and J. Clegg, personal communication), but does not appear to have been encountered during extensive surveys in Taiwan (Blackwell and Liu, 1970; Blackwell *et al*, 1972) and Japan (Shibata, Iuchi, and Miyaji, 1966).

The spread of haemoglobin D Punjab to its present day geographical pattern may be explained on the basis of population migration occurring in two phases. The first phase would cover the period of the appearance of the mutation and its distribution to northern India, Iran, and Turkey and may reflect the pattern of migrations that occurred in that region during the times of Darius I of Persia or later during the Mogul invasions. The second would cover the spread of the gene to Western Europe and to the New World and may have resulted from intermarriage between Europeans and Indians over the last two or three centuries. Konigsberg *et al* (1965) have examined the evidence that this variant was introduced into Britain from the Punjab as a result of intermarriage of serving British troops with the Indian population during the 18th century. They concluded that the occasional examples of D Punjab found during surveys in Europeans (and especially in Britain, France, and Portugal) confirm the bonds that existed between these countries and India between the 17th and 20th centuries.

Though the earlier literature on haemoglobin variants contains many reports on the occurrence of Hb D in negro subjects, especially in America (Chernoff, 1956; Chernoff and Weichselbaum, 1958; Marder and Conley, 1959; Myerson, Harrison, and Lohmuller, 1959; McCormick, 1960; Cawein *et al*, 1966) only in a very few cases has this been identified as D Punjab (Rothman and Ranney, 1971; P. R. McCurdy and H. Lehmann, unpublished results). In fact the commonest variant with the properties of Hb D in American negroes is Hb G Philadelphia, and this appears to be widely distributed since it has also been found in the Congo (Dherte *et al*, 1959), in Nigeria (Lehmann and Nwokolo, 1959), and in the British West Indies

Eleven of them had come from England. However, only five had come from English persons (two from Leeds and three from London) while the remainder came from Indian immigrants.

TABLE IV
INCIDENCE AND DISTRIBUTION OF HAEMOGLOBIN D PUNJAB

Country	No.		Reference	Reference to Isolated Instances
	Examined	D Punjab		
United Kingdom	36,826	7	Table II	Ringelmann <i>et al</i> (1967); Schneider <i>et al</i> (1968)
Portugal	3042	0	Trincao and Ferreira (1962)	Table III; Preto <i>et al</i> (1961); Baglioni (1962)
France	—	—	—	Table III; Labie and Rosa (1969); Wajcman <i>et al</i> (1969)
Holland	991	1	De Jong and Went (1968)	—
Denmark	5000	1	Sick <i>et al</i> (1967)	—
Norway	3000	1	Monn <i>et al</i> (1968)	—
Sweden	4171	0	Nilsson and Eriksson (1972)	—
Finland	4418	0	Nilsson and Eriksson (1972)	—
Russia	317	0	Nilsson and Eriksson (1972)	—
Germany	—	—	—	Martin <i>et al</i> (1960); Kohne <i>et al</i> (1972)
Switzerland	—	—	—	Table III
Italy	13,187 3000	0 0	Quattrin <i>et al</i> (1966) Licenziati (1969)	Table III; Bowman and Ingram (1961); Baglioni (1962); Ventruto <i>et al</i> (1963); Cortesi <i>et al</i> (1966); Ricco <i>et al</i> (1970); Rossi and Morelli (1970)
Bulgaria	—	—	—	Cholakov and Kanchev (1970)
Yugoslavia	2860	2	Sadikario <i>et al</i> (1969)	—
Greece	1000	0	Fessas (1959)	R. Rieder (personal communication)
Turkey	—	—	—	Baglioni (1962); Özsoylu (1970)
Lebanon	2938	0	Cabannes <i>et al</i> (1965)	Table III
Israel	397 190	0 0	Dreyfus and Pinkhas (1958) Beaven (1973)	Ramot <i>et al</i> (1969)
Iran	400 200 184	1 2 3	Rahbar <i>et al</i> (1967) Rahbar <i>et al</i> (1967) Lehmann <i>et al</i> (1973)	—
Indian sub-continent	1000 (Bengal)	1	Chatterjea (1959)	Table I; Table II
	129 (W. Pakistan)	1	Stern <i>et al</i> (1968)	Table III; Baglioni (1962); Marengo-Rowe <i>et al</i> (1968)
Thailand	2790	0	Flatz <i>et al</i> (1965)	Wasi <i>et al</i> (1968)
Indonesia	5400	0	Lie-Injo (1959)	L. E. Lie-Injo and J. Clegg (personal communication)
Taiwan	100,000	0	Blackwell and Liu (1970)	—
Japan	130,000	0	Shibata <i>et al</i> (1966)	—
Australia	—	—	—	Ungar <i>et al</i> (1973)
South Africa	3125	0	Botha and van Zyl (1966)	Table III
Canada	207,300	21	Table I	Table III
Greenland	153	0	Nilsson and Eriksson (1972)	—
Iceland	1000	0	Nilsson and Eriksson (1972)	—
USA	—	—	—	Stout <i>et al</i> (1964; see Arends, 1966); Rothman and Ranney (1971); R. R. McCurdy and H. Lehmann (unpublished results); Imamura and Riggs (1972)
Jamaica	10,000	0	Milner (1967)	—
	3449	0	Ahern <i>et al</i> (1973)	—
Venezuela	1666	0	Arends (1963)	Table III; Arends (1971)
Mexico	2336	0	Lisker <i>et al</i> (1966); Lisker (1971)	Schneider <i>et al</i> (1968); Bello <i>et al</i> (1969)

(Dance *et al*, 1964). Huisman (1969) has found 13 alpha-chain variants with the electrophoretic mobility of Hb D or G but only three beta-chain variants with a similar mobility amongst 8500 negroes from Georgia (USA) examined. Also frequent in negroes originating in West Africa is Hb Korle-Bu ($\beta 73 \text{ Asp} \rightarrow \text{Asn}$) which has been reported from Ghana (Konotey-Ahulu *et al*, 1968) as well as Martinique (Bookchin, Nagel, and Ranney, 1970). This variant is identical with Hb G Accra (Lehmann, Beale, and Boi-Doku, 1964; Boi-Doku, Kinderlerer, and Lehmann, 1972) and is the commonest form of Hb D or G so far found in the British West Indies (Milner, 1967; Ahern, Swan, and Ahern, 1973). It has been suggested that the Hb D found amongst negro subjects in South America (Venezuela—Arends, 1961; Colombia—Restrepo and Londoño, 1965; Brazil—de Araujo and Jamra, 1966) may have originated from the Iberian peninsula (de Araujo and Jamra, 1966). However, the Hb D found in a Colombian negro family (Restrepo and Londoño, 1965) has been identified as Hb Korle-Bu (Restrepo, 1971). The Hb D reported from some parts of Portuguese Africa (Trinçao, 1967; David and Trinçao, 1963) has not been characterized.

Hb D Punjab has not been reported amongst Arab populations in North Africa. It has been suggested that the Hb D that is present at appreciable frequency in the Arabo-Berbers of Algeria (Cabannes, 1965) may in fact be an alpha-chain variant (de Traverse and Coquelet, 1961; Konigsberg *et al*, 1965). More recently, however, two new variants of Hb D have been reported in the Algerian population, one Hb D Ouled Rabah ($\beta 19 \text{ Asn} \rightarrow \text{Lys}$) and the other D Iran ($\beta 22 \text{ Glu} \rightarrow \text{Gln}$) (Elion *et al*, 1973) the latter having also been reported in an Iranian (Rahbar, 1973) and in a Pakistani family (Rohe, Sharma, and Ranney, 1973).

A variant with the electrophoretic properties of Hb D has been found amongst some groups of American Indians in the USA (Pollitzer *et al*, 1959; Githens *et al*, 1961) but not in South America (Colombia, Venezuela, British Guiana—Arends, 1963; de Pinango and Arends, 1965; Arends, 1971, Restrepo, 1971; Mexico—Lisker, Zarate, and Loria, 1966, Lisker, 1971; Brazil—Salzano and Tondo, 1968; de Araujo, 1971; San Salvador—Bloch and Rivera, 1969). Arends (1966) has summarized the results of haemoglobin studies of Indians of the American continent. The only variant with the properties of Hb D that was found in Canadian Indians in Canada is Hb G Coushatta which was reported in a large family in Saskatchewan (Vella, Isaacs, and

Lehmann, 1967a; Vella and Guzak, 1968). It has since been found in 26 other Canadian Indians (12 each in Manitoba and Alberta and two in Ontario). This variant was originally discovered in the Alabama Coushatta Indians of Texas (Schneider *et al*, 1964; Bowman, Barnett, and Hite, 1967) and has more recently been reported in Korean and Chinese subjects (Blackwell *et al*, 1969).

Addendum

The following reported instances of Hb D Punjab were inadvertently omitted from Table IV: Greece (Deliyannis, Ballas, and Christakis, 1969) and Cuba (Uriarte, Perez Atencio, and Colombo, 1973).

REFERENCES

- Deliyannis, G. A., Ballas, A., and Christakis, I. (1969). Haemoglobin D in a Greek family. *Acta Haematologica*, **41**, 121–125.
- Uriarte, A., Peres Atencio, R., and Colombo, B. (1973). Haemoglobin D Punjab in a Cuban family and its interaction with haemoglobin S. *Acta Haematologica*, **50**, 315–320.
- We are grateful to Drs F. E. Boulton, J. B. Clegg, R. G. Huntsman, L. E. Lie-Injo, R. Rieder, and D. Tills for permission to quote their unpublished data.
- F. V. is grateful to A. Labossiere and Mrs M. Hurley for technical assistance and to the Medical Research Council of Canada for a grant.
- ### REFERENCES
- Ahern, E. J., Swan, A. V., and Ahern, V. N. (1973). The prevalence of the rarer inherited haemoglobin defects in adult Jamaicans. *British Journal of Haematology*, **25**, 437–444.
- Aksoy, M. and Lehmann, H. (1956). A further example of haemoglobin D in a Turkish family. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, **50**, 178–180.
- Araujo, J. T. de (1971). Geographical distribution and incidence of hemoglobin variants in Brazil. In *Genetical, Functional and Physical Studies of Hemoglobins*, ed. by T. Arends, G. Bemski, and R. L. Nagel, pp. 26–31. Karger, Basel.
- Araujo, J. T. de and Jamra, M. (1966). Incidence of abnormal hemoglobins in a sample of the Sao Paulo city population. *Blood*, **28**, 495.
- Arends, T. (1961). Frecuencia de las hemoglobinas anormales en Venezuela. *Archivos del Hospital Vargas*, **3**, 225–236.
- Arends, T. (1963). Search for abnormal hemoglobins in South American Indians living in malarious zones. In *Proceedings of the 7th International Congress on Tropical Medicine and Malaria*, **4**, 179–182. National Institute of Allergy and Infectious Diseases, Rio de Janeiro.
- Arends, T. (1966). Haemoglobinopathies, thalassaemia and glucose-6-phosphate dehydrogenase deficiency in Latin America and the West Indies. *New Zealand Medical Journal* (Haematology suppl.), **65**, 831–844.
- Arends, T. (1971). Epidemiology of hemoglobin variants in Venezuela. In *Genetical, Functional and Physical Studies of Hemoglobins*, ed. by T. Arends, G. Bemski, and R. L. Nagel, pp. 82–98. Karger, Basel.
- Arends, T., Layrisse, M., and Rincon, A. R. (1959). Sick cell-haemoglobin D disease in a Portuguese child. *Acta Haematologica*, **22**, 118–126.
- Atwater, J., Schwartz, I. R., and Tocantins, L. M. (1960). A variety of human hemoglobin with four distinct electrophoretic components. *Blood*, **15**, 901–908.
- Babin, D. R., Jones, R. T., and Schroeder, W. A. (1964). Haemoglobin D Los Angeles: $\alpha_2\beta_2^{121} \text{ Glu NH}_2$. *Biochimica et Biophysica Acta*, **86**, 136–143.
- Baglioni, C. (1962). Abnormal human haemoglobins. VII. Chemical studies on haemoglobin D. *Biochimica Biophysica Acta*, **59**, 437–449.

- Baglioni, C. and Ingram, V. M. (1961). Abnormal human haemoglobins. V. Chemical investigation of haemoglobins A, G, C, X from one individual. *Biochimica et Biophysica Acta*, **48**, 253-265.
- Beaven, G. H. (1973). Haemoglobin studies of Yemenite and Kurdish Jews in Israel. *Philosophical Transactions of the Royal Society of London, Series B*, **266**, 185-193.
- Bello, A., Dorantes, S., and Alvarez, C. (1969). Hemoglobin D Punjab en una familia Mexicana, con dos casos clinicos de hemoglobinopatia S-D. *Revista de Investigación Clínica*, **21**, 489-496.
- Bird, G. W. G. and Lehmann, H. (1956). Haemoglobin D in India. *British Medical Journal*, **1**, 514.
- Black, A. J. (1969). *A Survey of Umbilical Cord Blood for Haemoglobin Variants*. MD Thesis, Cambridge University.
- Blackwell, R. Q. and Liu, C. S. (1970). Hemoglobin G Taichung: $\alpha^{74}\text{Asp} \rightarrow \text{His}$. *Biochimica et Biophysica Acta*, **200**, 70-75.
- Blackwell, R. Q., Ro, I. H., Liu, C. S., Yang, H. J., Wang, C. C., and Huang, J. T. S. (1969). Hemoglobin variant found in Koreans, Chinese, and North American Indians: $\alpha\beta\beta_2$ ($^{22}\text{Glu} \rightarrow \text{Ala}$). *American Journal of Physical Anthropology*, **30**, 389-391.
- Blackwell, R. Q., Weng, M. I., Liu, C. S., Shih, T. B., and Wang, C. L. (1972). Hemoglobin G Chinese in Chinese subjects in Taiwan. *Vox Sanguinis*, **23**, 363-368.
- Bloch, M. and Rivera, H. (1969). Hemoglobinas anormales y deficiencia de glucosa-6-fosfato-dehidrogenasa en El Salvador. *Sangre*, **14**, 121-124.
- Boi-Doku, F. S., Kinderlerer, J., and Lehmann, H. (1972). In *Atlas of Protein Sequence and Structure*, ed. by M. Dayhoff, p. 73. National Biomedical Research Foundation, Silver Springs.
- Bookchin, R. M., Nagel, R. L., and Ranney, H. M. (1970). The effects of $\beta^{73}\text{Asn}$ on the interactions of sickling hemoglobin. *Biochimica et Biophysica Acta*, **221**, 373-375.
- Botha, M. C. and Zyl, L. J. van (1966). Abnormal haemoglobins in Cape Town. Occurrence and significance. *South African Medical Journal*, **40**, 753-756.
- Bowman, B. H., Barnett, D. R., and Hite, R. (1967). Hemoglobin G-Coushatta: a beta variant with a delta-like substitution. *Biochemical and Biophysical Research Communications*, **26**, 466-470.
- Bowman, B. H., Barnett, D. R., Hodgkinson, K. T., and Schneider, R. G. (1966). Chemical characterization of haemoglobin G_{Sst-1}. *Nature*, **211**, 1305-1306.
- Bowman, B. H. and Ingram, V. M. (1961). Abnormal human haemoglobins. VII. The comparison of normal human haemoglobins with haemoglobin D Chicago. *Biochimica et Biophysica Acta*, **53**, 569-573.
- Bowman, B. H., Oliver, C. P., Barnett, D. R., Cunningham, J. E., and Schneider, R. G. (1964). Chemical characterization of three hemoglobins G. *Blood*, **23**, 193-199.
- Cabannes, R. (1965). Répartition des hémoglobines anormales dans la partie ouest du continent Africain. In *Abnormal Haemoglobins in Africa*, ed. by J. H. P. Jonxis, pp. 291-317. Blackwell Scientific, Oxford.
- Cabannes, R., Taleb, N., Ghorra, F., and Schmitt-Beurrier, A. (1965). Études des types hémoglobiques dans la population du Liban. *Nouvelle Revue Française d'Hématologie*, **5**, 851-856.
- Cawein, M. J., Lappatt, E. J., Brangle, R. W., and Farley, C. H. (1966). Hemoglobin S-D disease. *Annals of Internal Medicine*, **64**, 62-70.
- Chatterjee, J. B. (1959). Haemoglobinopathy in India. In *Abnormal Haemoglobins*, ed. by J. H. P. Jonxis and J. F. Delafresnaye, pp. 322-329. Blackwell Scientific, Oxford.
- Chernoff, A. I. (1956). On the prevalence of hemoglobin D in the American negro. *Blood*, **11**, 907-909.
- Chernoff, A. I. (1958). The hemoglobin D syndromes. *Blood*, **13**, 116-127.
- Chernoff, A. I. and Pettit, N. (1965). The amino acid composition of hemoglobin. VI. Separation of the tryptic peptides of hemoglobin Knoxville No. 1 on Dowex-1 X-2 and Sephadex. *Biochimica et Biophysica Acta*, **97**, 47-60.
- Chernoff, A. I. and Weichselbaum, T. E. (1958). A microhemolyzing technic for preparing solutions of hemoglobin for paper electrophoretic analysis. *American Journal of Clinical Pathology*, **30**, 120-125.
- Cholakov, B. and Kanchev, K. N. (1970). A case of hemoglobinosis (In Bulgarian). *Savremenna Medicina*, **21**, 24-26.
- Cook, I. A. and Lehmann, H. (1973). Beta-thalassaemia and some rare haemoglobin variants in the highlands of Scotland. *Scottish Medical Journal*, **18**, 14-20.
- Cortesi, S., Vettore, L., Carta, S., Sorcini, M., and De Sandre, G. (1966). Associazione di emoglobinosi D Los Angeles $\alpha\beta$ -talassaemia in una famiglia Veneta. *Acta Medica Patavina*, **26**, 262-291.
- Dacie, J. V. (1954). *The Haemolytic Anaemias: Congenital and Acquired*, p. 146. Grune and Stratton, New York.
- Dance, N., Huehns, E. R., and Shooter, E. M. (1964). The chemical investigation of haemoglobins G Bristol and G Bristol/C. *Biochimica et Biophysica Acta*, **86**, 144-148.
- David, J. H., Santos and Trincão, C. (1963). Drepanocitemia, deficiência da desidrogenase da glucose-6-fosfato (G-6-PD) eritrocitária e paludismo no posto do Cuango (Lunda-Angola). *Anais do Instituto de Medicina Tropical*, **20**, 5-16.
- Dherte, P., Vandepitte, J., Ager, J. A. M., and Lehmann, H. (1959). Stanleyville I and II. Two new variants of adult haemoglobin. *British Medical Journal*, **2**, 282-284.
- Dreyfus, F. and Pinkhas, J. (1958). Search for abnormal haemoglobins in oriental Jews. *Lancet*, **2**, 1180.
- Eliou, J., Belkhdja, O., Wajcman, H., and Labie, D. (1973). Two variants of hemoglobin D in the Algerian population: Hemoglobin D Ouled Rabah $\beta^{19}\text{B1 Asn} \rightarrow \text{Lys}$ and hemoglobin D Iran $\beta^{22}\text{B4 Glu} \rightarrow \text{Gln}$. *Biochimica et Biophysica Acta*, **310**, 360-369.
- Fessas, P. (1959). The hereditary anaemias in Greece. In *Abnormal Haemoglobins*, ed. by J. H. P. Jonxis and J. F. Delafresnaye, pp. 260-270. Blackwell Scientific, Oxford.
- Flatz, G., Pik, C., and Sringam, S. (1965). Haemoglobinopathies in Thailand. II. Incidence and distribution of elevations of haemoglobin A₂ and haemoglobin F. A survey of 2790 people. *British Journal of Haematology*, **11**, 227-236.
- Gammack, D. B., Huehns, E. R., Lehmann, H., and Shooter, E. M. (1961). The abnormal polypeptide chains in a number of haemoglobin variants. *Acta Genetica et Statistica Medica*, **11**, 1-16.
- Githens, J. H., Knock, H. K., and Hathaway, W. E. (1961). Prevalence of abnormal hemoglobins in American Indian children. Survey in the Rocky Mountain area. *Journal of Laboratory and Clinical Medicine*, **57**, 755-758.
- Gouttas, A., Tseverenis, H., Papaspyrou, A., Pougouras, P., Fertakis, A., and Vorias, N. (1960). L'hémoglobinosse D en Grèce. *Sang*, **31**, 303-306.
- Huisman, T. H. J. (1969). Human hemoglobins. In *Biochemical Methods in Red Cell Genetics*, ed. by J. J. Yunis, pp. 391-504. Academic Press, New York.
- Huntsman, R. G., Hall, M., Lehmann, H., and Sukumaran, P. K. (1963). A second and a third abnormal haemoglobin in Norfolk. Haemoglobin G_{Norfolk} and Haemoglobin D_{Norfolk}. *British Medical Journal*, **1**, 720-722.
- Hynes, M. and Lehmann, H. (1956). Haemoglobin D in a Persian girl; presumably the first case of haemoglobin-D-thalassaemia. *British Medical Journal*, **2**, 923-924.
- Imamura, T. and Riggs, A. (1972). Identification of hemoglobin Oak Ridge with hemoglobin D Punjab (Los Angeles). *Biochemical Genetics*, **7**, 127-130.
- Itano, H. A. (1951). Third abnormal hemoglobin associated with hereditary hemolytic anemia. *Proceedings of the National Academy of Sciences*, **37**, 775-784.
- Itano, H. A. (1953). Solubilities of naturally occurring mixtures of human hemoglobin. *Archives of Biochemistry and Biophysics*, **47**, 148-159.
- Jacob, G. F., Lehmann, H., and Raper, A. B. (1956). Haemoglobin D in Indians of Gujerati origin in Uganda. *East African Medical Journal*, **33**, 135-138.
- Jong, W. W. W. de and Went, L. N. (1968). Haemoglobin J_{Baltimore} ($\alpha\beta_2$ $^{16}\text{Gly} \rightarrow \text{Asp}$) and haemoglobin D Punjab ($\alpha\beta_2$ $^{121}\text{Glu} \rightarrow \text{Gln}$) in two Dutch families. *Acta Genetica et Statistica Medica*, **18**, 429-443.
- Kohne, E., König, E., and Roggenbach, H. J. (1972). Hämoglobin D-Punjab bei einem deutschen Patienten. *Verhandlungen der Deutschen Gesellschaft für Innere Medizin*, **78**, 1596-1597.
- Konigsberg, W., Huntsman, R. G., Wadia, F., and Lehmann, H. (1965). Haemoglobin D β Punjab in an East Anglian family. *Journal of the Royal Anthropological Institute*, **95**, 295-306.
- Konotey-Ahulu, F. I. D., Gallo, E., Lehmann, H., and Ringelmann, B. (1968). Haemoglobin Korle-Bu ($\beta^{73}\text{Aspartic acid} \rightarrow \text{asparagine}$). *Journal of Medical Genetics*, **5**, 107-111.
- Labie, D. and Rosa, J. (1969). Structure de quelques hémoglobines anormales. *Annales de la Société Belge de Médecine Tropicale*, **49**, 99-107.

- Lehmann, H., Ala, F., Hedeyat, S., Montazemi, K., Nejad, H. K., Lightman, S., Kopeck, A. C., Mourant, A. E., Teesdale, P., and Tills, D. (1973). The hereditary blood factors of the Kurds of Iran. *Philosophical Transactions of the Royal Society of London, Series B*, **226**, 195–205.
- Lehmann, H., Beale, D., and Boi-Doku, F. S. (1964). Haemoglobin G_{Accra}. *Nature*, **203**, 363–365.
- Lehmann, H. and Nwokolo, C. (1959). The river Niger as a barrier in the spread eastwards of haemoglobin C: a survey of haemoglobins in the Ibo. *Nature*, **183**, 1587–1588.
- Licenziati, M. (1969). Incidenza delle emoglobinopatie nella Provincia di Napoli. *Minerva Medica*, **60**, 140–141.
- Liddell, J., Brown, D., Beale, D., Lehmann, H., and Huntsman, R. G. (1964). A new haemoglobin-Ja Oxford found during a survey of an English population. *Nature*, **204**, 269–270.
- Lie-Injo, L. E. (1959). Pathological haemoglobins in Indonesia. In *Abnormal Haemoglobins*, ed. by J. H. P. Jonxis and J. F. Delafresnaye, pp. 368–383. Blackwell Scientific, Oxford.
- Lisker, R. (1971). Distribution of abnormal hemoglobins in Mexico. In *Genetical, Functional and Physical Studies of Hemoglobins*, ed. by T. Arends, G. Bemski, and R. L. Nagel, pp. 8–14. Karger, Basel.
- Lisker, R., Zarate, G., and Loria, A. (1966). Studies on several genetic hematologic traits of Mexicans. IX. Abnormal hemoglobins and erythrocytic glucose-6-phosphate dehydrogenase deficiency in several Indian tribes. *Blood*, **27**, 824–830.
- McCormick, W. F. (1960). Abnormal hemoglobins. I. Incidence in Memphis and Western Tennessee, with special reference to autopsy material. *American Journal of Clinical Pathology*, **34**, 220–224.
- McCurdy, P. R. (1959). Clinical and physiologic studies in a negro with sickle cell hemoglobin D disease. *New England Journal of Medicine*, **262**, 961–964.
- McCurdy, P. R., Pearson, H., and Gerald, P. S. (1961). A new hemoglobinopathy of unusual genetic significance. *Journal of Laboratory and Clinical Medicine*, **58**, 86–94.
- Marder, V. J. and Conley, C. L. (1959). Electrophoresis of hemoglobin on agar gels: frequency of hemoglobin D in a negro population. *Bulletin of the Johns Hopkins Hospital*, **105**, 77–88.
- Marengo-Rowe, A. J., McCracken, A. W., and Flanagan, P. (1968). Complete suppression of haemoglobin A synthesis in haemoglobin D Los Angeles—beta thalassaemia. *Journal of Clinical Pathology*, **21**, 508–510.
- Martin, H., Heupke, G., Pfeiderer, G., and Wörner, W. (1960). Hämoglobin D in einer Frankfurter Familie. *Folia Haematologica*, **4**, 233–241.
- Milner, P. F. (1967). High incidence of haemoglobin G_{Accra} in a rural district of Jamaica. *Journal of Medical Genetics*, **4**, 88–90.
- Minnich, V., Cordonnier, J. K., Williams, W. J., and Moore, C. V. (1962). Alpha, beta, and gamma hemoglobin polypeptide chains during the neonatal period with description of a fetal form of D_α St Louis. *Blood*, **19**, 137–167.
- Monn, E., Gaffney, P. J., and Lehmann, H. (1968). Haemoglobin Sogn (β14 arginine). A new haemoglobin variant. *Scandinavian Journal of Haematology*, **5**, 353–360.
- Myerson, R. M., Harrison, E., and Lohmuller, A. W. (1959). Incidence and significance of abnormal hemoglobins. *American Journal of Medicine*, **26**, 543–546.
- Nilsson, L. O. and Eriksson, A. W. (1972). Screening for haemoglobin and lactate dehydrogenase variants in the Icelandic, Swedish, Finnish, Lappish, Mari, and Greenland Eskimo populations. *Human Heredity*, **22**, 372–379.
- Özsoylu, S. (1970). Homozygous hemoglobin D Punjab. *Acta Haematologica*, **43**, 353–359.
- Pinango, C. L. A. de and Arends, T. (1965). Hemoglobinas anormales en donantes de sangre nativos del Estado Bolívar. *Acta Científica Venezolana*, **16**, 215–218.
- Pollitzer, W. S., Chernoff, A. I., Horton, L. L., and Froehlich, M. (1959). Hemoglobin patterns in American Indians. *Science*, **129**, 216.
- Preto, R. V., Trincão, C., Melo, J., Cordeiro Ferreria, N., and Coutinho, P. (1961). Considerações sobre numa hemoglobinopatia familiar associada a talassemia-hemoglobina D numa família algarvia. *Boletim Clínico dos Hospitais Cívicos de Lisboa*, **25**, 483–486.
- Quattrin, N., Bianchi, P., Cimino, P., De Rosa, L., Dini, E., Montuori, R., and Ventruito, V. (1966). Le microcitemie ed altre emoglobinopatie in Campania. Cinque anni di recherche. *Riforma Medica*, **80**, 285–295.
- Rahbar, S. (1973). Haemoglobin D Iran: β₂₂ Glutamic acid → Glutamine (B4). *British Journal of Haematology*, **24**, 31–35.
- Rahbar, S., Beale, D., Isaacs, W. A., and Lehmann, H. (1967). Abnormal haemoglobins in Iran. Observation of a new variant—haemoglobin J Iran (α₂β₂⁷⁷ His → Asp). *British Medical Journal*, **1**, 674–677.
- Ramot, B., Rotem, J., Rahbar, S., Jacobs, A. S., Udem, L., and Ranney, H. M. (1969). Hemoglobin D Punjab in a Bulgarian Jewish family. *Israel Journal of Medical Sciences*, **5**, 1066–1070.
- Raper, A. B., Gammack, D. B., Huehns, E. R., and Shooter, E. M. (1960). Four haemoglobins in one individual. A study of the genetic interaction of Hb-G and Hb-C. *British Medical Journal*, **2**, 1257–1261.
- Restrepo, A. (1971). Frequency and distribution of abnormal hemoglobins and thalassemia in Colombia, South America. In *Genetical, Functional and Physical Studies of Hemoglobins*, ed. by T. Arends, G. Bemski, and R. L. Nagel, pp. 39–52. Karger, Basel.
- Restrepo, H. A. and Londoño, O. G. (1965). Sickle cell hemoglobin D disease in a negro Colombian patient. *Annals of Internal Medicine*, **62**, 1301–1306.
- Ricco, G., Gallo, E., Mazza, U., Papa, G., and Inglott, G. (1970). Su un caso di emoglobina D Punjab (α₂β₂¹²¹ Glu → Gln). *Bollettino della Società Italiana di Biologia Sperimentale*, **46**, 341–344.
- Ringelmann, B., Lewis, R. A., Lorkin, P. A., Kynoch, P. A. M., and Lehmann, H. (1967). Sickle-cell haemoglobin D Punjab disease: S from Ghana and D from England. *Acta Haematologica*, **38**, 324–331.
- Robinson, A. R., Robson, M., Harrison, A. P., and Zuelzer, W. W. (1957). A new technique for the differentiation of hemoglobin. *Journal of Laboratory and Clinical Medicine*, **50**, 745–752.
- Rohe, R. A., Sharma, V., and Ranney, H. M. (1973). Hemoglobin D Iran α₂β₂²² Glu → Gln in association with thalassemia. *Blood*, **42**, 455–462.
- Rossi, U. and Morelli, A. (1970). Emoglobine rare: frequenza presso la popolazione della provincia di Milano. *Folia Hereditaria et Pathologica*, **19**, 29–33.
- Rothman, M. C. and Ranney, H. M. (1971). Double heterozygosity for hemoglobin G (α₆₈ Lys β₂^A) and hemoglobin D (α₂β₂¹²¹ Gln). *Blood*, **37**, 177–183.
- Sadikario, A., Duma, H., Efremov, G., Mladenovski, B., Andreeva, M., Petkov, G., and Lazova, C. (1969). Thalassemias and abnormal hemoglobins in SR Macedonia. *Acta Haematologica*, **41**, 162–169.
- Salzano, P. M. and Tondo, C. V. (1968). Hemoglobin types of Brazilian Indians. *American Journal of Physical Anthropology*, **28**, 355–359.
- Schneider, R. G., Haggard, M. E., McNutt, C. W., Johnson, J. E., Bowman, B. H., and Barnett, D. R. (1964). Hemoglobin G_{Coushatta}: a new variant in an American Indian family. *Science*, **143**, 697–698.
- Schneider, R. G., Ueda, S., Alperin, J. B., Levin, W. C., Jones, R. T., and Brimhall, B. (1968). Hemoglobin D Los Angeles in two Caucasian families: hemoglobin SD disease and hemoglobin D thalassemia. *Blood*, **32**, 250–259.
- Shibata, S., Iuchi, I., and Miyaji, T. (1966). Abnormal hemoglobins discovered in Japan. *Bulletin of the Yamaguchi Medical School*, **13**, 1–17.
- Sick, K., Beale, D., Irvine, D., Lehmann, H., Goodall, P. T., and MacDougall, S. (1967). Haemoglobin G_{Copenhagen} and haemoglobin J_{Cambridge}. Two new β-chain variants of haemoglobin A. *Biochimica Biophysica Acta*, **140**, 231–242.
- Singer, K., Chernoff, A. I., and Singer, L. (1951). Studies on abnormal hemoglobins. I. Their demonstration in sickle cell anemia and other hematologic disorders by means of alkali denaturation. *Blood*, **6**, 413–429.
- Smith, E. W. and Conley, C. L. (1959). Sickle cell—hemoglobin D disease. *Annals of Internal Medicine*, **50**, 94–105.
- Stern, M. A., Kynoch, P. A. M., and Lehmann, H. (1968). β-Thalassemia, glucose-6-phosphate dehydrogenase deficiency and haemoglobin D-Punjab in Pathans. *Lancet*, **1**, 1284–1285.
- Stewart, J. N. and MacIver, J. G. (1956). Sickle cell—haemoglobin-D disease in a mulatto girl. *Lancet*, **1**, 23–25.
- Stout, C., Holland, C. K., and Bird, R. M. (1964). Hemoglobin D in an Oklahoma family. *Archives of Internal Medicine*, **114**, 296–300.

- Stuart, J. C., Schwartz, F. C. M., Little, A. J., and Raine, D. N. (1973). Screening for abnormal haemoglobins: a pilot study. *British Medical Journal*, **4**, 284-287.
- Sturgeon, P., Itano, H. A., and Bergren, W. R. (1955). Clinical manifestations of inherited abnormal hemoglobins. I. The interaction of hemoglobin-S with hemoglobin-D. *Blood*, **10**, 389-404.
- Traverse, P. M. de and Coquelet, M. L. (1961). Hémoglobine foetale anormale chez un enfant hétérozygote AD. Hypothèse structurale. *Comptes Rendus des Séances de la Société de Biologie et de ses Filiales*, **155**, 1888-1892.
- Trinçao, C. (1967). Hemoglobinas anormais nos territorios Portuguesas. *Boletim Clínico dos Hospitais Cívicos Lisboa*, **21**, 813-827.
- Trinçao, C. and Ferreira, C. N. (1962). Thalassaemia in Portugal. In *Proceedings of the 8th Congress of the European Society of Haematology*, pt. 2, 307a. Karger, New York.
- Ungar, B., Symons, H. S., Anderson, G. R., Stubbs, A. E., Wiltshire, B. G., and Lehmann, H. (1973). Haemoglobin D Punjab in a European family in Australia. *Medical Journal of Australia*, **1**, 354-356.
- Vella, F. (1961/1962). Abnormal haemoglobins, thalassaemia and erythrocyte glucose-6-phosphate dehydrogenase deficiency in Singapore and Malaya. *Oceania*, **32**, 219-225.
- Vella, F. (1967). Haemoglobin variants in Saskatchewan. *Clinical Biochemistry*, **1**, 118-134.
- Vella, F. and Guzak, P. (1968). Haemoglobin variants and thalassaemia in Saskatchewan Indians. *Clinical Biochemistry*, **2**, 153-157.
- Vella, F., Isaacs, W. A., and Lehmann, H. (1967a). Hemoglobin G Saskatoon: $\beta^{22} \text{Glu} \rightarrow \text{Ala}$. *Canadian Journal of Biochemistry*, **45**, 351-353.
- Vella, F., Lorkin, P. A., Carrell, R. W., and Lehmann, H. (1967b). A new hemoglobin variant resembling hemoglobin E. Hemoglobin E Saskatoon: $\beta^{22} \text{Glu} \rightarrow \text{Lys}$. *Canadian Journal of Biochemistry*, **45**, 1385-1391.
- Ventruto, V., Colombo, B., Caniglia, A., and De Rosa, L. (1963). Studio di tre famiglie con emoglobinosi D. *Haematologica*, **48**, 873-884.
- Wade, P. T., Jenkins, T., and Huehns, E. R. (1967). Haemoglobin variant in a Bushman: Haemoglobin D β -Bushman $\alpha\beta_{22}^{16} \text{Gly} \rightarrow \text{Arg}$. *Nature*, **216**, 688-690.
- Wajcman, H., Boigne, J. M., Labie, D., and Serigne, P. (1969). Hémoglobine D Los Angeles. Mise en évidence de l'anomalie de structure par une amélioration des méthodes. *Biochimica et Biophysica Acta*, **188**, 55-58.
- Wasi, P., Pootrakul, S., Na-Nakorn, S., Beale, D., and Lehmann, H. (1968). Haemoglobin D Los Angeles (D Punjab, $\alpha\beta_{22}^{121} \text{Glu NH}_2$) in a Thai family. *Acta Haematologica*, **39**, 151-158.
- Watson-Williams, E. J., Beale, D., Irvine, D., and Lehmann, H. (1965). A new haemoglobin D Ibadan (β -87 threonine $\rightarrow$ lysine) producing no sickle-cell haemoglobin D disease with haemoglobin S. *Nature*, **205**, 1273-1276.
- Weatherall, D. J., Sigler, A. T., and Baglioni, C. (1962). Four hemoglobins in each of three brothers. Genetic and biochemical significance. *Bulletin of the Johns Hopkins Hospital*, **111**, 143-156.
- White, J. C. and Beaven, G. H. (1954). A review of the varieties of human haemoglobin in health and disease. *Journal of Clinical Pathology*, **7**, 175-200.

Oxford Symposium

A symposium on 'The eye in the inborn errors of metabolism' will take place at Oxford University, Oxford, England from 13 to 16 April 1975. Speakers will include: Norman Ashton, Elmer Ballintine, Elaine Berman, Anthony Bron, Donald Bergsma, Elliot Berson, Robert P. Burns, Ronald E. Carr, David G. Cogan, Patrick I. Condon, Edward Cotlier, Harold Cross, Glyn Dawson, Monte A. Del Monte, Albert T. Franceschetti, Jules Francois, Alec Garner, Morton Goldberg, Brian Harcourt, William F. Hughes, Barrie Jay, Kenneth Kenyon, Gordon Klintworth, Toichiro Kuwabara, Irene Hussels-Maumenee, Michael D. Sanders, Jack D. Singer, George Spaeth, C. Takki, Ramesh Tripathi, Ruth van Heyningen, J. M. Walshe, Warren Wilson, Jonathan Wirschafter, Vernon Wong, Wolfgang Zeman. A limited number of free papers will be accepted. Ophthalmologists, paediatricians, or basic scientists are invited to attend; register in advance. For information regarding registration or participation, contact the symposium organizers; Edward Cotlier, MD, University of Illinois Eye and Ear Infirmary, 1855 W. Taylor Street, Chicago, Illinois 60612, USA, or Anthony Bron, FRCS, University of Oxford Nuffield Laboratory of Ophthalmology, Walton Street, Oxford OX2 6AN, England.