Haemoglobin D Ouled Rabah (β 19[B1] Asn→Lys) in a Tuareg tribe of the Southern Sahara

A. MAURAN-SENDRAIL, Ph. LEFEVRE-WITIER, H. LEHMANN, AND R. CASEY

From Centre d'Hématologie du C.N.R.S., Hôpital Purpan, 31300—Toulouse, France; and Department of Clinical Biochemistry, Addenbrooke's Hospital, Hills Road, Cambridge, England

SUMMARY Haemoglobin Ouled Rabah found only once before in an Algerian has been discovered at high frequency in the Kel Kummer Tuaregs in Mali. The tribe represents a 'genetic isolate'.

In November to December 1971, an anthropological and genetic study was carried out of the Kel Kummer Tuaregs. These Tuaregs, numbering 367 individuals, belong to a local leader clan nomadising around Menaka (Mali). The clan is highly endogamous, 85% of the genes in the community being derived from 25 'founders' who can be traced to the beginning of the genealogies spanning 15 generations. Thus, the Kel Kummer Tuaregs constitute a 'genetic isolate' built up some 300 years ago (Chaventre, 1971, 1972; Degos, 1973).

This report concerns the finding of an unusual variant of human haemoglobin (Hb) at a high frequency in this community. This variant, Hb Ouled Rabah (β 19[B1] Asn→Lys) has been reported once before in an Algerian from the Aures mountains (Elion *et al.*, 1973) and may prove to be a useful anthropological marker.

Methods

(a) THE SURVEY

The sample studied represented about 76% of the Tuareg population.

Genealogical tables showing the family relations between carriers of Hb Ouled Rabah were drawn by A. Chaventre (Fig. 1).

The following methods were used routinely: (i) Electrophoresis of haemolysate using paper, cellulose acetate, starch gel (pH 9.2), and agar gel (pH 6.2) (Lehmann and Huntsman, 1974); (ii) the Itano solubility test (Itano, 1953); (iii) Hb F estimation by

alkali denaturation (Singer *et al.*, 1951); and (iv) Hb A₂ estimation (Marengo-Rowe, 1965).

(b) HAEMOGLOBIN STRUCTURAL STUDIES

Studies were carried out on haemolysate prepared from an individual who had been shown, from the survey, to be heterozygous for the haemoglobin variant.

DEAE-Sephadex chromatography of the haemolysate was carried out according to Huisman and Dozy (1965); the variant haemoglobin was further purified, after DEAE-Sephadex chromatography, by paper electrophoresis.

The purified Hb D was converted to globin (Winterhalter and Huehns, 1964) and the latter was separated into α - and β -chains by CM-cellulose chromatography in the presence of 8M urea (Clegg *et al.*, 1966). The α - and β -chains were aminoethylated (Raftery and Cole, 1966), freed of urea and salts by dialysis against 0.5% (v/v) formic acid in water at 4°C, and recovered by freeze-drying. The isolated aminoethylated chains were digested with trypsin (1/50 (w/w), 37°, 2h, in 0.08 NH₄HCO₃); insoluble material was removed by centrifugation and the supernatant was freeze-dried and 'fingerprinted' in the usual manner (Beale, 1967).

Appropriate peptides (see results) were eluted from a preparative-scale 'fingerprint' which had been lightly stained with 0.02% (w/v) ninhydrin in acetone (containing 1% (v/v) pyridine) with constant-boiling HCl and hydrolysed at 108°C in a sealed tube for 16 hours. After removal of excess HCl *in vacuo* over NaOH pellets, the amino acid composition of the dried hydrolysate was determined (Spackman *et al.*, 1958) using a Locarte amino acid analyser.

Fig. 1 Genealogical tree of the Kel Kummer tribe studied.

Results

(a) THE SURVEY

In 285 samples studied the abnormal haemoglobin was found together with Hb A in 57; a single Hb D band was found in 3 cases, presumably representing the homozygous state. None of the specimens showed a raised Hb A₂, but 3 (1%) showed a raised level of Hb F.

(b) THE HEMOGLOBIN VARIANT

Examination of the haemolysate by paper electrophoresis showed the presence of Hb A, a single Hb A₂, and a slow-moving (D) haemoglobin. The relative proportions of the three fractions from DEAE-Sephadex were Hb A₂, 2.5%, Hb A, 50.5%, and Hb D, 47.0%. The presence of only one Hb A₂ suggested that the amino acid substitution was in the β -, rather than the α -, chain.

CM-cellulose chromatography in 8M urea of the globin from the purified Hb D showed the α -chains to elute in the normal position; the β -chains, however, eluted later than normal, suggesting a loss of one negative charge, or a gain of one positive charge, per β^D -chain.

The 'fingerprint' of the aminoethylated (AE)

α -chains was normal. Comparison of that of the AE β^D -chains with the pattern obtained from AE β^A -chains, however, showed the presence of a new, negatively-charged peptide which gave a positive staining reaction for arginine (β^D 18-30 in Fig. 2). The amino acid composition of this peptide resembled that of β^A Tp III, except that one of the two residues of aspartic acid normally found on analysis of β^A Tp III had been replaced by a residue of lysine (Table 1). The electrophoretic mobilities of the variant peptide and the whole haemoglobin made it clear that there had been an increase of one positive charge per

Table 1 Amino acid composition of peptide β TPIII from Hb A and peptides β 18-30 and β 20-30 from Hb Ouled Rabah

Amino acid	β^D 18-30	β^A TpIII	β^D 20-30
Asp	1.0	2	1.0
Glu	2.1	2	2.0
Gly	2.7	3	2.8
Ala	1.0	1	1.0
Val	2.9	3	2.0
Leu	1.0	1	1.0
Lys	1.2	—	—
Arg	1.0	1	1.0

One amino acid residue is approximately 48 nmol for β^D Tp 18-30 and 16 nmol for β^D 20-30.

Table 2 Amino acid sequence of the tryptic peptide $\beta 18$ –30 from Hb D Ouled Rabah

Residue No. Helical No.	18 A14	19 B1	20 B2	21 B3	22 B4	23 B5	24 B6	25 B7	26 B8	27 B9	28 B10	29 B11	30 B12
Residue:													
Hb ^A	↓Val	Asn	Val	Asp	Glu	Val	Gly	Gly	Glu	Ala	Leu	Gly	Arg↓
Hb ^D Ouled Rabah	↓Val	Lys↓	Val	Asp	Glu	Val	Gly	Gly	Glu	Ala	Leu	Gly	Arg↓

↓ Indicates the positions of hydrolysis by trypsin.

β -chain, indicating that asparagine $\beta 19$, rather than aspartic acid $\beta 21$ (see Table 2) had been substituted (asparagine is converted to aspartic acid during the acid hydrolysis which precedes amino acid analysis).

There was apparently no peptide absent from the 'fingerprint' of the AE β^D chain; it is, however, to be expected that some tryptic hydrolysis would normally take place between lysine $\beta^D 19$ and valine $\beta 20$, and the products, $\beta^D 18$ –19 (Val-Lys) and $\beta 20$ –30 (see Table 2), would be expected to appear on the 'fingerprint' at positions overlapping β^A Tp VI ($\beta 60$ –61, Val-Lys) and β^A Tp III ($\beta 18$ –30), respectively. The spot which apparently corresponded to β^A Tp III was, therefore, eluted, hydrolysed, and analysed; its amino acid composition (Table 1) was that expected for $\beta 20$ –30, showing that limited hydrolysis had occurred between lysine $\beta 19$ and valine $\beta 20$ (Table 2). From the relative yields of the two

peptides $\beta^D 18$ –30 and $\beta^D 20$ –30, it could be calculated that the extent of hydrolysis at lysine $\beta 19$ was about 25% of that at arginine $\beta 30$. This slower rate of hydrolysis probably results from the proximity of aspartic acid $\beta 21$ and glutamic acid $\beta 22$ (Table 2), as pointed out by Elion *et al.* (1973).

These results show that this variant is Hb Ouled Rabah ($\beta 19$ (B1) Asn $\rightarrow$ Lys) (Elion *et al.*, 1973).

Discussion

In the Tuareg of Arabo-Berber tribes so far investigated, the only unusual haemoglobins to have been identified (electrophoretically) have been Hb S, Hb C, and Hb K. The frequency of these variants (2 to 3%) is low compared with that observed in the sedentary people of the oases (10 to 20%) (Cabannes, 1964; Cabannes and Ruffie, 1961; Cabannes *et al.*,

Table 3 Abnormal haemoglobins in the Tuareg tribe

Total	AD	DD	%	$q^2\beta^D$	Thalassaemia				
					$A_2 \uparrow$	$F \uparrow$	$(A_2 + F) \uparrow$	Total	%
285	57	3	21	0.1105	—	3	—	3	1

Fig. 2 'Fingerprint' of the tryptic peptides of the AE β chain from Hb D Ouled Rabah. Electrophoresis at pH 6.4, 53 V/cm for 1 hour. Ascending chromatography in pyridine-isoamyl alcohol-water (6:6:7, by vol.) for 20 hours. Peptides located by staining with 0.2% ninhydrin in acetone containing 1% v/v pyridine.

*=origin.

1967, 1969a, b; Ruffie *et al.*, 1959a, b, 1962, 1963; Treilhou, 1969). Table 3 shows the haemoglobin distribution observed in the present survey (see also Mauran-Sendrail and Lefevre-Witier, 1973), from which it can be seen that the frequency of Hb D in the Tuareg tribe studied is very high. Though the Hb D was identified as Hb Ouled Rabah in only a limited number of cases, it is likely that the variant affects the bulk of the tribe, which is a genetic isolate with extremely intricate family ties, as shown in Fig. 2 and previously discussed elsewhere (Chaventre 1971, 1972; Landre and Valat, 1972; Langaney *et al.*, 1973). While it is possible that the variant might have a certain selective value for example against *Plasmodium falciparum* infection (from which the Kel Kummer Tuaregs suffer every year), thus leading to its observed high frequency, the fact that the population is highly inbred could be the sole reason for the high incidence of Hb Ouled Rabah.

The study of different blood markers has revealed a clearly Mediterranean genetic structure in the Kel Kummer Tuaregs, close to that of Moroccan Chleuh or Mauritanian Reguibat (Degos, 1973; Constans and Lefevre-Witier, 1973). The finding of Hb D Ouled Rabah would fall within that genetic pattern; but while it is noteworthy that the male founder of the tribe came from Northern Sahara, the probability of transmission of Hb D Ouled Rabah by that founder (5.7%) should be compared with the probable contribution of the Tuareg group already settled in that area (24.2%) (Jacquard, 1972).

In conclusion, it is felt that Hb Ouled Rabah may prove to be a useful anthropological marker in the study of the tribes of the Sahara.

References

- Beale, D. (1967). A partial amino acid sequence for sheep haemoglobin A. *Biochemical Journal*, **103**, 129-140.
- Cabannes, R. (1964). Les types hémoglobiniques des populations de la partie occidentale du continent africain. *Monographie du Centre d'Hématologie due C.N.R.S.*, p. 148.
- Cabannes, R., Larrouy, G., Fernet, P., and Sendrail, A. (1969a). Étude hémotypologique des populations sédentaires de la Saoura (Sahara Occidentale) II. Les hémoglobines. *Bulletin et Mémoires de la Société d'Anthropologie de Paris*, **4**, 139-142.
- Cabannes, R., Larrouy, G., and Sendrail, A. (1969b). Étude hémotypologique des populations du Massif du Hoggar et du plateau de l'Air III. Les hémoglobines. *Bulletin et Mémoires de la Société d'Anthropologie de Paris*, **4**, 143-146.
- Cabannes, R., Lefevre-Witier, Ph., and Sendrail, A. (1967). Étude des hémoglobines dans les populations du Tassili N'ajjer (Sahara Central). *Bulletin et Mémoires de la Société d'Anthropologie de Paris*, **1**, 434-439.
- Cabannes, R., and Ruffie, J. (1961). Les types hémoglobiniques dans les populations des massifs montagneux du Sahara Central (Hoggar et Tibesti). *L'Anthropologie*, **65**, 467-483.
- Chaventre, A. (1971). Présentation d'un isolat targui. *Population*, **1-2**, 188-189.
- Chaventre, A. (1972). Les Kel Kummer: isolat social. *Population*, **4-5**, 771-783.
- Clegg, J. B., Naughton, M. A., and Weatherall, D. J. (1966). Abnormal human haemoglobins. Separation and characterisation of the α and β chains by chromarography, and the determination of the two new variants, Hb Chesapeake and Hb J (Bangkok). *Journal of Molecular Biology*, **19**, 91-108.
- Constans, J., and Lefevre-Witier, Ph. (1973). Nouvelles données hémotypologiques en pays touareg. I. Les types d'haptoglobine du village d'Idélès (Ahaggar, Sahara Algérien) et chez les Kel Kummer nomades (touareg Iwellemmeden, Kel Attaram, Mali). In *IXth International Congress of Anthropological and Ethnological Sciences, Chicago, September 1973*.
- Degos, L. (1973). Structure génétique du système Hl-A chez les Kel Kummer. *Population*, **6**, 1111-1114.
- Elion, J., Belkhdja, O., Wajcman, H., and Labie, D. (1973). Two variants of Hb D in the Algerian population: Hb D Ouled Rabah β^{19} (B₁) Asn→Lys and Hb D Iran β^{22} (B₄) Glu→Gln. *Biochimica et Biophysica Acta*, **310**, 360-364.
- Huisman, T. H. J., and Dozy, A. M. (1965). Studies on the heterogeneity of hemoglobin. IX. The use of tris (hydroxymethyl)amino-methane HCl buffers in the anion-exchange chromatography of hemoglobins. *Journal of Chromatography*, **19**, 160-169.
- Itano, H. A. (1953). Solubilities of naturally occurring mixtures of human hemoglobin. *Archives of Biochemistry*, **47**, 148-159.
- Jacquard, A. (1972). Evolution du patrimoine génétique des Kel Kummer. *Population*, **4-5**, 784-800.
- Landre, M. T., and Valat, M. T. (1972). Un isolate du Sud Sahara: les Kel Kummer. Tracé automatique des généalogies. *Population*, **4-5**, 801-803.
- Langaney, A., Chaventre, A., Lefevre-Witier, Ph., and Jacquard, A. (1973). Filiation présumée, filiation biologique et détermination des phénotypes hémotypologiques chez les Kel Kummer. *Population*, **6**, 1115-1124.
- Lehmann, H., and Huntsman, R. G. (1974). *Man's Haemoglobins*, 2nd ed. North-Holland, Amsterdam.
- Marengo-Rowe, A. J. (1965). Rapid electrophoresis and quantitation of haemoglobin on cellulose acetate. *Journal of Clinical Pathology*, **18**, 790-792.
- Mauran-Sendrail, A., and Lefevre-Witier, Ph. (1973). Nouvelles données hémotypologiques en pays touareg. II. Étude des hémoglobines dans le village d'Idélès (Ahaggar, Sahara Algérien) et dans une fraction Kel Kummer (Touareg Iwellemmeden Kel Attaram, Mali). In *IXth International Congress of Anthropological and Ethnological Sciences, Chicago, September 1973*.
- Raftery, M. A., and Cole, R. D. (1966). On the aminoethylation of proteins. *Journal of Biological Chemistry*, **241**, 3457-3461.
- Ruffie, J., Ducos, J., and Larrouy, G. (1962). Étude hémotypologique des populations Chaambas de la région du M'zab (département des oasis). *Bulletin et Mémoires de la Société d'Anthropologie de Paris*, **3**, XIe s., 354-371.
- Ruffie, J., Ducos, J., Ranque, J., and Rioux, J. (1959a). Étude séro-anthropologique des populations du Tibesti (Sahara Oriental). In *Proceedings of the 8th Congress of the International Society of Blood Transfusion, Tokyo*, pp. 195-202.
- Ruffie, J., Ducos, J., and Sicart, M. (1959b). Étude séro-anthropologique des populations Réguibat (Sahara Occidental). In *Proceedings of the 8th Congress of the International Society of Blood Transfusion, Tokyo*, pp. 187-194.
- Ruffie, J., Ducos, J., and Vergnes, H. (1963). Étude hémotypologique des populations du Tidikelt (Sahara Central). *Bulletin et Mémoires de la Société d'Anthropologie de Paris*, **3**, XIe s., 531-544.

Singer, K., Chernoff, A. I., and Singer, L. (1951). Studies on abnormal hemoglobins. II. Their identification by means of the method of fractional denaturation. *Blood*, **6**, 429-439.

Spackman, D. H., Stein, W. H., and Moore, S. (1958). Automatic recording apparatus for use in the chromatography of amino acids. *Analytical Chemistry*, **30**, 1190-1206.

Treilhou, J. P. (1969). Nomades et sédentaires d'Agadès. Étude hémotypologique. Thesis, University of Toulouse.

Winterhalter, K. H., and Huehns, E. R. (1964). Preparation, properties and specific recombination of α - β globin subunits. *Journal of Biological Chemistry*, **239**, 3699-3705.

Requests for reprints to Professor H. Lehmann, Department of Clinical Biochemistry, Addenbrooke's Hospital, Hills Road, Cambridge CB2 2QR.