Taurodontism and Klinefelter's syndrome

YOSUKE KOMATZ, TADAO TOMOYOSHI, OSAMU YOSHIDA, ATHUSHI FUJIMOTO, AND KAZUSADA YOSHITAKE

From the Departments of Urology and Oral Surgery, Faculty of Medicine, Kyoto University, Kyoto, Japan

SUMMARY The incidence of taurodontism in 31 patients with XXY Klinefelter's syndrome was studied. Taurodont molars were observed in 6 of the 31 cases (19.4%), a significantly higher rate than among the controls. Though taurodontism is not an obligatory finding in Klinefelter's syndrome it is believed to be one of the anomalies frequently observed in connection with this condition.

Taurodontism is an anomaly of the shape of the teeth. It is found in multirrooted teeth and is characterised radiologically by short roots, elongated body, and enlarged pulp chamber. The word 'taurodontism' was first used by Keith (1913) to describe the teeth of prehistoric man, the Neanderthals, and the Heidelberg. Anthropologically, it has been viewed with interest in connection with the evolution of man (Shaw, 1928; Hamner *et al.*, 1964; Barker, 1976).

It is believed that taurodontism was first discussed in relation to Klinefelter's syndrome when Law *et al.* (1969) reported that the 2 conditions often appear concurrently. Since then, cases of Klinefelter's syndrome accompanied by taurodontism have been reported by several authors. These cases have included not only the standard XXY type, but also the XXY/XXXY, XXXY, XXXXY, and XXXXY/XXXY variants (Witkop, 1971; Keeler, 1973; Stewart, 1974; Feichtinger and Rossiwall, 1977).

Taurodontism, however, is not an obligatory finding in Klinefelter's syndrome and its incidence is, as yet, unclear. In this report, 31 patients with XXY Klinefelter's syndrome were studied to ascertain the incidence of taurodontism in this condition.

Materials and methods

Thirty-one patients with Klinefelter's syndrome, treated at the Infertility Clinic of the Department of Urology, Kyoto University Hospital, were studied. Their ages ranged from 21 to 42. Chromosome studies were performed on cells from peripheral blood cultures, according to the method of Moorhead *et al.* (1960). In each case, 30 cells were examined and a minimum of 16 cells with the modal chromosome number were karyotyped. The karyotype of all the cases was 47,XXY.

Received for publication 13 February 1978

The controls consisted of 48 males and 63 females without phenotypic anomalies, chosen from patients treated at the Oral Surgery Clinic of this hospital. The ages of the controls, both males and females, ranged from 25 to 42.

Panorex radiographs were used to measure the length of the 'body', as defined by Shaw (1928), as well as that of the crown of the permanent mandibular molars. A molar with a crown/body ratio of more than 0.5 was designated as a taurodont tooth. Maxillary molars were not used for this measurement because of the difficulty of obtaining clear radiographs. In all cases, the measurements were performed by one examiner (AF) who did not know whether the patient belonged to the Klinefelter's syndrome group or to the control group.

Results

Taurodontism was observed in 6 of the 31 Klinefelter's syndrome patients (19.4%), while it was seen in 2.1% of the control males and 4.8% of the control females, indicating a significantly higher ratio in the Klinefelter's syndrome group (Table 1). Of the Klinefelter's syndrome patients with taurodontism, 5 were affected bilaterally (Fig.) and 4 unilaterally. Of the 9 taurodont teeth found in the 6 cases, 7 were

Table 1 Incidence of taurodontism in Klinefelter's syndrome and in controls

	No. of affected cases (%)
Klinefelter's syndrome n = 31	6 (19.4)
Control males n = 48	1 (2.1)
Control females n = 63	3 (4.8)

Fig. 1 Taurodontism affecting the mandibular left second and third, and right second molars (Case 1).

Table 2 Classification of 6 cases of Klinefelter's syndrome with taurodontism

Case no.	Age	Karyotype	Affected teeth
(1) ST	30	XXY	8 7 8
(2) TU	30	XXY	7 7
(3) KK	21	XXY	7
(4) SK	32	XXY	7
(5) TM	23	XXY	7
(6) NY	39	XXY	7

mandibular second molars, while the remaining 2 were mandibular third molars (Table 2). The average crown/body ratio of the taurodont teeth of the Klinefelter's group (0.72 ± 0.04) and the controls (0.62 ± 0.03) were computed to compare the degree of taurodontism in the 2 groups. No significant difference was observed.

Pectus excavatum was noted in one Klinefelter's syndrome patient with taurodontism (Case 4, SK), while the remaining 30 patients examined did not have any clinically or radiographically visible structural defect of the thoracic cage, the spine, or the hand.

Discussion

It has been said that racial factors are important in the incidence of taurodontism, and that this incidence is low in modern Caucasians and high in Capoids, Australoids, and Mongoloids (Witkop, 1971).

The only study on the incidence of taurodontism among the Japanese was conducted on deciduous teeth. It was reported that the incidence is 0.5% to 1.6% and that it is higher among females than males (Daito and Hieda, 1971; Hitomi *et al.*, 1971). However, there has been no population study on permanent

teeth, and no indisputable data on the incidence of taurodontism among the Japanese is available.

In 1971, on the basis of reports by Law *et al.* (1969), Heida and Daito undertook a chromosomal analysis of a 4-year-old whose mandibular first deciduous molar was a taurodont tooth, and reported that in this case the karyotype was 47,XXY. Thus, the relation between Klinefelter's syndrome and taurodontism was noted even in Japan.

Keeler (1973) reported that in 6 cases of X chromosome aneuploidy (3 XXY, 1 XXXY, 1 XXYY and 1 XXXXY/XXXY), taurodont molars were observed in each. Stewart (1974) reported on 1 XXY and 1 XXXXY case, both with taurodontism, and speculated, on the basis of this and past studies, that the greater the number of X chromosomes, the greater the incidence and degree of taurodontism would be. Feichtinger and Rossiwall (1977) studied 8 individuals with X chromosome aneuploidy and found taurodontism in only 1 XXY/XXXY case. They also reported that, among 6 XXY cases and 1 XXX case, taurodontism was not observed. Based on these results, they concluded that there is some relation between additional X chromosomes and the appearance of taurodontism, but that taurodontism is not an obligatory finding in standard XXY Klinefelter's syndrome.

In the 31 XXY cases we studied, the incidence of taurodontism was approximately 20%, slightly less than that of gynaecomastia in patients with Klinefelter's syndrome. Thus, we believe this is one of the anomalies frequently associated with this condition.

We acknowledge the considerable assistance of Dr T. Abe, Department of Preventive Medicine, Kyoto Prefectural University of Medicine.

References

- Barker, B. C. W. (1976). Taurodontism: the incidence and possible significance of the trait. *Australian Dental Journal*, **21**, 272-276.
- Daito, M., and Hieda, T. (1971). Taurodont teeth in primary dentition. *Japanese Journal of Pedodontics*, **9**, 94-106.
- Feichtinger, C. H., and Rossiwall, B. (1977). Taurodontism in human sex chromosome aneuploidy. *Archives of Oral Biology*, **22**, 327-329.
- Hamner, J. E., Wintrop, C. J., Metro, P. S., and Ohio, C. (1964). Taurodontism. Report of a case. *Oral Surgery, Oral Medicine, and Oral Pathology*, **18**, 409-418.
- Hieda, T., and Daito, M. (1971). Klinefelter's syndrome in a case of taurodontism. *Japanese Journal of Pedodontics*, **9**, 79-85.
- Hitomi, K., Kato, K., Nagasaka, N., and Kurosu, K. (1971). A study of taurodontism in deciduous teeth. *Japanese Journal of Pedodontics*, **9**, 126-135.
- Keeler, C. (1973). Taurodont molars and shovel incisors in Klinefelter's syndrome. *Journal of Heredity*, **64**, 234-236.
- Keith, A. (1913). Problems relating to the teeth of earlier forms of prehistorical man. *Proceedings of the Royal Society of Medicine*, **6**, 103-119.
- Law, D. B., Lewis, T. M., and Davis, J. M. (1969). *An Atlas of Pedodontics*. Saunders, Philadelphia.
- Moorhead, P. S., Nowell, P. S., Batipps, D. M., and Hungerford, D. A. (1960). Chromosome preparations of leukocytes cultured from human peripheral blood. *Experimental Cell Research*, **20**, 613-616.
- Shaw, J. C. W. (1928). Taurodont teeth in South African races. *Journal of Anatomy*, **62**, 476-498.
- Stewart, R. E. (1974). Taurodontism in X chromosome aneuploidy syndromes. *Clinical Genetics*, **6**, 341-344.
- Witkop, C. J. (1971). Manifestations of genetic diseases in human pulp. *Oral Surgery, Oral Medicine, and Oral Pathology*, **32**, 278-316.

Requests for reprints to Dr Yosuke Komatz, Department of Urology, Faculty of Medicine, Kyoto University, Sakyo-ku, Kyoto 606, Japan.