TABLE Frequency of mothers pregnant with a male with male fetal cells in their peripheral blood. Data on interphase and metaphase counts are presented separately

	Interphases in uncultured cells	Interphases after PHA stimulation	Metaphases after PHA stimulation
Walknowska <i>et al</i> ⁴			19/21 cases
de Grouchy and Trébuchet ⁵			8/11
Zimmerman and Schmickel ⁶	No good correlation between sex and fluorescent bodies		
Schindler <i>et al</i> ⁷			4/9
Schröder and de la Chapelle ⁸	7/9	7/9	
Grosset <i>et al</i> ⁹		42/45	
Zilliacus <i>et al</i> ¹⁰	7/11		0/10

metaphases, with quinacrine staining, in the peripheral blood of pregnant women after stimulation with PHA, but no previous purification of lymphocytes. Whether these cells are lymphocytes or not has already been discussed earlier by Schröder and Herzenberg.¹²

M KIRSCH-VOLDERS

Vrije Universiteit Brussel,
Laboratorium voor Antropogenetika,
Pleinlaan 2, 1050 Brussels, Belgium

References

- Schröder J. Transplacental passage of blood cells. *J Med Genet* 1975;**12**:230-42.
- Turunen O, Lundqvist C, de la Chapelle A. Fetal cells in the maternal blood: differential effects of mitogens on maternal and fetal lymphocytes. *Excerpta Medica Int Cong Ser* No 397, 1976:162-3.
- Herzenberg LA, Bianchi DW, Schröder J, Cann HM, Iverson GM. Fetal cells in the blood of pregnant women: detection and enrichment by fluorescence-activated cell sorting. *Proc Natl Acad Sci USA*, 1979;**76**:1453-5.
- Walknowska J, Conte FA, Grumbach MM. Practical and theoretical implication of fetal/maternal lymphocyte transfer. *Lancet* 1969;**i**:1119-22.
- de Grouchy J, Trébuchet C. Transfusion foeto-maternelle de lymphocytes sanguins et détection du sexe du foetus. *Ann Genet (Paris)* 1971;**14**:133-7.
- Zimmerman A, Schmickel R. Fluorescent bodies in maternal circulation. *Lancet* 1971;**i**:1305.
- Schindler AM, Graf E, Martin-du-Pan R. Prenatal diagnosis of fetal lymphocytes in the maternal blood. *Obstet Gynecol* 1972;**40**:340-6.
- Schröder J, de la Chapelle A. Fetal lymphocytes in the maternal blood. *Blood* 1972;**39**:153-62.
- Grosset L, Barrelet V, Odartchenko N. Antenatal fetal sex determination in maternal blood during early pregnancy. *Am J Obstet Gynecol* 1974;**120**:60-3.
- Zilliacus R, de la Chapelle A, Schröder J, Tiilikainen A, Kohne E, Kleihauer E. Transplacental passage of foetal blood cells. *Scand J Haematol* 1975;**15**:333-8.
- Schröder J, Schröder E, Cann H. Fetal cells in the maternal blood. *Hum Genet* 1977;**38**:91-7.
- Schröder J, Herzenberg LA. Fetal cells in the maternal circulation: prenatal diagnosis by cell sorting using a fluorescence-activated cell sorter (FACS). In: Milunsky A, ed. *Hereditary disorders and the fetus. Diagnosis, prevention and treatment*. New York: Plenum, 1977.

Recurrence risk of neural tube defects

SIR,

With reference to Dr Seller's paper (page 245) on recurrence risks for neural tube defects (NTD) derived from a population of women presenting themselves for prenatal diagnostic tests,¹ I would like to confirm some of her conclusions from the experience in South Wales. Among the 4221 amniocenteses carried out between 1974 and 1980, the 927 carried out because the mother had had one or more previous children with an NTD led to the detection of 24 recurrences, giving an apparent recurrence risk of 1 in 39 (2.58%) (table). However, a follow-up of all the pregnancies revealed a further seven cases that were missed (in four because the lesion was closed) giving a total risk of recurrence derived from this population of 1 in 30 (3.34%),

TABLE Amniocenteses in South Wales 1974-1980

Amniocenteses	NTD detected				NTD detection rate	NTD undetected	Total NTD rate
	No	An	SBC	E			
Indication							
Previous NTD	927	14	10	—	1.39	7	1.30
Family history of NTD	596	3	2	—	1.119	—	1.119
Parent with NTD	40	—	—	—	—	—	—
Raised serum AFP	774	47	45	1*	1.8.3	5	1.7.9
Other CNS malformations	75	1	1	—	1.37	—	1.37
Anxiety	151	1	—	—	1.151	—	1.151
Cytogenetic	1523	1	3	—	1.508	3	1.218
Other	127	2	—	—	1.63	—	1.63
Total	4221	69	61	1	1.32	15	1.30

A, anencephaly; SBC, spina bifida cystica; E, encephalocele. *A case of Meckel-Gruber syndrome.

a very similar one to that obtained by Dr Seller. In those who had amniocentesis because of a family history of NTD (mostly with an affected second or third degree relative) the overall risk was 1 in 119 (0.83%), while in those where the amniocentesis was performed for cytogenetic indication the incidence of NTD turned out to be 1 in 218 (0.46%), suggesting a population risk of 4.6 per 1000 births. The latter would be in keeping with the fallen population incidence of NTD in the area from 7.67 per 1000 births between 1956 and 1962² and 7.0 per 1000 births between 1964 and 1966³ to just over 4 per 1000 births 12 years later.⁴

The risk of recurrence of 1 in 30 is substantially less than the 1 in 19.2 found in a large family study in South Wales reported in 1968.⁵ This lower risk must be a reflection not only of the considerably higher social class distribution of couples coming for amniocentesis than the general population of those who have a child with an NTD,⁶ but also of the reduced population incidence in recent years. It is probable that, in addition, some minor cases of spina bifida cystica and encephalocele may not have been reported in the follow up.⁷ With some caution this lower risk of recurrence in amniocentesis cases should be reflected in risks that could now be used in genetic counselling.

I wish to thank Dr Seller for letting me see the typescript of her paper.

K M LAURENCE

Department of Child Health,
Welsh National School of Medicine,
Heath Park,
Cardiff CF4 4XN

References

- ¹ Seller MJ. Recurrence risks for neural tube defects in a genetic counselling clinic population. *J Med Genet* 1981;**18**:245-8.
- ² Laurence KM, Carter CO, David PA. The major central nervous system malformations in South Wales. I. Incidence, local variation and geographical factors. *Br J Prev Soc Med* 1968;**22**:146-60.
- ³ Richards IDG, Lowe CR. Incidence of congenital defects in South Wales 1964-66. *Br J Prev Soc Med* 1971;**25**:59-64.
- ⁴ Laurence KM. Provision of population screening services for neural tube defects. *Health Trends* 1981;**13**: (in press).
- ⁵ Carter CO, David PA, Laurence KM. A family study of major central nervous system malformations in South Wales. *J Med Genet* 1968;**5**:81-106.
- ⁶ Laurence KM, Morris J. The effect of the introduction of prenatal diagnosis on the reproductive history of women at increased risk from neural tube defects. *J Prenatal Diagnosis* 1981;**1**:51-60.
- ⁷ Laurence KM. Prevention and prenatal diagnosis of neural tube defects. In: Persaud TVN, ed. *Advances in the study of birth defects*. vol 7. Lancaster: MTP Press, 1981.