- ¹² Bouvet JP, Leveque D, Bernetieres F, Gros JJ. Vascular origin of Poland syndrome? A comparative rheographic study of the vascularization of the arms of eight patients. *Eur J Pediatr* 1978;**128**:17-26.
- ¹³ Castilla EE, Paz JE, Orioli IM. Pectoralis major muscle defect and Poland complex. *Am J Med Genet* 1979;**4**: 263-9.
- ¹⁴ McGillivray BC, Lowry RB. Poland syndrome in British Columbia: incidence and reproductive experience of affected persons. *Am J Med Genet* 1977;**1**:65-74.
- ¹⁵ David TJ. Debendox does not cause the Poland anomaly. *Arch Dis Child* 1982;**57**:479-80.
- ¹⁶ Sugiura Y. Poland's syndrome. Clinico-roentgenographic study on 45 cases. *Congen Anom* 1976;**16**:17-28.
- ¹⁷ Gnamey D. Le syndrome de Poland (considerations etiologiques). *Lille Med* 1974;**19**:953-6.
- ¹⁸ Mohlbauer W, Wangerin K. Zur Embryologie und Ätiologie des Poland- und Amazonensyndromes. *Handchirurgie* 1977;**9**:147-51.

Craniosynostosis

SIR,

In their interesting paper on families with craniosynostosis, Professor Carter and co-workers¹ suggest that our higher incidence of apparent autosomal dominant coronal synostosis may result from our inclusion of patients with Saethre-Chotzen and other syndromes among our study group.² Certainly, if the minor hand anomalies to which we referred have the same aetiology as the craniosynostosis, then some families had 'private' syndromes. We agree that family number 1 has what the authors call the 'split face syndrome', and that it is like the family reported by Slover and Sujansky.³ However, in the interest of clarity, we would like to point out that we did not make a diagnosis of Saethre-Chotzen syndrome unless at least one family

member had the typical syndactyly. Therefore, as the title implied, our families with Saethre-Chotzen syndrome were not included in the paper and we do not believe that families 4, 8, or 10 had this condition.

The degree to which the reported heterogeneity of the craniosynostosis syndromes represent true genetic heterogeneity must await biochemical or linkage markers for the genes or both. In the meantime it remains true that patients who have craniosynostosis in association with other dysmorphic features, notably of the hands, are more at risk to represent a single gene mutation than are those with isolated craniosynostosis.

ALASDAIR HUNTER
Division of Genetics,
Children's Hospital of Eastern Ontario,
Ottawa, Ontario, Canada K1H 8L1.

References

- ¹ Carter CO, Till K, Fraser V, Coffey R. A family study of craniosynostosis, with probable recognition of a distinct syndrome. *J Med Genet* 1982;**19**:280-5.
- ² Hunter AGW, Rudd NL. Craniosynostosis. II. Coronal synostosis; its familial characteristics and clinical findings in 109 patients lacking bilateral polysyndactyly or syndactyly. *Teratology* 1977;**15**:301-10.
- ³ Slover R, Sujansky E. Fronto-nasal dysplasia with coronal synostosis in three sibs. *Birth Defects* 1979; **XI**(5B):75-83.

Pericentric inversion of chromosome 13

SIR,

In 1972 a paper from our laboratory described a large family with a pericentric inversion of chromosome 13, leading to a duplication deficiency

FIG 1 Pedigree of family.

FIG 2 (a) Chromosome 13 from a carrier. The inverted chromosome is shown with both G and NOR banding. (b) The recombinant chromosome 13, type 1. (G banding.) (c) The recombinant chromosome 13, type 2. (Conventional, G, and NOR banding.)

associated with congenital malformations in three family members.¹

Since then all carriers have been followed and fetal cells cultured from all known pregnancies, including two spontaneous abortions (fig 1). The inversion and breakpoints have been confirmed with banding techniques (fig 2a, b).

The second type of recombinant chromosome postulated in our paper has been found in one spontaneous abortion (fig 2c).

We wish to add this new information and show the revised pedigree.

HALLA HAUKSDÓTTIR*, ÁSTRÓS ARNARDÓTTIR*,
MARGRÉT STEINARSDÓTTIR*,
ELÍN GUÐMUNDSDÓTTIR*, SÆVAR HALLDÓRSSON†,
AUÐÓLFUR GUNNARSSON‡, AND
MARGARETA MIKKELSEN§

*The Chromosome Laboratory, Department
of Pathology, University of Iceland, †Department
of Paediatrics, St Joseph's Hospital,
‡Department of Gynaecology and Obstetrics,
Landspítalinn, 101 Reykjavík, Iceland; and
§John F Kennedy Institutttet, G1 Landevej 7-9,
Glostrup, Copenhagen, Denmark.

Reference

- Hauksdóttir H, Halldórsson S, Jensson Ó, Mikkelsen M, McDermott A. Pericentric inversion of chromosome No 13 in a large family leading to duplication deficiency causing congenital malformations in three individuals. *J Med Genet* 1972;9:413-21.

Adjacent 2 translocation involving 13q and 21q

SIR,

The article in *Journal of Medical Genetics* entitled 'Adjacent 2 translocation involving 13q and 21q' (1982;19:314-5) states that this case is the first involving chromosomes 13 and 21 with an adjacent 2 disjunction in the infant and a balanced reciprocal translocation involving the long arms of chromosomes 13 and 21 in the mother.

We have studied a female carrier of a translocation in which the long arms of chromosomes 13 and 21 were involved.¹ Identification with G banding (GTG) was not conclusive enough to enable us to establish definite breakpoints but, together with R banding (RBA), would suggest the following karyotype: 46,XX,t(13;21)(q21;q21).

The offspring of this woman suggest that this translocation carries a high risk. The first child died just after delivery in another hospital without cytogenetic study. The second child had dysmorphic features with partial trisomy 13 and partial monosomy 21 owing to an adjacent 2 meiotic disjunction. His karyotype was 46,XY,-21,+der(13),t(13;21)(q21;q21). The third child had the phenotype of Down's syndrome because of a 3:1 segregation and his karyotype was 47,XY,+21,t(13;21)(q21;q21).

F PRIETO AND L BADIA
Sección de Genética,
Servicio de Hematología y Hemoterapia,
Hospital Infantil,
Ciudad Sanitaria de la Seguridad Social 'La Fe',
Valencia, Spain.

Reference

- Prieto F, Badia L, Asensi F, Roques V. Two reciprocal translocations t(9p+;13q-) and t(13q-;21q+). A study of the families. *Hum Genet* 1980;54:7-11.

Pyloric stenosis: children vs sibs

SIR,

We have reported¹ findings in the relatives of patients with pyloric stenosis which showed, for female patients, more children affected than sibs. This is unexpected on a simple multifactorial threshold model and has led us and others to speculate whether there may be some direct maternal effect, though there is no indication, on the small series available, that maternal half-sibs are more often affected than paternal half-sibs.² We have continued to follow the children of the female patients born between 1933 and 1949 (but not those born between 1921 and 1932, who are not likely to have had further children) and the relatively high risk to children has now disappeared. The data on