linkage study using two cloned DNA sequences. *J Med Genet* 1983;**20**:249-51.

³ Kosambi DD. The estimation of map distances from recombination values. *Ann Eugen* 1944;**12**:172-5.

⁴ Rao DC, Morton NE, Lindsten J, Hultén M, Yee S. A mapping function for man. *Hum Hered* 1977;**27**:99-104.

⁵ Renwick JH. Progress in mapping human autosomes. *Br Med Bull* 1969;**25**:65-73.

Haemoglobin O Arab (B121 Glu-Lys) in Turkish Cypriot population

SIR,

Hb O Arab (B121 Glu-Lys) has been reported in different ethnic groups including Sudanese, Arabs, American Negroes, Bulgarians, and recently in Turkish people.¹⁻⁷ It has not previously been reported in the Turkish Cypriot population.

We investigated blood samples of 1365 apparently healthy Turkish Cypriots by cellulose acetate electrophoresis.⁸ When an abnormal haemoglobin was identified, citrate agar electrophoresis was performed.⁹ Structural analysis was performed when indicated.¹⁰

Two haemoglobin O Arab traits and one Hb O Arab β -thalassaemia combination were detected in three different families. Further study of 24 subjects from these three families revealed 13 Hb O Arab traits and two Hb O Arab β -thalassaemia combinations. It was observed that the subjects with Hb O Arab had negroid characteristics (negroid Turkish Cypriots have the same physical characteristics as African Negroes; they represent less than 1% of all Turkish Cypriots). We investigated a further 19 subjects with negroid characteristics, two with Hb S trait, one with Hb H disease, and five with β -thalassaemia trait, but no further Hb O Arab traits were found in these subjects.

Hb O Arab was first detected in an Arab living in Israel, then in Egypt, Aden, Bulgaria, Rumania, Hungary, among American Negroes, and recently in Turkey.¹⁻⁷ It is believed to have originated in the Sudan and to have spread from there to West Africa and to many countries once occupied by or in close contact with the Ottoman Empire.^{6,7}

When the history of the Ottoman Empire is considered the occurrence of Hb O Arab in Cyprus is not surprising. In 1570 the Ottomans conquered Cyprus and Anatolian Turks migrated to the island. The island was ruled by the Ottomans from the 16th to the end of the 19th century and during this period many people from Sudan and Egypt emigrated to Anatolia, Syria, Lebanon, and Cyprus.¹¹ Identification of Hb O Arab in Turkish Cypriots may indicate an admixture of African blood in the past.

Moreover, some Turkish Cypriots show negroid physical characteristics.

We are grateful to T H J Huisman from the Medical College of Georgia, USA, for the analysis of abnormal variants.

ŞÜKRÜ CİN, NEJAT AKAR, AYTEN ARCASOY,
AYHAN O ÇAVDAR, AND SEHER DEDEOĞLU
*Pediatric Haematology and
Oncology Research Unit,
Medical School of Ankara University,
Ankara, Turkey.*

References

- Vella F, Beale D, Lehmann H. Hemoglobin O Arab in Sudanese. *Nature* 1966;**209**:308-9.
- Kamel KA, Hoerman KC, Awany AY. Hemoglobin $\alpha_2\beta_2$ 121 Lys: chemical identification in an Egyptian family. *Science* 1967;**156**:397-8.
- Ibrahim SA, Mustafa D. Sickle-cell haemoglobin O disease in a Sudanese family. *Br Med J* 1967;**3**:715-7.
- Milner PF, Miller C, Grey R, Seakins M, DeJong WW, Went LN. Hemoglobin O Arab in four negro families and its interaction with hemoglobin S and hemoglobin C. *N Engl J Med* 1970;**283**:1417-25.
- Kantchev KN, Tcholakov B, Baglioni C, Colombo B. Hemoglobin O Arab in Bulgaria. *Nature* 1965;**205**:187-8.
- Lehmann H, Huntsman RG. *Man's haemoglobins*. Amsterdam: North Holland, 1974.
- Prozorova-Zamani V, Özsoylu Ş, Aksoy M, et al. Hb E and Hb E like variants in individuals from Turkey. *Hemoglobin* 1981;**5**:743-8.
- Cin Ş, Akar N, Arcasoy A, Dedeoğlu S, Çavdar AO. Kıbrıs Türk Toplumunda talassemia, anormal hemoglobinler ve G6PD enzim eksikliği insidansı. *Doğa* 1983;**7**:21-36.
- Schneider RG. Identification of hemoglobins and hemoglobinopathies by electrophoresis on cellulose acetate plates impregnated with citrate agar. *Clin Chem* 1974;**20**:1111-5.
- Huisman THJ, Jonxis JHD. *The hemoglobinopathies. Techniques of identification*. New York: Marcel Dekker, 1977.
- Kutay C. *Türkiye. İstiklal ve Mücadeleleri Tarihi (History of Turkish nation)*. Istanbul: Istanbul Matbaası, 1958.

Incidence of pyloric stenosis in British Columbia over a 12 year period

SIR,

In a recent letter, Carter and co-workers¹ mentioned that the incidence of pyloric stenosis (infantile hypertrophic pyloric stenosis, IHPS) may be falling in the general population. We would like to comment on a recent study² done in British Columbia (BC) which investigated changes and seasonal variation in the incidence of IHPS over a 12 year period.

The BC study² was based on a cohort of 1234 livebirths with IHPS (988 males, 246 females) born