Dr Andrew Czeizel has sent us a copy of the Hungarian medical weekly *Orvosi Hetilap* of 10 February 1985 (pp 343–4) in which there is an obituary of Professor C O Carter written by Dr Czeizel, saying how much they miss him in Hungary.

Lateral reading 12

MALE SEXUAL DIFFERENTIATION IN MICE LACKING H-Y ANTIGEN (McLaren A, *et al.* *Nature* 1984;**312**:552)

“The sexual phenotype of an adult mammal depends on whether the fetal gonad has differentiated as a testis or as an ovary. Because individuals of XY or XXY sex chromosome constitution develop as males, while XX and XO individuals develop as females, the presence of a Y chromosome seems normally to be required for testis differentiation and its absence to be necessary for differentiation of an ovary. The nature of the hypothetical Y dependent substance responsible for masculinisation of the indifferent gonad has been a matter for debate. A male specific transplantation antigen, H-Y, has been known for many years, and more recently a serologically detected antigen, also male specific, has been reported. Those who believe that the two are antigenically distinct refer to the latter as SDM (serologically detected male) antigen, but many refer to both as H-Y antigen. The hypothesis that H-Y is itself the Y dependent testis inducer, although supported by little or no direct evidence, is economical and hence attractive. H-Y antigen is frequently stated to be the substance responsible for primary sex determination. We report here that H-Y is absent from certain mice that develop testes and are of indisputably male phenotype, hence this transplantation antigen is unlikely to be responsible for testis determination.”

PREVENTION OF MELANOMA BY RECOGNITION OF ITS PRECURSORS (Greene *et al.* *N Engl J Med* 1985;**312**:91–7. Fitzpatrick *et al.* *N Engl J Med* 1985;**312**:115–6)

It is well known that the incidence of cutaneous malignant melanoma is rising rapidly throughout the world. It is also well established that there are helpful characteristics of pigmented lesions to decide whether they are likely to be benign or malignant. Thus, uniform tan or brown lesions with smooth or regular borders and a stable growth

history indicate a benign condition whereas variegated dark brown or especially black, grey, or blue lesions with irregular borders suggest melanoma.

Now the emphasis has shifted to the recognition of people with an increased risk of melanoma by reason of their family history or the presence of one or more precursor lesions. These are the subject of a paper in the *New England Journal of Medicine* of 10 January 1985 (Greene *et al.*) and an editorial (Fitzpatrick *et al.*).

A large number of dermatological names is given to the at risk lesions and there are coloured plates of these together with those of the innocent types. With them are detailed descriptions of the lesions shown—forming an atlas of great practical use.

ABNORMAL METABOLISM OF γ TRACE ALKALINE MICROPROTEIN (Grubb A, *et al.* *N Engl J Med* 1984;**311**:1547)

“In 1935 Arnason described 10 families with a high incidence of cerebral haemorrhage and concluded that a hereditary form of the disease was present in these families. Further clinicopathological investigations of the disease revealed an autosomal dominant inheritance and a connection between the disease and a special form of amyloidosis confined to the cerebral vasculature. Recently, the fibrillar components of the amyloid deposits in the cerebral arteries of the three patients were isolated and shown to consist of large fragments of the alkaline microprotein γ trace. This protein is present in a number of neuroendocrine cells, and its concentration in the cerebrospinal fluid is 5.5 times that in the plasma of healthy males.

In order to investigate whether patients with hereditary cerebral haemorrhage with amyloidosis have an abnormal metabolism of γ trace in the central nervous system, we measured concentrations of γ trace in cerebrospinal fluid from nine patients with the disease, 44 healthy subjects, and 60 patients with other cerebrovascular disorders, multiple sclerosis, or prolapsed intervertebral discs. The patients with hereditary cerebral haemorrhage with amyloidosis had the lowest values for γ trace (mean 2.5 mg/l; range 1.7 to 3.5), and the mean value was only about one third of that in the reference groups (mean 7.6 mg/l; range 3.1 to 18.5). The cerebrospinal fluid concentration of another microprotein, β_2 microglobulin, with about the same size as γ trace,

did not differ between the groups. No structural differences between the γ trace in spinal fluid from patients with hereditary cerebral haemorrhage with amyloidosis and control subjects could be revealed. The data are therefore consistent with the hypothesis that the basic defect in this disease is an abnormal metabolism of γ trace, and they suggest that the abnormality resides in the catabolic processing of γ trace. Furthermore, the results strongly indicate that the concentration of γ trace in the cerebrospinal fluid can be used as an index of high sensitivity and specificity in the diagnosis of this lethal disorder."

LONG TERM TREATMENT OF CEREBROTENDINOUS XANTHOMATOSIS WITH CHENODEOXYCHOLIC ACID (Berginer VM, Salen G, Shefer S. *N Engl J Med* 1984;311:No 26)

Cerebrotendinous xanthomatosis is a rare, recessively inherited lipid storage disease. The major clinical features are tendon xanthomas, cataracts, dementia, pyramidal paresis, cerebellar ataxia, abnormalities on the electroencephalogram and cerebral computed axial tomographic scan, premature atherosclerosis, endocrine hypofunction, and pulmonary dysfunction, associated with increased levels of plasma and tissue cholestanol (5 α -dihydro derivative of cholesterol) and defective bile acid synthesis. Defective synthesis of bile acid results in the virtual absence of chenodeoxycholic acid in the bile and the excretion of large amounts of bile acid precursors (bile alcohols) conjugated with glucuronic acid in bile and urine. In this paper the authors describe the favourable outcome of treating 17 patients with chenodeoxycholic acid and conclude that long term therapy with chenodeoxycholic acid may correct the biochemical abnormalities and arrest and possibly reverse the progression of cerebrotendinous xanthomatosis. There is an editorial on the subject on p 1694 of the same issue.

METABOLIC OXIDATION PHENOTYPES AS MARKERS FOR SUSCEPTIBILITY TO LUNG CANCER (Ayesh R, Idle JR, Ritchie JC, Crothers MJ, Hetzel MR. *Nature* 1984;312:169-70)

The authors have previously published a series of 106 bronchial carcinoma patients who had been studied by means of a debrisoquine hydroxylation phenotyping test and who were compared with 258 healthy subjects. In the present article 245 patients (all smokers) with histologically proven bronchial cancer have been studied, and their results

compared with those obtained in 234 smokers with chronic airflow limitation but without evidence of carcinoma.

The debrisoquine hydroxylation test yields the 'metabolic ratio', that is, the amount of unchanged debrisoquine excreted in the urine during eight hours following a standard oral dose divided by the amount of the main metabolite 4-hydroxy debrisoquine in the same sample. This ratio exhibits a bimodal frequency distribution curve in the British population and the 9% of persons with values higher than about 12.6 ('poor metabolisers') are autosomal Mendelian recessives.

The authors describe three findings. (1) The control group yielded a frequency distribution closely similar to that given by healthy volunteer subjects. (2) There were significantly fewer 'poor metaboliser' phenotypes among the cancer patients than among the controls. (3) The frequency distribution curve in extensive metabolisers is markedly shifted towards lower ratios in cancer patients as compared with controls.

The big difficulty with interpreting these data is what might be termed the 'epiphenomenon problem'. Are the changes indicative that a selected set of genotypes within the population develops cancer, or do they reflect changes in metabolism which arise as a result of having cancer?

The authors themselves discuss this problem. The arguments brought forward against the epiphenomenon hypothesis are not fully convincing. Many bronchial carcinomas have endocrine activity which might explain the observed phenomena. The authors do not state whether the total recoveries of drug plus metabolites are the same in the cancer and control groups. If less unchanged drug but the same amount of metabolite were excreted this would cause a lowering of the observed ratio. Endocrine influences might affect the disposition of debrisoquine. The account does not make it quite clear whether patients were tested before or after chest surgery, but by implication it appears to be the former. In that case an interesting avenue is unexplored, that is, to compare what happens after tumour resection (in persons in whom there is no evidence of metastases) with the preoperative result. Another possibly interesting facet is unreported, namely the frequency distribution of the ratio in different histological types of cancer.

If these criticisms can be satisfactorily answered then the authors may have gone some of the way towards providing an affirmative answer to the question in the title of their original publication. The idea that persons who are more able to oxidise foreign compounds may be more prone to develop cancer is appealing because it is known that hydro-

carbons require oxidation to become active carcinogens. This line of enquiry need not now be restricted to the 'debrisoquine' polymorphism because an analogous but independent hydroxylation polymorphism can be detected with the drugs mephenytoin and nirvanol.

(Summarised by Dr D A Price Evans)

LIVER TRANSPLANTATION TO PROVIDE LOW DENSITY LIPOPROTEIN RECEPTORS AND LOWER PLASMA CHOLESTEROL IN A CHILD WITH HOMOZYGOUS FAMILIAL HYPERCHOLESTEROLAEMIA (Bilheimer DW, *et al.* *N Engl J Med* 1984;311**:1658)**

A 6 year old girl with severe hypercholesterolaemia and atherosclerosis had two defective genes at the low density lipoprotein (LDL) receptor locus, as determined by biochemical studies of cultured fibroblasts. One gene, inherited from the mother, produced no LDL receptors; the other gene, inherited from the father, produced a receptor precursor that was not transported to the cell surface and was unable to bind LDL. The patient degraded intravenously administered ^{125}I -LDL at an extremely low rate, indicating that her high plasma LDL cholesterol level was caused by defective receptor mediated removal of LDL from plasma. After transplantation of a liver and a heart from a normal donor, the patient's plasma LDL cholesterol level declined by 81% from 988 to 184 mg/dl. The

fractional catabolic rate for intravenously administered ^{125}I -LDL, a measure of functional LDL receptors in vivo, increased by 2.5-fold. Thus, the transplanted liver, with its normal complement of LDL receptors, was able to remove LDL cholesterol from plasma at a nearly normal rate.

The authors conclude that a genetically determined deficiency of LDL receptors can be largely reversed by liver transplantation.

BONE MARROW TRANSPLANTATION IN THE MUCOPOLYSACCHARIDOSIS TYPE VI SYNDROME (Krivit *et al.* *N Engl J Med* 1984;311**: 1606-11)**

In this paper, Krivit *et al* report the results of bone marrow transplantation in a 13 year old girl with the above syndrome. The procedure provided a source of enzymatically normal cells which altered for the better the metabolic and clinical course of the disease. The paper is followed (pp 1629-30 of the same issue) by an editorial by Dr Barranger on marrow transplantation in genetic disease in general. The conclusions seem rather ambiguous but the main message is that neurological syndromes are not likely to benefit. Additionally, it is pointed out that the lessons learnt from bone marrow transplantation will forecast the results that may be expected from gene transfer in the future.

C A CLARKE