We thank Dr Ahmed Al-Ansari for endocrinological assessment.

References

- ¹ Shoval AR. Mixed gonadal dysgenesis. In: Rashad MN, Morton WRM, eds. *Genital anomalies*. Springfield, Illinois: Charles C Thomas, 1969:476-89.
- ² Jones HWJ, Scott WW. *Hermaphroditism, genital anomalies and related endocrine disorders*. Baltimore: Williams & Wilkins, 1971:290-317.
- ³ Davidoff F, Federman DD. Mixed gonadal dysgenesis. *Pediatrics* 1973;**52**:725-42.
- ⁴ Simpson JL. *Disorders of sexual differentiation: etiology and clinical delineation*. New York: Academic Press, 1976:237-9.
- ⁵ Al-Awadi SA, Cuschieri A, Farag TI, et al. Mixed gonadal dysgenesis and sex chromosome mosaicism with multiple cell lines including structural abnormalities of the Y chromosome. *Clin Genet* 1983;**23**:172-6.
- ⁶ de la Chapelle A. Sex chromosome abnormalities. In: Emery AEH, Rimoin DL, eds. *Principles and practice of medical genetics*. Edinburgh: Churchill Livingstone, 1983:193-215.
- ⁷ Al-Awadi SA, Farag TI, Naguib KK, et al. A five year experience with gonosomal abnormalities in Kuwait. *J Kwt Med Assoc* 1984;**18**:225-32.
- ⁸ Klebs E. *Handbuch der Pathologischen Anatomie*. Berlin: Hirschwald, 1876.
- ⁹ Harnden DG, Armstrong CN. The chromosomes of a true hermaphrodite. *Br Med J* 1959;**ii**:1287.
- ¹⁰ Hungerford DS, Donnelly AJ, Nowell PC, Beck S. The chromosome constitution of a human phenotypic intersex. *Am J Hum Genet* 1959;**2**:215-36.
- ¹¹ Gartler SM, Waxman SH, Giblett E. An XX/XY human hermaphrodite resulting from double fertilization. *Proc Natl Acad Sci USA* 1962;**48**:332-5.
- ¹² Van Niekerk WA. *True hermaphroditism*. New York: Harper, 1974.
- ¹³ Manuel MA, Allie A, Jackson WPU. A true hermaphrodite with XX/XY chromosome mosaicism. *S Afr Med J* 1965;**39**:414-9.
- ¹⁴ Massimo L, Vianello MG. Chromosome studies in two cases of intersexuality. *Human Chromosome Newsletter* 1965;**15**:15-16.
- ¹⁵ Race RR, Sanger R. *Blood groups in man*. 6th ed. Oxford: Blackwell, 1975:531-9.
- ¹⁶ Fitzgerald PH, Brehaut LA, Shannan FT, Angus HE. Evidence of XX/XY sex chromosome mosaicism in a child with true hermaphroditism. *J Med Genet* 1970;**7**:383-8.
- ¹⁷ Park JJ, Jones HW Jr, Bias WG. True hermaphroditism with ambiguous external genitalia. *Am J Obstet Gynecol* 1970;**108**:1197-205.
- ¹⁸ Kakati S, Sharma T, Udupa KN, Chaudhuri SPR. A true hermaphrodite with XX/XY mosaicism. *Ind J Med Res* 1971;**59**:104-6.
- ¹⁹ Benirschke K, Naftalin F, Gittes R, Khuder G, Yen SSC, Allen FH. True hermaphroditism and chimerism. *Am J Obstet Gynecol* 1972;**113**:449-58.
- ²⁰ Grace JH. A white hermaphrodite in South Africa. *S Afr Med J* 1973;**47**:1553-4.
- ²¹ Shanfield I, Young RB, Hume DM. True hermaphroditism with XX/XY mosaicism: report of a case. *J Pediatr* 1973;**83**:471-3.
- ²² de la Chapelle A, Schröder J, Rantanen P, et al. Early fusion of two human embryos. *Ann Hum Genet* 1974;**38**:63-75.
- ²³ De Marchi M, Carbonara AO, Carozzi F, et al. True hermaphroditism with XX/XY chromosome mosaicism: report of a case. *Clin Genet* 1976;**10**:265-72.
- ²⁴ Fitzgerald PH, Donald RA, Kirk RL. A true hermaphrodite dispermic chimera with 46,XX and 46,XY karyotypes. *Clin Genet* 1979;**15**:89-96.
- ²⁵ Dewald G, Haymond MW, Spurbeck JL. Origin of 46,XX/46,XY chimerism in a human true hermaphrodite. *Science* 1980;**207**:321-3.
- ²⁶ Shah VC, Krishna Murthy DS, Sabita Roy, Contractor PM, Shah VA. True hermaphrodite 46,XX/46,XY: clinical, cytogenetic and histopathological studies. *Ind J Pediatr* 1982;**49**:885-90.
- ²⁷ Borgoankar DS. *Chromosomal variation in man. A catalog of chromosomal variants and anomalies*. 3rd ed. New York: Alan R Liss, 1980:483-5.
- ²⁸ de Grouchy J, Turleau C. *Clinical atlas of human chromosomes*. 2nd ed. New York: John Wiley, 1984:406-8.
- ²⁹ Schinzel A. *Catalogue of unbalanced chromosome aberrations in man*. Berlin: Walter de Gruyter, 1984:812-3.
- ³⁰ Tippet P. Human chimeras. In: *Chimeras in developmental biology*. London: Academic Press, 1984:165-78.
- ³¹ Ford CE. Mosaicism and chimeras. *Br Med Bull* 1969;**25**:104-9.

Correspondence and requests for reprints to Dr Patricia Tippet, MRC Blood Group Unit, Wolfson House, University College London, 4 Stephenson Way, London NW1 2HE.

Interstitial del(13)(q21-3q31) associated with psychomotor retardation, eczema, and absent suck and swallowing reflex

P J PEET, R RODRIGUES PEREIRA, J O VAN HEMEL, AND A J M HOOGEBOOM

Department of Paediatrics, St Clara Hospital, and Department of Clinical Genetics, Erasmus University, Rotterdam, The Netherlands.

SUMMARY A patient with a deletion (13)(q21-3q31) showed only eczema and ab-

sent suck and swallowing reflex, in contrast to other well documented cases with a similar deletion. Apparently there is wide clinical variability in patients with deletions in this area.

Received for publication 24 November 1986.
Accepted for publication 28 November 1986.

Cases with partial deletions of the long arm of chromosome 13 constitute a clinically and cytogenetically heterogeneous group. We describe a patient with a $\text{del}(13)(\text{q}21\cdot3\text{q}31)$ and compare our findings with well documented cases of deletions in the same region.

Case report

The proband, a boy, was the fourth child of healthy, unrelated parents. After an uneventful pregnancy the infant was born at term weighing 2635 g. In the neonatal period he was observed in another hospital for one week for possible sepsis. Three months later he was admitted because of failure to thrive and feeding difficulties.

He was a small boy (weight 3720 g, length 54 cm, head circumference 37 cm) with extensive eczema (fig 1). Retardation of psychomotor development, muscular hypotonia, weak tendon reflexes, absent suck and swallowing reflex, and inability to follow with the eyes were present. He had a partly depigmented right iris and low set, normal ears. No cardiac murmur was audible. There were recurrent

FIG 2 GTG and RBA banded chromosome 13.

respiratory and urinary tract infections. Radiography revealed no urinary tract anomalies. A CT scan of the brain was normal.

CYTOGENETIC STUDIES

GTG and RBA banded chromosomes from blood lymphocytes and skin fibroblasts showed a $46,\text{XY},\text{del}(13)(\text{pter}\rightarrow\text{q}21\cdot3::\text{q}31\rightarrow\text{qter})$ karyotype in all metaphases analysed (fig 2).

Both parents had normal chromosomes.

FIG 1 The proband at five months.

FIG 3 Schematic diagram of $\text{del}(13)(\text{q}21\text{--}\text{q}31)$.

TABLE Clinical and cytogenetic features in 13q monosomy.

	Present case	Reference				
		1	2	3	4	5
Extent of deletion	21:3→31	21→31	21→31	21:3→31.2	14→31	21→22
Sex	M	F	F	F	M	F
Birth weight <2800 g	+	+	+	+	+	
Psychomotor retardation	+	+	+	+		+
Hypotonia	+			+		+
Generalised dermatitis	+					
Microcephaly	+		+	+		+
Absent suck and swallowing reflex	+					
Partial iris depigmentation	+					
Low set or malformed ears	+	+	+	+	+	
Short neck	-	+	+	+		
Prominent nasal bridge	-		+	-		+
Broad nasal bridge	-		+	+	+	
Hypertelorism	-			+	+	
Microphthalmia	-			-		+
Epicanthus	-			+		
Retinoblastoma	-			-	+	-
Hypoplastic 5th digit + clinodactyly	-			+		-
Cardiac defect	-		+			-
Multiple intestinal malformations	-				+	-

Discussion

Clinical symptoms associated with 13q deletions are variable. Comparison of cases with deletions in the same region as our patient (table)¹⁻⁵ does not help to establish a definite clinical pattern⁶; psychomotor retardation seems to be the most constant finding.

Gross intestinal malformations are possibly associated with deletions in the region q13→q31,^{4,7} and deletion of band 13q14 is associated with retinoblastoma.⁷ Our case shows absence of suck and swallowing reflexes which has not previously been reported, and extensive eczema; dermatitis has been reported in one other case.⁷

More distal deletions may give a recognisable clinical picture. Deletion of band 13q34 is probably responsible for the craniofacial dysmorphism classically described in the 13q monosomy syndrome.⁶ Thumb hypoplasia is variably present in deletion 13q34.⁶ The wide phenotypical variability is also demonstrated by the normal phenotype in a case with a deletion of band 13q21.⁸ This was explained by the fact that this band is one of the latest replicating, which may indicate that it carries genetic material which is not transcriptionally active.

Establishing a definite phenotype for deletions in this region, if at all possible, will certainly require more cases studied by high resolution banding techniques.

The authors would like to thank Professor M F Niermeijer for his critical comments and helpful suggestions.

References

- Nielsen J, Homma A, Christiansen F, Rasmussen K, Saldana Garcia P. Deletion long arm 13. *Hum Genet* 1977;37:339-45.
- Serena-Lungarotti M, Calabro A, Mariotti G, Mastroiacovo P, Provenzano S, Dallapiccola B. Interstitial deletion 13q syndrome: a report on two unrelated patients. *Hum Genet* 1979;52:269-74.
- Weisswichter P, Stögmänn W. Interstitielle Stückverlust am Chromosom Nr 13: Ein neues Syndrom? *Klin Pädiatr* 1979;191:318-24.
- Yunis E, Zuñiga R, Ramirez E. Retinoblastoma, gross internal malformations and deletion 13q14→q31. *Hum Genet* 1981;56:283-6.
- Noel B, Quack B, Rethoré MO. Partial deletions and trisomies of chromosome 13: mapping of bands associated with particular malformations. *Clin Genet* 1976;9:593-602.
- Niebuhr E. Partial trisomies and deletions of chromosome 13. In: Yunis JJ, ed. *New chromosomal syndromes*. New York: Academic Press, 1977:273-99.
- Yunis JJ, Ramsay N. Retinoblastoma and subband deletion of chromosome 13. *Am J Dis Child* 1978;132:161-3.
- Couturier J, Morichon-Delvallez N, Dutrillaux B. Deletion of band 13q21 is compatible with normal phenotype. *Hum Genet* 1985;70:87-91.

Correspondence and requests for reprints to Dr R Rodrigues Pereira, Department of Paediatrics, St Clara Ziekenhuis, Olympiaweg 350, 3078 HT Rotterdam, The Netherlands.