Sex reversal in a child with a 46,X,Yp+ karyotype: support for the existence of a gene(s), located in distal Xp, involved in testis formation

Tsutomu Ogata, J Ross Hawkins, Anne Taylor, Nobutake Matsuo, Jun-ichi Hata, Peter N Goodfellow

Human Molecular Genetics Laboratory, Imperial Cancer Research Fund, PO Box 123, Lincoln's Inn Fields, London WC2A 3PX.

T Ogata
J R Hawkins
A Taylor
P N Goodfellow

Department of Paediatrics, Keio University School of Medicine, Tokyo 160, Japan.
N Matsuo

Department of Pathology, Keio University School of Medicine, Tokyo 160, Japan.
J Hata

Correspondence to Dr Ogata.

Received 19 June 1991.
Revised version accepted 30 September 1991.

Abstract

We report on a sex reversed Japanese child with a 46,X,Yp+ karyotype, minor dysmorphic features, and no testicular development. The Yp+ chromosome was derived by translocation of an Xp fragment (Xp21-Xp22.3) to Yp11.3. This has resulted in deletion of distal part of the Y chromosome pseudoautosomal region (DXYS15-telomere) and duplication of the X specific region (DXS84-PABX) and proximal part of the pseudoautosomal region (MIC2-DXYS17). No deletion of the Y specific region was detected nor was any mutation found in SRY. Cytogenetic analysis suggests that the proximal part of the Xp fragment is the most distal part of the short arm of the Yp+ chromosome (Xp21→Xp22.3::Yp11.3→Yqter). No chromosomal mosaicism was detected.

These results are similar to previous reports of sex reversal in four subjects with a 46,Y,Xp+ karyotype. We conclude that the sex reversal is a direct, or indirect, consequence of having two active copies of the distal part of Xp and may indicate the presence of a gene(s) which

acts in the testis determination or differentiation pathway.

Sex determination in man, and other mammals, is chromosomally based: males have an X and a Y chromosome, females have two X chromosomes. Correlation between phenotype and karyotype in subjects with unusual sex chromosome constitutions has shown that the Y chromosome carries a gene, TDF (testis determining factor), essential for testis formation and male sex determination. Recent molecular analysis of the genomes of XX males and XY females has provided strong evidence that the Y located gene SRY is TDF,¹⁻³ and this has been confirmed by transgenic mice experiments.⁴ However, not all cases of sex reversal can be explained by alterations in SRY and it would be predicted that both 'gain of function' and 'loss of function' mutations in other genes in the sex determination pathway may cause sex reversal. Another theoretical possibility is that dosage of critical genes may affect sex determination. In this report, we describe a patient with sex reversal and an extra fragment of Xp, which is translocated to Yp.

Case report

A phenotypic female child was born to non-consanguineous parents at 40 weeks of gestation after an uncomplicated pregnancy and delivery. The parents and older sister were clinically normal. The birth length was 48 cm and weight 2900 g. From birth, she was admitted frequently to a local hospital with recurrent fever.

At 2 years 3 months of age, the patient was referred to Keio University Hospital because of high fever, lymphadenopathy, and erythematous rashes. Physical examination showed a weak child with marked hypotonia. External genitalia were feminine and there was no clitoromegaly or labial fusion. Dysmorphic features included frontal bossing, antimongoloid slant, large, low set ears with thick auricular folds, and cleft palate. Psychomotor development was severely retarded. Laboratory studies showed anaemia (Hb 9.5 g/dl), thrombocytopenia ($7.1 \times 10^9/\mu\text{l}$), positive antinuclear antibody (2560X, homogeneous pattern) and anti-DNA antibody (320X), decreased complement level (C3 0.3 g/l), and immunoglobulin A (IgA) deficiency (<0.01 g/l). After a diagnosis of systemic lupus erythematosus (SLE),

Figure 1 Histological finding of the gonad showing gonadoblastoma and ovarian stroma (haematoxylin-eosin). Nests composed of germ cells and immature sex cord derivatives contain rounded hyaline bodies. These nests are embedded in immature ovarian stroma. No testicular structure is seen.

The copy number of each locus.

Locus (probe)	Enzyme	RFLP	Patient	Sister	Father	Mother	Reference
Pseudoautosomal region							
<i>DXYS14</i> (29C1)	<i>TaqI</i>	+	1	2	2	2	9
<i>DXYS15</i> (113D)	<i>TaqI</i>	+	1	2	2	2	10
<i>DXYS17</i> (601)	<i>TaqI</i>	+	3	2	2	2	11
<i>MIC2</i> (19B)	<i>TaqI</i>	—	3 (1.47)	2 (1.08)	2 (1.05)	2 (0.98)	12
X specific region							
<i>PABX</i> (Hf0.2)	<i>SstI</i>	—	2 (1.35)	2 (1.24)	1 (0.65)	2 (1.47)	13
<i>DXS143</i> (dic56)	<i>BclI</i>	+	2	2	1	2	14
<i>DXS9</i> (RC8)	<i>TaqI</i>	—	2 (1.05)	2 (1.20)	1 (0.60)	2 (1.02)	15
<i>DXS43</i> (pD2)	<i>PvuII</i>	—	2 (0.85)	2 (0.80)	1 (0.38)	2 (0.78)	16
<i>DXS41</i> (p99-6)	<i>PstI</i>	—	2 (0.96)	2 (1.10)	1 (0.49)	2 (1.12)	16
<i>ZFX</i> (pPB, pMF-1)	<i>EcoRI</i>	—	2 (1.06)	2 (1.19)	1 (0.60)	2 (1.05)	17
<i>DXS164</i> (pERT87-1)	<i>PvuII</i>	—	2 (0.23)	2 (0.21)	1 (0.09)	2 (0.19)	18
<i>DXS84</i> (754)	<i>PvuII</i>	—	2 (0.30)	2 (0.28)	1 (0.12)	2 (0.28)	19
<i>OTC</i> (cDNA probe)	<i>PvuII</i>	—	1 (0.12)	2 (0.27)	1 (0.11)	2 (0.23)	20
<i>DXS7</i> (L1.28)	<i>TaqI</i>	—	1 (0.63)	2 (1.31)	1 (0.74)	2 (1.25)	21
Y specific region							
<i>PABY</i> (Hf0.2)	<i>SstI</i>	—	1 (0.60)	0	1 (0.75)	0	13
<i>SRY</i> (pY53.3)	<i>EcoRI</i>	—	1 (0.75)	0	1 (0.84)	0	1
<i>DYS104</i> (27a)	<i>EcoRI</i>	—	1 (0.64)	0	1 (0.73)	0	22
<i>ZFY</i> (pPB, pMF-1)	<i>EcoRI</i>	—	1 (1.23)	0	1 (1.25)	0	17

The copy number of each locus was determined by the presence of RFLP (*DXYS14*, *DXYS15*, *DXYS17*, and *DXS143*) or by the comparison of band intensity (other loci). The values in parentheses represent the ratio of the band intensity between each locus and autosomal TK gene. The loci are arranged from telomere to centromere.

she received corticosteroid therapy. At 2 years 6 months of age, a human chorionic gonadotrophin test (3000 IU/m²/dose i m for three consecutive days) was done, yielding no testosterone response (<0.5→<0.5 nmol/l). At three years of age, she exhibited persistent SLE-like symptoms and died of cachexia.

Macroscopic examination of the internal genitalia at necropsy showed that Mullerian duct derivatives (fallopian tubes, uterus, and upper portion of vagina) were normally developed and Wolffian duct derivatives were absent. Streak gonads were observed in the place of ovaries. Other organs were normal. Microscopic examination of the gonads showed ovarian stroma and gonadoblastoma (fig 1). Testicular development and ovarian germ cells were absent. In the extragonadal organs, severe necrotising vasculitis characteristic of polyarteritis nodosa was observed.

Methods

CYTOGENETIC STUDIES

Chromosome analysis was performed on 50 peripheral blood lymphocytes of the patient, her older sister, and parents using G banding.^{5,6} In the patient, high resolution G banding was also performed with ethidium bromide.⁷

SOUTHERN BLOT ANALYSIS

Genomic DNA was extracted from blood cells of all the family members. Southern transfer, probe hybridisation, and autoradiography were carried out by the standard methods.⁸ The restriction enzymes and probes used are

shown in the table. The copy number of each locus was determined by comparison of band intensity or by the presence of restriction fragment length polymorphisms. Band intensity was measured by a laser densitometer (Ultrascan, LKB), using the bands for the autosomal gene TK (probe kindly provided by Dr Y-F Lau) and Xq gene F8C²³ as intensity controls. For paternity testing, a minisatellite probe Msl²⁴ was hybridised with *AluI* digested DNA.

MUTATIONAL ANALYSIS OF SRY

SRY of this patient was subjected to both single strand conformational polymorphism (SSCP) analysis^{25,26} and DNA sequencing (fig 2). For SSCP analysis, polymerase chain reaction (PCR) amplifications were performed with primers XES10 and XES11 flanking the SRY open reading frame,¹ generating a 778 bp fragment. Amplifications were performed with 0.5 to 1.0 µg genomic DNA under standard conditions⁸ in a reaction volume of 50 µl. After an initial incubation of two minutes at 94°C, reactions were cycled for 80 seconds at 94°C, 1.5 minutes at 60°C, and 2.5 minutes at 71°C for 32 cycles. Primer sequences were: XES10 5'-GAGCTCGAGAATTCG-GTGTGTGAGGGCGGAGAAATGC-3' and XES11 5'-GAGCTCGAGAATTCGTAGC-CAATGTTACCCGATTGTC-3'. Two-fifths of the amplified DNA was fractionated on a 0.6% agarose gel. The amplified fragment was excised from the gel and melted and 1 µl of this DNA was reamplified with primers XES7 and XES2, generating a 609 bp fragment. PCR

Figure 2 A schematic map of SRY. The boxed region represents the open reading frame of SRY extending from nucleotide positions 354–1022 of the genomic clone of pY53.3. The positions of oligonucleotide PCR primers are indicated by horizontal arrows. The positions of the restriction endonuclease sites *DdeI* (D), *HpaII* (Hp), *XbaI* (X), *HinfI* (H), *TaqI* (T), and *PstI* (P) are marked between the primers XES7 and XES2. The positions of these sites are: D, 536 and 672; Hp, 495; X, 648; H, 654; T, 798; and P, 874.

Figure 3 The X and Yp+ chromosomes of the patient by high resolution G banding.

Figure 4 Southern blot analysis (P=patient, S=sister, F=father, M=mother). (A) *TaqI* digests probed with 29C1. The paternal *DXYS14* locus has not been inherited by the patient, though the maternal locus is present. (B) *SstI* digests probed with Hf0.2. The patient has both the X specific 4.5 kb band and the Y specific 3.2 kb band, with a band intensity ratio approximating 2:1. (C) *EcoRI* digests probed with pY53.3. The patient is positive for *SRY*. (D) *BclI* digests probed with dic56. RFLP is shown for the patient as well as her sister and mother, showing the presence of two copies of this locus. (E)–(I) *PvuII* digests probed with pERT87-1, 754, and probes for OTC, TK, and F8C, respectively (same filter). The patient has two copies of *DXS164*, *DXS84* and TK, and one copy of OTC and F8C.

conditions were as described previously.² Primer sequences were: XES7 5'-CCCGAATT-CGACAAATGCAATCATATGCTTCTGC-3' and XES2 5'-CTGTACCGGTCCCGTTG-CTGCGGTG-3'. One microlitre of the PCR product was digested with *DdeI*, and double digested with *HinfI* and *TaqI*, and *HpaII* and *PstI* in a 10 µl volume. SSCP analysis was as described previously.²

For DNA sequencing, two-fifths of the XES10/XES11 DNA amplification was digested with *EcoRI* to cleave the 5' ends of the primers. Digested DNA was fractionated on a 0.6% agarose gel. The DNA fragments were excised from the gel and ligated into the *EcoRI* site of pUC19. Ligated plasmids were transformed into *E. coli* DH5α. DNA from a single colony was purified by CsCl gradient centrifugation. Purified DNA was sequenced as double stranded DNA by the dideoxy chain termination method²⁷ on one strand using synthetic oligonucleotide primers and Sequenase (USB).

Results

CYTOGENETIC STUDIES

The patient's karyotype was 46,X,Yp+ in all of the 50 cells examined. On the elongated Yp, four extra dark bands were visible by high resolution G banding, the most distal band being the largest (fig 3). The karyotypes of the older sister and the parents were normal.

SOUTHERN BLOT ANALYSIS

Representative results are shown in fig 4 and summarised in the table. In the patient, only a single copy was detected for *DXYS14* and *DXYS15* in the distal part of the pseudoautosomal region (PAR), whereas three copies were found for *DXYS17* and *MIC2* in the proximal part of the PAR. The X specific loci from *PABX* to *DXS84* were present in two copies. The Y specific loci were present in a single copy as expected. Paternity was confirmed by the minisatellite analysis.

MUTATIONAL ANALYSIS OF SRY

In the SSCP analysis, none of the three different restriction enzyme digests gave an abnormal banding pattern as compared with normal male controls (fig 5). The DNA sequence was also completely normal (data not shown).

Discussion

Our results suggest that the paternal distal Xp segment (Xp21-p22.3) was translocated to Yp11.3 and inverted to form the Yp+ chromosome (Xp21→Xp22.3::Yp11.3→Yqter) (fig 6). Although the Y chromosome is missing the distal part of the PAR, no deletion of the Y specific region was detected nor was any mutation found in *SRY*. This strongly indicates that the impaired testis formation and the resultant female development of our patient occurred in the presence of *SRY*.

Figure 5 SSCP analysis of SRY. Shown is DNA from the patient described here, normal male DNA, and DNA from two XY females with apparently normal SRY genes. Amplified DNA was digested with HpaII and PstI.

This patient is similar to four non-mosaic sex reversed patients with a 46,Y,Xp+ karyotype. In spite of the presence of a morphologically normal Y chromosome, two sibs with 46,Y,dup(X)(p21→pter)²⁸ and two patients with 46,Y,dup(X)(p21.2→p22.2) and 46,Y,dup(X)(p21.2→p22.3) respectively²⁹ exhibited a female phenotype. Furthermore, the two sibs were examined for gonadal structure and confirmed to lack testis formation. In contrast to the four patients, all other reported non-mosaic patients with a partial X chromosome duplication between Xp21.2 and Xqter showed male sex development in the presence of a normal Y chromosome.^{30–36}

The four sex reversed patients and our patient have a similar duplication of an active X specific segment encompassing distal Xp21

and proximal Xp22 (the translocated Xp segment in our patient lacks the inactivation centre³⁷ and the duplicated X chromosome segments in karyotypically male patients have been reported to escape inactivation³⁶). Thus, it appears logical to assume that the same mechanism inhibiting testis formation is operating in the four patients with dup(Xp) and in our patient with Yp+. Although latent mosaicism in the gonad or a position effect on SRY might be possible in our patient, there is no evidence for either mechanism (the associated alteration of PAR is unlikely to affect sexual phenotype, since neither monosomy nor trisomy of the PAR influences testis formation^{38–39}). Similarly, although it might be possible in the four patients with dup(Xp) that latent mosaicism or a mutation of SRY existed, or that the breakage of Xp caused a gene disruption which acted as a dominant inhibitor for testis formation, such a mechanism also remains speculative.

If a causal relationship exists between two active doses of the Xp distal region and impaired testis formation, this implies that a gene or genes subject to X inactivation, involved in testis formation, exist in this region and two active copies of the gene(s) hinder the testis determination or differentiation process. Under this hypothesis, patients with only one active copy of the gene(s), for example, 47,XXY and 48,XXXY, masculinise like normal 46,XY males, whereas patients with two active copies of the gene(s), for example, 46,Y,dup(Xp) and 46,X,Yp+, result in sex reversal. Because no evidence for global developmental disruption was found in our patient, it appears that this gene(s) functions mainly, if not exclusively, in the gonad. Furthermore, it is possible that some SRY positive XY females may be explained by cryptic duplications of the gene(s) proposed here. It is also possible that other types of alteration in the gene(s) would cause sex reversal. Although Bernstein *et al*²⁸ ascribed defective testis formation of two sibs with dup(Xp) to absent H-Y antigen, and Scherer *et al*²⁹ regarded two copies of ZFX as the cause of sex reversal in two patients with dup(Xp), both hypotheses are untenable at present for the following reasons. (1) It has been shown that H-Y antigen is not required for testis determination.^{40–41} (2) ZFX has been shown to escape inactivation,⁴² so that if two copies of ZFX result in sex reversal, Klinefelter patients should develop as females.

In the present case, polyarteritis nodosa (autoimmune inflammatory disease) and IgA deficiency were observed. Interestingly, the association between sex chromosome aberrations and immune related diseases has been described previously.^{43–45} However, it is uncertain at this time whether the immune related complications of our patient were directly related to the Yp+ chromosome.

The authors would like to thank Dr T Ojima for providing important clinical data.

1 Sinclair AH, Berta P, Palmer MS, *et al*. A gene from the human sex-determining region encodes a protein with

Figure 6 A schematic representation of the generation of the Yp+ chromosome. Striped, stippled, and white areas depict X specific, Y specific, and pseudoautosomal regions, respectively. Chromosomal breakage occurred in the X and Y pseudoautosomal regions between DXYS15 and DXYS17 and in the X specific region between DXS84 and OTC. The Xp fragment (Xp21–Xp22.3) was translocated to Yp and inverted to form the Yp+ chromosome. Note that the most distal dark band of the Yp+ chromosome in fig 2 corresponds well to Xp21.

- homology to a conserved DNA-binding motif. *Nature* 1990;346:240-4.
- 2 Berta P, Hawkins JR, Sinclair AH, *et al*. Genetic evidence equating SRY and the testis-determining factor. *Nature* 1990;348:448-50.
 - 3 Jager RJ, Anvert M, Hall K, Scherer G. A human XY female with a frame shift mutation in the candidate testis-determining gene SRY. *Nature* 1990;348:452-4.
 - 4 Koopman P, Gubbay J, Vivian N, *et al*. Male development of chromosomally female mice transgenic for Sry. *Nature* 1991;351:117-21.
 - 5 Seabright M. A rapid banding technique for human chromosomes. *Lancet* 1971;ii:971-2.
 - 6 Seabright M. Improvement of trypsin method for banding chromosomes. *Lancet* 1973;i:1249-50.
 - 7 Ikeuchi T. Inhibitory effect of ethidium bromide on mitotic chromosome condensation and its application to high-resolution chromosome banding. *Cytogenet Cell Genet* 1984;38:56-61.
 - 8 Sambrook J, Fritsch EF, Maniatis T. *Molecular cloning: a laboratory manual*. 2nd ed. New York: Cold Spring Harbor Laboratory Press, 1989.
 - 9 Cooke HJ, Brown WRA, Rappold GA. Hypervariable telomeric sequences from the human sex chromosomes are pseudoautosomal. *Nature* 1985;317:687-92.
 - 10 Simmler M-C, Rouyer F, Vergnaud G, *et al*. Pseudoautosomal DNA sequences in the pairing region of the human sex chromosomes. *Nature* 1985;317:692-7.
 - 11 Rouyer F, Simmler M-C, Johnsson C, Vergnaud G, Cooke HJ, Weissenbach J. A gradient of sex linkage in the pseudoautosomal region of the human sex chromosomes. *Nature* 1986;319:291-5.
 - 12 Goodfellow PJ, Darling SM, Thomas NS, Goodfellow PN. A pseudoautosomal gene in man. *Science* 1986;234:740-3.
 - 13 Ellis NA, Goodfellow PJ, Pym B, *et al*. The pseudoautosomal boundary in man is defined by an Alu repeat sequence inserted on the Y chromosome. *Nature* 1989;337:81-4.
 - 14 Middlesworth W, Bertelson C, Kunkel LM. An RFLP detecting single copy X-chromosome fragment, dic56, from Xp22-Xpter. *Nucleic Acids Res* 1985;13:5723.
 - 15 Murray JM, Davies KE, Harper PS, Meredith L, Mueller CP, Williamson R. Linkage relationship of a cloned DNA sequence on the short arm of the X chromosome to Duchenne muscular dystrophy. *Nature* 1982;300:69-71.
 - 16 Aldridge J, Kunkel LM, Bruns G, *et al*. A strategy to reveal high frequency RFLPs along the human X chromosome. *Am J Hum Genet* 1984;36:546-64.
 - 17 Palmer MS, Berta P, Sinclair AH, Pym B, Goodfellow PN. Comparison of human ZFY and ZFX transcripts. *Proc Natl Acad Sci USA* 1990;87:1681-5.
 - 18 Monaco AP, Bertelson CJ, Middlesworth W, *et al*. Detection of deletions spanning the Duchenne muscular dystrophy locus using a tightly linked DNA segment. *Nature* 1985;316:842-5.
 - 19 Hofker MH, Wapenaar MC, Goor N, Bakker E, van Ommen G-JB, Pearson PL. Isolation of probes detecting restriction fragment length polymorphism from X chromosome-specific libraries: potential use for diagnosis of Duchenne muscular dystrophy. *Hum Genet* 1985;70:148-56.
 - 20 Horwich AL, Fenton WA, Williams KR, *et al*. Structure and expression of a complementary DNA for the nuclear coded precursor of human mitochondrial ornithine transcarbamylase. *Science* 1984;224:1068-74.
 - 21 Davies KE, Pearson PL, Harper PS, *et al*. Linkage analysis of two cloned DNA sequences flanking the Duchenne muscular dystrophy locus on the short arm of the human X chromosome. *Nucleic Acids Res* 1983;11:2303-12.
 - 22 Pritchard CA, Goodfellow PJ, Goodfellow PN. Isolation of a sequence which maps close to the human sex determining gene. *Nucleic Acids Res* 1987;15:6159-69.
 - 23 Gitschier J, Wood W, Goralka T, *et al*. Characterization of the human factor VIII gene. *Nature* 1984;312:326-30.
 - 24 Wong Z, Wilson V, Patel I, Povey S, Jeffreys AJ. Characterization of a panel of highly variable minisatellites cloned from human DNA. *Ann Hum Genet* 1987;51:269-88.
 - 25 Orita M, Iwahana K, Kanazawa H, Hayashi K, Sekiya T. Detection of polymorphisms of human DNA by gel electrophoresis as single-strand conformation polymorphisms. *Proc Natl Acad Sci USA* 1989;86:2766-70.
 - 26 Orita M, Suzuki Y, Sekiya T, Hayashi K. Rapid and sensitive detection of point mutations and DNA polymorphisms using the polymerase chain reaction. *Genomics* 1989;5:874-9.
 - 27 Sanger F, Nicklen S, Coulson AR. DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci USA* 1977;74:5463-7.
 - 28 Bernstein R, Jenkins T, Dawson B, *et al*. Female phenotype and multiple abnormalities in sibs with a Y chromosome and partial X chromosome duplication: H-Y antigen and Xg blood group findings. *J Med Genet* 1980;17:291-300.
 - 29 Scherer G, Schempp W, Baccichetti C, *et al*. Duplication of an Xp segment that includes the ZFX locus causes sex inversion in man. *Hum Genet* 1989;81:291-4.
 - 30 Steinbach P, Horstmann W, Scholz W. Tandem duplication dup(X)(q13q22) in a male proband inherited from the mother showing mosaicism of X inactivation. *Hum Genet* 1980;54:309-13.
 - 31 Nielson KB, Langkjer F. Inherited partial X chromosome duplication in a mentally retarded male. *J Med Genet* 1982;19:222-3.
 - 32 Vejerslev L, Rix M, Jespersen B. Inherited tandem duplication dup(X)(q131-q212) in a male proband. *Clin Genet* 1985;27:276-81.
 - 33 Schwartz S, Schwartz MF, Panny SR, Peterson CJ, Waters E. Inherited X-chromosome inverted tandem duplication in a male traced to a grandparental mitotic error. *Am J Hum Genet* 1986;38:741-50.
 - 34 Cremers FPM, Pfeiffer RA, van de Pol TJR, *et al*. An interstitial duplication of the X chromosome in a male allows physical fine mapping of probes from the Xq13-q22 region. *Hum Genet* 1987;77:23-7.
 - 35 Thode A, Partington MW, Yip M-Y, Chapman C, Richardson VF, Turner G. A new syndrome with mental retardation, short stature and an Xq duplication. *Am J Med Genet* 1989;30:239-50.
 - 36 Schmidt M, Du Sart D, Kalitsis P, *et al*. Duplications of the X chromosome in males: evidence that most parts of the X chromosome can be active in two copies. *Hum Genet* 1991;86:519-21.
 - 37 Brown CJ, Lafreniere RG, Powers VE, *et al*. Localization of the X inactivation centre on the human X chromosome in Xq13. *Nature* 1991;349:82-4.
 - 38 Curry CJR, Magenis RE, Brown M, *et al*. Inherited chondrodysplasia punctata due to a deletion of the terminal short arm of an X chromosome. *N Engl J Med* 1984;311:1010-5.
 - 39 Jacobs PA, Strong JA. A case of human intersexuality having a possible XXY sex determining mechanism. *Nature* 1959;183:302-3.
 - 40 McLaren A, Simpson E, Tomonari K, Chandler P, Hogg H. Male sexual differentiation in mice lacking H-Y antigen. *Nature* 1984;312:552-5.
 - 41 Simpson E, Chandler P, Goulmy E, Distèche CM, Ferguson-Smith MA, Page DC. Separation of the genetic loci for H-Y antigen and for testis determination on human Y chromosome. *Nature* 1987;326:876-8.
 - 42 Schneider-Gadicke A, Beer-Romero P, Brown LG, Nussbaum R, Page DC. ZFX has a gene structure similar to ZFY, the putative human sex determinant, and escapes X inactivation. *Cell* 1989;57:1247-58.
 - 43 Stern R, Fishman J, Brusman H, Kunkel HG. Systemic lupus erythematosus associated with Klinefelter's syndrome. *Arth Rheum* 1977;20:18-22.
 - 44 Lenoble L, Kaplan G. Systemic lupus erythematosus in a woman with primary 47XXX hypogonadism. *Rev Med Intern* 1987;8:430-2.
 - 45 Donti E, Nicoletti I, Venti G, *et al*. X-ring Turner's syndrome with combined immunodeficiency and selective gonadotropin defect. *J Endocrinol Invest* 1989;12:257-63.