LETTERS TO THE EDITOR

X linked α thalassaemia/mental retardation (ATR-X) syndrome

We read with interest the series of reports in the journal on the ATR-X syndrome.¹⁻⁴ The facial features are similar to affected subjects with another X linked MR syndrome in a Canadian family described by us a few years ago.⁵ Cole *et al*² suggested that our family may have the ATR-X syndrome. We studied the three living affected males in the family again. All showed normal haematological indices, no detectable haemoglobin H on electrophoresis, and no cells containing haemoglobin H inclusions using 1% brilliant cresyl blue preparations of peripheral blood smears.

In addition, our patients appear to be less severely retarded than the British families described. The possibilities are that these two disorders may (1) be the result of different mutations, (2) be allelic, or (3) represent an X chromosome contiguous gene syndrome with different sized deletions. In respect to the last hypothesis, high resolution chromosome analysis in our laboratories (Winnipeg and Calgary) and in Houston (courtesy of David H Ledbetter) showed no deletions. Identification of the gene locus and molecular studies will ultimately resolve the issue.

ALBERT E CHUDLEY
Section of Clinical Genetics,
Children's Hospital,
840 Sherbrook Street,
Winnipeg, Manitoba R3A 1S1, Canada.

R BRIAN LOWRY
Division of Medical Genetics,
Alberta Children's Hospital,
1820 Richmond Road SW,
Calgary, Alberta T2T 5C7, Canada.

- Gibbons RJ, Wilkie AOM, Weatherall DJ, Higgs DR. A newly defined X linked mental retardation syndrome associated with α thalassaemia. *J Med Genet* 1991;28:729-33.
- Cole TRP, May A, Hughes HE. α thalassaemia/mental retardation syndrome (non-deletion type): report of a family supporting X linked inheritance. *J Med Genet* 1991;28:734-7.
- Wilkie AOM, Gibbons RJ, Higgs DR, Pembrey ME. X linked α thalassaemia/mental retardation: spectrum of clinical features in three related males. *J Med Genet* 1991;28:738-41.
- Donnai D, Clayton-Smith J, Gibbons RJ, Higgs DR. The non-deletion α thalassaemia/mental retardation syndrome: further support for X linkage. *J Med Genet* 1991;28:742-5.
- Chudley AE, Lowry RB, Hoar DI. Mental retardation, distinct facial changes, short stature, obesity, and hypogonadism: a new X linked mental retardation syndrome. *Am J Med Genet* 1988;31:741-51.

49,XXXXY syndrome: behavioural and developmental profiles

I read with great interest the recent paper by Lomelino and Reiss,¹ reporting behavioural, psychological, and cognitive profiles of two cases of 49,XXXXY and one of 48,XXXXY/49,XXXXY mosaicism.

In 1964² I reported two cases of severely mentally handicapped patients with XXXXY syndrome, who suffered from similar developmental profiles. Case 1 improved very little in cognition, behaviour, and language

development since 1964 and died from bronchopneumonia at the age of 23 years.

Case 2 is now 44 years of age and intellectually he has improved slightly since 1964. Assessed on the Merrill Palmer Scale, his MA is 5.11 years but he still cannot read or write. His verbal comprehension and expression has shown some improvement. Results of testing on the Reynell Development Language Scales were: raw scores 42, equivalent age 3.03 years. The main characteristics which make him difficult to understand are the very low volume of his voice (he mostly speaks only in a whisper) and his shyness.

However, there are increasing problems with his behaviour. He suffers from temper tantrums, kleptomania, and self-injury, particularly to his hands. He constantly picks a chronic hypostatic ulcer³ on his leg. He is often untruthful and mischievous. The patient has been caught setting fire on many occasions and therefore requires constant supervision.

Sometimes he becomes verbally abusive, swearing and threatening violence, and will resort to throwing things.

Physically he is prone to falls and sustained a fracture of the neck of the right femur and fracture of the left patella, which is adding to his frustrations.

J JANCAR
Mental Handicap Services,
Stoke Park Hospital,
Stapleton,
Bristol BS16 1QU.

- Lomelino CA, Reiss AL. 49,XXXXY syndrome: behavioural and developmental profiles. *J Med Genet* 1991;28:609-12.
- Jancar J. Mentally defective males with XXXXY chromosomes. In: *Proc Int Copenhagen Conf on SS Ment Retard*. Copenhagen: Øster, 1964; 179-95.
- Jancar J. Hypostatic ulceration and male sex chromosomal anomalies. *BMJ* 1978;2:434.

BOOK REVIEWS

All titles reviewed here are available from the BMJ Bookshop, PO Box 295, London WC1H 9TE. Prices include postage in the UK and for members of the British Forces Overseas, but overseas customers should add 15% to the value of the order for postage and packing. Payment can be made by cheque in sterling drawn on a UK bank, or by credit card (Mastercard, Visa or American Express) stating card number, expiry date, and full name.

The Triumph of the Embryo. Lewis Wolpert. (£14.95.) Oxford: Oxford University Press. 1991.

This book is intended as an introduction to developmental biology for non-specialist readers. The majority of the book is devoted to the mechanisms of embryonic development, but cancer, ageing, and evolution are also covered from a developmental perspective. The reader is taken through the stages of embryonic development by careful discussion of important experiments and well chosen analogies, for example, gastrulation and origami. Professor Wolpert's belief that embryonic development can be described by a series of relatively simple mechanisms, and his sheer enthusiasm for the subject, provide the basis for a clear and comprehensible

account of complex processes, such as limb formation and central nervous system development.

This is an excellent book. It is informative and readable and fulfils its intended purpose (triumphantly). I would recommend it to lay readers wanting a clear and comprehensible account of embryonic development and to any health professional who wanted a gentle introduction to the subject.

E R MAHER

Antenatal Diagnosis of Fetal Abnormalities. Ed J O Drife, D Donnai. (Pp 400; £60.00, DM190.) Berlin: Springer-Verlag. 1991.

An academic paediatrician who recently reviewed advances in genetics commented that the subject had become much too important to be left to geneticists. His realisation only came lately; thus obstetricians should congratulate themselves on being farsighted enough to have established (with perhaps more than a little prompting from a few pioneering geneticists) a multidisciplinary subspecialty which, paradoxically for a subspecialty, has grown to touch upon the lives of nearly everyone in the general population. This, of course, means that workers in the field carry a very heavy burden of responsibility, so it is timely that the Royal College of Obstetricians and Gynaecologists should, in its 23rd study group meeting, examine the state of antenatal diagnosis.

This book contains papers presented at the meeting and edited transcripts of the resulting discussions. Its title is brief but do not be misled into anticipating that its contents will merely be another account of the latest clinical and laboratory techniques, although there are certainly many expert and critical expositions of these. One only has to run through the list of the meeting's participants, noting a philosopher, an economist, a psychologist, a civil servant, and several epidemiologists, to realise that a wide brief is being addressed. The broad sweep of the contribution is reflected in a list of 19 conclusions and recommendations at the end of the book. As well as statements on clinical procedures and laboratory tests, such as ultrasound examination, fetal blood sampling, and maternal serum screening, there are more general statements supporting the organisation of regional genetic centres and fetal medicine centres, calling for improved professional and lay education in genetics and recommending that ethicists should be involved in the introduction of new techniques. Two other recommendations are, firstly, each district general hospital should have a consultant obstetrician with a special interest in and responsibility for the various aspects of prenatal diagnosis and, secondly, that appropriate support, obstetric follow up, and genetic counselling should be available to women who undergo termination of pregnancy for fetal abnormality. I thought there might be a statement on the relative merits of amniocentesis and chorionic villus sampling, given that there is discussion of this at several points in the text, but perhaps the study group did not wish to prejudge the then unpublished MRC trial comparing these procedures.

The book's 25 chapters, which embrace almost the whole subject, are arranged in seven sections and comprise reviews in the