Comparison of adult height between patients with XX and XY gonadal dysgenesis: support for a Y specific growth gene(s)

Tsutomu Ogata, Nobutake Matsuo

Abstract

Adult height was compared between published cases of patients with XX gonadal dysgenesis (XXGD) and those with XY gonadal dysgenesis (XYGD). The mean adult height of XYGD patients (171.0 cm (SD 7.8), n=27) was significantly greater than that of XXGD patients (164.4 cm (7.7), n=27) ($p<0.01$). This finding supports the existence of a Y specific growth gene(s) which promotes statural growth independently of the effects of gonadal sex steroids.

Pure XX gonadal dysgenesis (XXGD) and XY gonadal dysgenesis (XYGD) are disorders of sexual development characterised by a rudimentary streak gonad and an apparently normal karyotype.¹ Consequently, affected subjects of both sexes have gonadal steroid deficiency irrespective of sex chromosome complement. Thus, it is believed that phenotypic differences between XXGD and XYGD are directly caused by the difference in sex chromosome complement, independently of the effects of gonadal sex steroids. On the basis of this notion, we compared adult heights between patients with XXGD and those with XYGD.

Methods

This investigation was based on height data of published cases of patients with XXGD and those with XYGD. The selection criteria used were: (1) apparently normal karyotype, (2) morphological confirmation of streak gonad either in the patient or in her affected family members, (3) lack of apparent Turner stigmata, (4) recorded height obtained between 20 and 50 years of age, (5) no description of sex steroid therapy before 20 years of age, and (6) absence of other associated disorders that may affect statural growth.

The adult height (mean (SD)) was compared before and after excluding non-Caucasian patients to allow for the influence of racial height variation. Statistical significance was determined by the two tailed *t* test. In addition, the age and the publication year were compared between XXGD and XYGD patients using Wilcoxon's rank-sum test, to examine the influence of secular height change (about 1 to 2 cm per decade).²⁻⁴

Results

Twenty-seven patients were identified for both XXGD⁵⁻¹⁸ and XYGD.^{7,8,15,19-34} The mean adult height was 164.4 cm (SD 7.7) (range 148 to 180 cm) for the patients with XXGD and 171.0 cm (SD 7.8) (range 150 to 183 cm) for those with XYGD. This adult height difference of 6.6 cm was statistically significant ($p<0.01$). Between the two groups, the distribution of ages was comparable (XXGD, mean 25.9 years, range 20 to 37; XYGD, mean 26.2 years, range 20 to 38) as was that of publication years (XXGD, mean 1968, range 1961 to 1982; XYGD, mean 1973, range 1964 to 1984).

Twenty-two patients with XXGD and 24 patients with XYGD appeared to be Caucasians of various nationalities; the remaining patients were two American blacks,^{11,31} five South Americans,^{17,18,30} and one African.²⁷ The mean adult height of the 22 XXGD patients was 164.3 cm (SD 7.7) (range 148 to 175 cm) and that of the 24 XYGD patients was 172.0 cm (SD 7.0) (range 156 to 183 cm). This height difference of 7.7 cm was also significant ($p<0.01$). Between the two subgroups, the distribution of ages was comparable (XXGD, mean 26.5 years, range 20 to 37; XYGD, mean 26.4, range 20 to 38) but that of publication years was statistically different (XXGD, mean 1967, range 1961 to 1971; XYGD, mean 1972, range 1964 to 1984) ($p<0.05$), implying a small influence of secular height change (about 0.5 to 1.0 cm) on the adult height difference.

Discussion

The present study showed a significant difference in the mean adult height between XXGD and XYGD patients. This adult height difference is considered to be a direct consequence of a growth promoting effect of the Y chromosome. This notion is supported by the fact that patients with 47,XYY are taller than normal 46,XY males.³⁵ In contrast, the possibility that the X chromosome has a growth suppressing effect is excluded by the fact that patients with 45,X are invariably short³⁶ and those with 47,XXX tend to be tall.³⁷

If the Y chromosome is responsible for the adult height difference between XXGD and XYGD patients, it is expected that a growth gene(s) unrelated to gonadal sex steroids exists in the Y specific region. In this context, non-mosaic adult patients with 46,X,del(Yq) are informative: in spite of a similar male sex development, those with apparently large Yq

Human Molecular Genetics Laboratory, Imperial Cancer Research Fund, PO Box 123, Lincoln's Inn Fields, London WC2A 3PX.
T Ogata

Department of Paediatrics, Keio University School of Medicine, Tokyo 160, Japan.
N Matsuo

Correspondence to Dr Ogata.

Received 1 November 1991.
Revised version accepted 24 January 1992.

deletions had short stature³⁸⁻⁴⁰ and those with apparently small Yq deletions did not have short stature.⁴¹⁻⁴⁵ This finding is compatible with the presence of a Y specific growth gene(s) which promotes statural growth independently of the effects of gonadal sex steroids, although other mechanisms may also contribute to the growth failure associated with large chromosomal deletions. In agreement with this view, recent molecular analysis in patients with Y chromosome aberrations has also argued for the presence of the growth gene(s) in the proximal part of Yq.⁴⁶ Although another growth gene(s) has been suggested on the tip of Yp distal to the testis determining gene,⁴⁷ genotype-phenotype correlations in patients with sex chromosome abnormalities have indicated that the growth gene(s) on the tip of Yp is not Y specific but pseudoautosomal.^{48,49}

Our results suggest that the putative Y specific growth gene(s) may exert an influence of about 7 cm on adult height. This value is in close agreement with the adult height difference between normal males (174.7 cm (SD 6.7))⁵⁰ and SRY positive XX males (166.1 cm (SD 3.3), n = 8, 20 to 50 years of age)⁵¹ in the United Kingdom. Since the SRY positive XX males have normal external genitalia indicative of sufficient testosterone secretion⁵¹ and are expected to have two copies of the pseudoautosomal growth gene(s), the height decrease of the XX males can be simply explained as a loss of the Y specific growth gene(s). In addition, it is known that patients with testicular feminisation syndrome (TFS) are 5 to 10 cm taller than normal females, despite having similar sex steroid hormone status.⁵²⁻⁵⁴ This height increase of TFS patients is also explicable by the effect of the Y specific growth gene(s).

The adult height difference between XXGD and XYGD patients was smaller than that between normal males and females (about 13 cm).^{2,50,55} This indicates that sex dimorphism in gonadal steroids also contributes to the sex difference in the mean adult height of normal subjects. Since the mean adult height of XXGD patients is similar to the adult female standards and that of XYGD patients is shorter than the adult male standards,^{2,50,55} it appears that ovarian oestrogens play a relatively minor role in statural growth, whereas testicular androgens have the potential to increase the adult height. In favour of this notion, it has been reported that androgens have a more beneficial effect on the production of somatomedin-C (a potent stimulus for linear growth) than oestrogens.⁵⁶ Furthermore, this notion can explain the adult height differences between XX males and normal females and between TFS patients and normal males.

It should be pointed out, however, that possible variations in several factors influencing the adult height, such as secular trend, racial group, parental height, and socioeconomic status,^{3,4,57,58} are not eliminated between the XXGD and XYGD patients analysed. The distributions of ages and publication years imply a possible small secular height change between the two groups of patients. In addition, although the exclusion of non-Caucasian

patients allowed the height comparison in a more homogeneous population, intra-Caucasian height variation still exists.^{2,50,55} Furthermore, no parental height was available in the published reports, nor was there any indication of socioeconomic status. Thus, it is uncertain at this time whether the adult height difference between XXGD and XYGD patients (about 7 cm) is totally attributable to the Y specific growth gene(s). Further accumulation of patients will be required for the more precise estimation of the quantitative effect of the Y specific growth gene(s).

The authors are grateful to Dr Peter Goodfellow, Laboratory of Human Molecular Genetics, Imperial Cancer Research Fund, for his invaluable advice on this work.

- Grumbach MM, Conte FA. Disorders of sexual differentiation. In: Wilson JD, Foster DW, eds. *Williams textbook of endocrinology*. 7th ed. Philadelphia: Saunders, 1985:312-401.
- Prader A, Largo RH, Molinari L, Issler C. Physical growth of Swiss children from birth to 20 years of age. *Helv Paediatr Acta* 1989;suppl 52:1-125.
- Bielicki T. Physical growth as a measure of the economic well-being of populations: the twentieth century. In: Falkner F, Tanner JM, eds. *Human growth*. 2nd ed. New York: Plenum, 1985:283-305.
- van Wieringen JC. Secular growth changes. In: Falkner F, Tanner JM, eds. *Human growth*. 2nd ed. New York: Plenum, 1985:307-31.
- Jacobs PA, Harnden DG, Buckton KE, et al. Cytogenetic studies in primary amenorrhea. *Lancet* 1961;i:1183-8.
- Ferguson-Smith MA, Alexander DS, Bowen P, et al. Clinical and cytogenetic studies in female gonadal dysgenesis and their bearing on the cause of Turner's syndrome. *Cytogenetics* 1964;3:355-83.
- Kinch RAH, Plunkett ER, Smout MS, Carr DH. Primary ovarian failure: a clinicopathological and cytogenetic study. *Am J Obstet Gynecol* 1965;91:630-44.
- Boczkowski K, Teter J. Clinical, histological and cytogenetic observations in pure gonadal dysgenesis. *Acta Endocrinol (Copenh)* 1966;51:497-510.
- Serment H, Lafargue P, Piana L, Bossi G, Felce A, Hartung M. Essai de classification des insuffisances ovariennes primitives. *Rev Fr Endocrinol Clin* 1966;7:461-86.
- Greenblatt RB, Byrd JR, McDonough PG, Mahesh VB. The spectrum of gonadal dysgenesis. *Am J Obstet Gynecol* 1967;98:151-72.
- Board JA, Estep HL, Young RB, Borland DS. Gonadal dysgenesis in a eunuchoid XX female. *Virg Med Month* 1967;94:91-5.
- Boczkowski K. The syndrome of pure gonadal dysgenesis. *Med Gynecol Soc* 1968;3:96-104.
- Judd HL, Scully RE, Atkins L, Neer RM, Kliman B. Pure gonadal dysgenesis with progressive hirsutism. *N Engl J Med* 1970;282:881-5.
- Boczkowski K. Pure gonadal dysgenesis and ovarian dysplasia in sisters. *Am J Obstet Gynecol* 1970;106:626-8.
- Boczkowski K. Further observations on the syndrome of pure gonadal dysgenesis. *Am J Obstet Gynecol* 1970;106:1177-83.
- Simpson JL, Christakos AC, Horwith M, Silverman FS. Gonadal dysgenesis in individuals with apparently normal chromosomal complements: tabulation of cases and compilation of genetic data. *Birth Defects* 1971;7:215-28.
- Nazareth HRS, Farah LMS, Cunha AJB, Viera FJPB. Pure gonadal dysgenesis (type XX): report on a family with four affected sibs. *Hum Genet* 1977;37:117-20.
- Youlton R, Michelsen H, Be C, Cruz-Coke R. Pure XX gonadal dysgenesis in identical twins. *Clin Genet* 1982;21:262-5.
- Court-Brown WM, Harnden DG, Jacobs PA, Maclean N, Mantle DJ. Abnormalities of the sex chromosome complement in man. *Medical Research Council Special Report Series* 1964; No 305:220.
- Cohen MM, Shaw MW. Two XY siblings with gonadal dysgenesis and a female phenotype. *N Engl J Med* 1965;272:1083-8.
- Dewhurst CJD. The XY female. *J Obstet Gynecol Br Cwlth* 1967;74:353-66.
- Guinet P, Putelat R, Laurent C, Tourniare A, Nivelon A, Eyraud MT. Une observation de dysgenésie gonadique avec gonocytome caryotype XY. *Ann Endocrinol (Paris)* 1967;28:375-80.
- Lieberman LL, Bresler VM, Verlinskaya DK, et al. Karyotype and clinical symptoms in gonadal dysgenesis. *Genetica* 1967;7:83-100.
- Rigo S, Martin FIR, MacDonald IA, Cowling DC. Chromosomal analysis and gonadotropin studies in primary amenorrhea. *Med J Aust* 1968;55:515-9.
- Sternberg WH, Barclay DL, Kloepfer HW. Familial XY gonadal dysgenesis. *N Engl J Med* 1968;278:695-700.

- 26 Stanesco V, Maximilian C, Florea I, Ciovirneche M. Trois soeurs avec dysgenesie gonadale pure et caryotype XY. *Ann Endocrinol (Paris)* 1968;29:449-59.
- 27 Smith M, Pepler WJ. Female phenotype associated with XY karyotype: report of a case with gonadal dysgenesis. *S Afr Med J* 1969;43:1414-5.
- 28 Boczkowski K, Teter J, Sternadel Z. Sibship occurrence of XY gonadal dysgenesis with dysgerminoma. *Am J Obstet Gynecol* 1972;113:952-5.
- 29 Nazareth HRS, Moreira-Filho CA, Cunha AJB, Viera-Filho JPB, Lengyel AMJ, Lima MC. H-Y antigen in 46,XY pure testicular dysgenesis. *Am J Med Genet* 1979;3:149-54.
- 30 Moreira-Filho CA, Toledo SPA, Bagnolli VR, Frota-Pes-soa O, Bisi H, Wajntal A. H-Y antigen in Swyer syndrome and the genesis of XY gonadal dysgenesis. *Hum Genet* 1979;53:51-6.
- 31 Phansay SA, Satterfield R, Jorgenson RJ, et al. XY gonadal dysgenesis in three siblings. *Am J Obstet Gynecol* 1980;138:133-8.
- 32 Moltz L, Schwartz U, Pickartz H, Hammerstein J, Wolf U. XY gonadal dysgenesis: aberrant testicular differentiation in the presence of H-Y antigen. *Obstet Gynecol* 1981;58:17-25.
- 33 Fisher RA, Salm R, Spencer RW. Bilateral gonadoblastoma/dysgerminoma in a 46XY individual: case report with hormonal studies. *J Clin Pathol* 1982;35:420-4.
- 34 Nagamani M, van Dinh T, Hannigan EV. XY gonadal dysgenesis with gonadoblastoma and dysgerminoma. *South Med J* 1984;77:1205-7.
- 35 Court Brown WM. Males with an XYY sex chromosome complement. *J Med Genet* 1968;5:341-59.
- 36 Brook CGD, Murset G, Zachmann M, Prader A. Growth in children with 45,XO Turner's syndrome. *Arch Dis Child* 1974;49:789-95.
- 37 Linden MG, Bender BG, Harman RJ, Mrazek DA, Robinson A. 47,XXX: what is the prognosis? *Pediatrics* 1988;82:619-30.
- 38 Langmaid H, Laurence KM. Deletion of the long arms of the Y chromosome with normal male development and intelligence. *J Med Genet* 1974;11:208-11.
- 39 Yunis E, Garcia-Conti FL, de Caballero OMT, Giraldo A. Yq deletion, aspermia, and short stature. *Hum Genet* 1977;39:117-22.
- 40 Kosztolanyi G, Trixler M. Yq deletion with short stature, abnormal male development, and schizoid character disorder. *J Med Genet* 1983;20:393-4.
- 41 Alvesalo L, de la Chapelle A. Tooth sizes in two males with deletions of the long arm of the Y-chromosome. *Ann Hum Genet* 1981;45:49-54.
- 42 Meisner LF, Inhorn SL. Normal male development with Y chromosome long arm deletion (Yq-). *J Med Genet* 1972;9:373-7.
- 43 Soudek D, Laraya P. C and Q bands in long arm of Y chromosomes; are they identical? *Hum Genet* 1976;32:339-41.
- 44 Tiepolo L, Zuffardi O. Localization of factors controlling spermatogenesis in the nonfluorescent portion of the human Y chromosome long arm. *Hum Genet* 1976;34:119-24.
- 45 Cohen G, Manuel A, Cohen M, Fagan K, Grunstein H. A deletion of heterochromatin only of the Y chromosome in an azoospermic male. *Hum Genet* 1983;64:297-300.
- 46 Bardoni B, Zuffardi O, Guioli S, et al. A deletion map of the human Yq11 region: implications for the evolution of the Y chromosome and tentative mapping of a locus involved in spermatogenesis. *Genomics* 1991;11:443-51.
- 47 Buhler EM. A synopsis of the human Y chromosome. *Hum Genet* 1980;55:145-75.
- 48 Ballabio A, Bardoni B, Carrozzo R, et al. Contiguous gene syndrome due to deletion in the distal short arm of the human X chromosome. *Proc Natl Acad Sci USA* 1989;86:10001-5.
- 49 Ogata T, Goodfellow P, Petit C, Aya M, Matsuo N. Short stature in a girl with a terminal Xp deletion distal to DXY5:5: localisation of a growth gene(s) in the pseudo-autosomal region. *J Med Genet* 1992;29:455-9.
- 50 Tanner JM, Whitehouse RH, Takaishi M. Standards from birth to maturity for height, weight, height velocity, and weight velocity: British children, 1965. *Arch Dis Child* 1966;41:454-71 (part I), 1966;41:613-35 (part II).
- 51 Ferguson-Smith MA, Cooke A, Affara NA, Boyd E, Tolmie JL. Genotype-phenotype correlations in XX males and their bearing on current theories of sex determination. *Hum Genet* 1990;84:198-202.
- 52 Varrela J, Alvesalo L, Vinkka H. Body size and shape in 46,XY females with complete testicular feminization. *Ann Hum Biol* 1984;11:291-301.
- 53 Smith DD, Marokus R, Graham JM Jr. Tentative evidence of Y-linked statural gene(s). *Clin Pediatr* 1985;24:189-92.
- 54 Zachmann M, Prader A, Sobel EH, et al. Pubertal growth in patients with androgen insensitivity: indirect evidence for the importance of estrogens in pubertal growth of girls. *J Pediatr* 1986;108:694-7.
- 55 Tanner JM, Davies PSW. Clinical longitudinal standards for height and height velocity for North American children. *J Pediatr* 1985;107:317-29.
- 56 Caufriez A, Copinschi G. Somatomedins and steroids. *Horm Res* 1986;24:185-8.
- 57 Mueller WH. The genetics of size and shape in children and adults. In: Falkner F, Tanner JM, eds. *Human growth*. 2nd ed. New York: Plenum, 1985:145-68.
- 58 Eveleth PB. Population differences in growth: environmental and genetic factors. In: Falkner F, Tanner JM, eds. *Human growth*. 2nd ed. New York: Plenum, 1985:221-39.