Usher syndrome type I associated with bronchiectasis and immotile nasal cilia in two brothers

Dominique Bonneau, Françoise Raymond, Christine Kremer, Jean-Michel Klossek, Josseline Kaplan, Françoise Patte

Abstract

Usher syndrome type I is an autosomal recessive disease characterised by congenital sensorineural deafness, involvement of the vestibular system, and progressive visual loss owing to retinitis pigmentosa. Here we report the association of this disease with bronchiectasis, chronic sinusitis, and reduced nasal mucociliary clearance in two sibs and we suggest Usher syndrome type I could be a primary ciliary disorder.

(*J Med Genet* 1993;30:253-4)

Usher syndrome (US) is an autosomal recessive disease which combines congenital sensorineural deafness and progressive visual loss owing to retinitis pigmentosa (RP).¹ It is considered to be the most frequent cause of deaf-blindness in adults and affects 3 to 6% of deaf children.² Two distinct clinical and genetic subtypes, Usher types I and II, have been described.¹ Usher syndrome type I (USI) is defined by profound congenital hearing impairment with unintelligible speech, vestibular dysfunction, and early onset of RP within the first or at the beginning of the second decade. In Usher type II, deafness is moderate with intelligible speech and RP occurs later. Here we report the association of USI with immotile cilia syndrome in two sibs and suggest this disease could be a primary ciliary disorder.

Case reports

The two brothers (II-1 and II-2, fig 1) were born in 1943 and 1947 to non-consanguineous parents without any family history. They both exhibited the same progression of the disease. Profound sensorineural deafness (120 dB loss)

was diagnosed at the age of 18 months in both of them and they have a total absence of intelligible speech.

Ophthalmic signs were identical in both of them: hemeralopia and visual field loss were first noted before 15 years of age and retinitis pigmentosa was diagnosed on concentric loss of visual field, typical bone spicule on fundal examination, and unrecordable electroretinogram.

Vestibular function was impaired as both had loss of balance; standard caloric testing showed absence of nystagmic responses in the younger patient and was uninterpretable in the older because of spontaneous nystagmus.

In addition, they both had had chronic bronchitis with productive cough from childhood and x rays showed bronchial dilatation predominantly in the lower lobes of the lungs (fig 2) together with maxillary sinusitis.

Département de
Génétique Médicale,
Centre Hospitalier
Universitaire, BP 577,
86021 Poitiers Cedex,
France.
D Bonneau
F Raymond

Service d'ORL,
Centre Hospitalier
Universitaire,
Poitiers, France.
J M Klossek
C Kremer

Service de
Pneumologie, Centre
Hospitalier
Universitaire,
Poitiers, France.
F Patte

Unité de Recherche
sur les Handicaps
Génétiques de
l'Enfant, INSERM
U12, Hôpital des
Enfants-Malades,
Paris, France.
J Kaplan

Correspondence to
Dr Bonneau.

Received 6 May 1992.
Revised version accepted
13 July 1992.

Figure 1 Pedigree of family.

Figure 2 Pulmonar radiographs of II-1 showing bilateral bronchiectasis and normal position of the heart.

Intrathoracic organs were in the normal position.

Nasal mucociliary clearance was assessed using Puchelle's method by measuring the progression of a coloured index placed on the inferior nasal turbinate³; in two consecutive tests progression of the index was less than 10 mm after 20 minutes in both patients (normal = 0.5 to 24 mm/minute).

Electron microscopy of nasal and bronchial biopsies did not show any ultrastructural abnormality of cilia; semen samples for study of spermatozoa velocity and ultrastructure were impossible to obtain.

Discussion

These two brothers exhibited typical Usher syndrome type I, bronchiectasis, chronic sinusitis, and reduced nasal mucociliary clearance. To our knowledge, such an association has not previously been reported and suggests USI could be the result of primary ciliary dyskinesia. This might account for the combined involvement of three different sensory systems since photoreceptors, auditory hair cells, and vestibular hair cells develop from ciliated progenitors.⁴

Several other reports indicate that a generalised abnormality of cilia is present in patients with Usher syndrome. (1) Sperm motility, velocity, and structure have been found to be abnormal.⁴ (2) A high proportion of abnormal axonemes has been found in retinal photoreceptor cells in one patient.⁴ (3) A decrease of outer ciliar cells has been reported at the lower part of the cochlea.⁵ Arden and Fox⁶ have found an increased incidence of abnormal ultrastructure of axonemes in nasal cilia in 11 subjects with retinitis pigmentosa; interestingly, six of their 11 patients were suffering from Usher syndrome.

We have recently proposed that a gene for USI could be located on the long arm of chromosome 14 (q32) for linkage with the DNA probe MLJ14 at locus *DS14S13* (lod

score = 3.20 at $\theta = 0$).⁷ However, further linkage analysis clearly supports the view that US type I is a genetically heterogeneous condition. Heterogeneity test HOMOG allowed us to separate our families into two groups: one linked to chromosome 14 and another one unlinked. Interestingly, eight of the 10 families linked to 14q32 originated from the same area of western France, suggesting a founder effect.⁸ It is of particular interest to note that a mouse gene controlling the position of heart and viscera has recently been mapped to a homologous region to the human 14q32 region,⁹ especially as a normal ciliary function is thought to be required during embryogenesis for normal positioning of the heart and viscera. It is therefore tempting to hypothesise that a relationship could exist between Usher syndrome type I and Kartagener syndrome, especially as cases of Kartagener syndrome have been reported either in association with retinitis pigmentosa¹⁰ or congenital deaf mutism.¹¹

1 Fishman GA, Kumar A, Joseph ME, Torok N, Anderson RJ. Usher's syndrome: ophthalmic and neuro-otologic findings suggesting genetic heterogeneity. *Arch Ophthalmol* 1983;101:1367-74.

2 Boughman JA, Vernon M, Shaver KA. Usher syndrome: definition and estimate of prevalence from two high-risk populations. *J Chron Dis* 1983;36:595-603.

3 Puchelle E, Aug F, Bertrand A. Comparison of three methods for measuring nasal mucociliary clearance in man. *Acta Otolaryngol* 1981;91:297-303.

4 Hunter DG, Fishman GA, Mehta RS, Kretzer FL. Abnormal sperm and photoreceptor axonemes in Usher's syndrome. *Arch Ophthalmol* 1986;104:383-9.

5 Shinkawa H, Nadol JB. Histopathology of the inner ear in Usher's syndrome as observed by light and electron microscopy. *Ann Otol Rhinol Laryngol* 1986;95:313-8.

6 Arden GB, Fox B. Increased incidence of abnormal nasal cilia in patients with retinitis pigmentosa. *Nature* 1979;279:534-6.

7 Kaplan J, Gerber S, Bonneau D, et al. Location of Usher type I gene on chromosome 14 q by linkage with D14S13 (MLJ14 probe). *11th International Workshop on Human Gene Mapping, London, 18-22 August, 1991*.

8 Kaplan J, Gerber S, Bonneau D, et al. A gene for Usher syndrome type I maps to chromosome 14 q. *Genomics* (in press).

9 Bruekner M, D'Eustachio P, Horwich AL. Linkage mapping of a mouse gene "iv", that controls left-right asymmetry of the heart and viscera. *Proc Natl Acad Sci USA* 1989;86:5035-8.

10 Segal P, Kikiela M, Mrzyglod S, Zeromeska-Zbierska I. Kartagener syndrome with familial eye changes. *Am J Ophthalmol* 1963;55:1043-9.

11 Lake K, Sharma OP. Kartagener syndrome and deaf mutism: an unusual association. *Chest* 1973;64:661-3.