Syndrome of the month

Albright's hereditary osteodystrophy

Louise C Wilson, Richard C Trembath

History and terminology

In 1942 Albright *et al*¹ investigated a patient with seizures, hypocalcaemia, and hyperphosphataemia consistent with a diagnosis of hypoparathyroidism. Unexpectedly, repeated injections of bovine parathyroid extract failed to alter the serum calcium or phosphate levels. They concluded the patient was resistant to parathyroid hormone (PTH) rather than deficient and coined the term pseudohypoparathyroidism (PHP). Additional features, (a short stocky build, round face, cutaneous ossification, and metacarpophalangeal abnormalities) were sufficiently distinctive for two unrelated but phenotypically similar patients with hypocalcaemia to be recalled and shown to have PTH resistance. This physical appearance is known as Albright's hereditary osteodystrophy (AHO).

A decade later, Albright *et al*² described a further patient with typical features of AHO but normal serum calcium and phosphate. Recognising the phenotypic similarities but absence of PTH resistance, he named this disorder pseudopseudohypoparathyroidism (PPHP).

Subsequent reports of families where both conditions were segregating suggested an aetiological link.

Aetiology

PTH, like TSH and several other hormones,³

exerts its effects by stimulating adenylyl cyclase to produce the intracellular second messenger cAMP. Their receptors are coupled to adenylyl cyclase by a signal transducing protein known as Gs, one of a large family of heterotrimeric GTP binding G proteins.⁴⁻⁶ Gs comprises an α , β , and γ subunit (fig 1). The α subunit appears to determine both the receptor and effector specificity.

In 1980 two groups showed that Gs α activity in erythrocyte membranes from patients with PHP was reduced to 50% of normal controls.^{7,8} Similar reductions have been found in membranes of platelets,⁹ fibroblasts,¹⁰ and renal cortex¹¹ from patients with PHP. Intriguingly, equivalent Gs α reductions have been found in membranes from patients with PPHP, despite their lack of hormone resistance.¹²

Diagnosis and classification

A diagnosis of PHP may be inferred in a patient who has hypocalcaemia, hyperphosphataemia,

Departments of
Genetics and
Medicine, University
of Leicester, Leicester
LE1 7RH, UK
L C Wilson
R C Trembath

Mothercare Unit for
Clinical Genetics and
Fetal Medicine,
Institute of Child
Health, 30 Guilford
Street, London
WC1N 1EH, UK
L C Wilson

Correspondence to
Dr Wilson, Department
of Genetics, University
of Leicester, Leicester,
LE1 7RH, UK.

Figure 1 Diagram illustrating the hormone receptor-Gs-adenylyl cyclase pathway.

Figure 2 Male infant with AHO.

Table 1 Classification of pseudohypoparathyroidism

Type	Response to PTH	Phenotype	Multiple hormone resistance	Gsα levels (% controls)	Aetiology
PHP Ia	Blunted cAMP Blunted Pi	AHO	Frequent	50	Deactivating Gsα gene mutation ? PTH receptor
PHP Ib	Blunted cAMP Blunted Pi	No AHO	No	100	? Defective adenyl cyclase
PHP Ic	Blunted cAMP Blunted Pi	AHO	Frequent	100	? Distal to adenyl cyclase
PHP II	Normal cAMP Blunted Pi	No AHO	No	100	?phospholipase C pathway ¹⁷
PPHP	Normal cAMP Normal Pi	AHO	No	50	Deactivating Gsα gene mutation

PHP, even when resistance to PTH has been established through PTH infusion testing.

Clinical features

Typical patients with AHO (figs 2 and 3) have a pleasant, affable nature, short, stocky build, round face with a low, flat nasal bridge, short neck, ectopic ossification, brachymetaphalangism, and developmental delay, with or without endocrine abnormalities, and these subjects pose little diagnostic difficulty. However, aspects of the phenotype may vary considerably from one patient to another.

Figure 3 Child with AHO showing round face and short nose with low, flattened nasal bridge.

GENERAL APPEARANCE

Fitch¹⁹ noted short stature in 39% of patients with AHO below the age of 18 years and 62% of adults. These findings and longitudinal studies^{20,21} suggest that advanced skeletal maturation is a contributing factor. There is typically a shortening of the extremities.^{22,23} Occipitofrontal circumference is generally within the normal range but macrocephaly has been described.¹⁹ Obesity is usual and often arises in early childhood.¹⁹

NEUROMUSCULAR

Mental retardation has been reported in 52/66 (70%) hypocalcaemic and 19/64 (30%) normocalcaemic patients with AHO.¹⁹ In PHP Ia, mild to moderate mental retardation occurred in 9/14 (64%) patients.²⁴ There appeared to be a strong correlation with reduced Gsα activity since no patients with PHP Ib had mental retardation in spite of equivalent serum calcium and phosphate abnormalities. While untreated hypothyroidism may contribute to the degree

and raised serum immunoactive PTH levels, providing renal function is normal. Resistance can be shown more directly by assessing the renal response to exogenous PTH.¹³⁻¹⁵ Normally an intravenous bolus of PTH produces a rapid rise in plasma and urinary cAMP with a slower rise in urinary phosphate excretion. Subjects with PHP have blunting of one or both of these responses (table 1).

It is important to note that PHP is not always associated with AHO and in such instances the aetiology is likely to be different. The classification of PHP is therefore based on the response to PTH testing and Gsα levels (table 1).¹⁶⁻¹⁸ The majority of those with PTH resistance and AHO have PHP Ia.

Persons with PPHP have, in addition to the AHO phenotype, a normal renal response to PTH but reduced levels of Gsα (table 1). Use of the term PPHP to denote any person with AHO who is normocalcaemic is unhelpful since periods of normocalcaemia are frequent in

Figure 4 Multiple, small, plaque-like foci of intradermal ossification on the dorsal aspect of the thigh in a child with AHO.

Figure 5 Plain radiograph of hand of a 4.7 year old female with AHO showing a large focus of subcutaneous ossification, brachymetaphalangism (involving all the metacarpals), phalangeal shortening, cone shaped epiphyses, and advanced bone age.

of mental retardation, it is unlikely to be the primary cause.^{18 24}

Our own observations confirm that mental retardation in AHO may range from moderately severe delay to entirely normal educational ability, but is in general mild. Of 16 adults, many have needed remedial help or special schooling but 15/16 (94%) have subsequently been able to live independently, remain in employment and, care for families of their own.

Seizures, usually grand mal, occur in AHO and may be independent of hypocalcaemia. Tetany and laryngeal stridor¹ may occur secondary to hypocalcaemia and the latter, if unrecognised, may result in unnecessary tracheostomy.

ECTOPIC CALCIFICATION

Spranger²⁵ noted cutaneous calcification in 42% of reported patients with PHP and 27% with normocalcaemic AHO, intracranial calcification occurring in 33% and 4% respectively.

Cutaneous calcification may be intradermal or subcutaneous, is frequently ossified,^{1 26-28} and has been noted within days of birth.²³ It is usually painless and often appears as multiple small superficial plaques (fig 4), easier to feel than see, with a predilection for the scalp, hands and feet, periarticular regions, abdomen, and chest wall. Larger lumps may occur, occasionally disrupting underlying structures (fig 5).

Intracranial calcification commonly involves the basal ganglia but may involve other structures including the dentate nucleus,²⁹ falx cerebri,¹ dura, cerebral cortex, and cerebellum.³⁰ Calcification may occur at

Figure 6 Short, broad thumb nails of an adult with AHO.

Figure 7 Hands of an adult with AHO showing marked shortening of the fourth metacarpal and distal phalanges with knuckle dimples in the clenched fists.

other sites including the sclera, choroid,³¹ and cardiac ventricular septum.³²

Soft tissue ossification appears to arise by a mechanism independent of serum calcium and phosphate levels since it occurs in PPHP, frequently precedes the biochemical abnormalities in PHP, and is rare in idiopathic hypoparathyroidism. Excision has been recommended for potentially troublesome or disfiguring foci^{19 28} and, unlike myositis ossificans, surgery does not appear to aggravate the condition (Wilson and Trembath, unpublished observations).³³ In contrast intracranial calcification is probably related to altered calcium/phosphate ratio.

BRACHYMETAPHALANGISM

Hand abnormalities in the PHP and PPHP forms of AHO appear indistinguishable,³⁴ involve both the phalanges and metacarpals, and may be asymmetrical. Shortening of the distal phalanx of the thumb, estimated to occur in 75% of AHO patients,³⁴ is as frequent as metacarpal shortening. This manifests as a short broad thumb nail (fig 6) which may be quan-

Figure 8 Feet of a child with AHO showing asymmetrical metatarsal shortening and shortened distal phalanges.

tified as a width to length ratio.³⁵ Metacarpal shortening most commonly involves the fourth then fifth metacarpals and least often the second,^{23,34} but any combination or indeed shortening of all the metacarpals may be observed (figs 5 and 7). When marked it may be recognised by knuckle dimples in the clenched fist (fig 7). However, subtle or generalised shortening is more reliably detected by measurement of bone lengths on x ray and metacarpal pattern profiles.³⁴ It is important to note that the characteristic hand changes evolve with age and may not be apparent until at least 4 years.

Similar abnormalities of the feet (fig 8) occur in at least 70% of persons with AHO, shortening of the fourth then third metatarsals occurring most frequently.²⁹ Phalangeal shortening also occurs but is more difficult to evaluate owing to the wide range of normal variation.

OTHER FEATURES

Eye

Cataracts and lenticular opacities have been

described in PHP probably related to the hypocalcaemia/hyperphosphataemia.²⁵ Strabismus occurred in 10% in one series.³⁶ Scleral and choroidal calcification have been observed.³¹

Hearing/olfaction

In our experience a history of glue ear requiring grommets is common in AHO perhaps because of midfacial hypoplasia. Sensorineural hearing loss has been identified in some patients with PHP Ia.³⁷ Weinstock *et al*³⁸ reported impaired olfaction in 5/5 patients with PHP Ia but 0/7 patients with PHP Ib; however, it is rarely recognised by patients.

Dental

The dental manifestations in PHP include small crowns, thin hypoplastic enamel, pulp calcifications, widened root canals, and delayed eruption of permanent teeth,^{29,39-41} and resemble those of idiopathic hypoparathyroidism.

SKELETAL AND RADIOLOGICAL FEATURES

Radiological findings are summarised in table 2. Notably, in persons with PHP Ia, evidence of osteomalacia or rickets may arise owing to low levels of 1,25 vitamin D, secondary to renal PTH resistance. Conversely, findings of generalised osteoporosis and occasionally osteitis fibrosa cystica^{29,42} suggest some preservation of the skeletal remodelling response to the raised levels of circulating PTH.^{18,42,43}

In both forms of AHO, bone age measured by the carpal bone (CB) score or the radius-ulna-short (RUS) bone score is generally advanced,^{19,21,29} the discrepancy with chronological age being greatest for the RUS score.²¹ The cone shaped, absent, and prematurely fused epiphyses in the hands and feet are thought to contribute to metacarpal and phalangeal shortening and, combined with decreased longitudinal growth, to explain the increasing prominence of these bony changes with age.

Exostoses are frequent in AHO but are not usually numerous and are generally distinguishable radiologically from multiple familial exostoses.²⁹

Endocrinological features

Resistance to hormones other than PTH occurs in PHP Ia with effects on the thyroid and gonadal axes being of clinical importance.

THYROID

Hypothyroidism secondary to TSH resistance is common in PHP Ia and has been the presenting feature in some neonates.^{44,45} In a series of 13 persons with PHP Ia,⁴⁶ 69% were found to have reduced T4, 69% reduced T3, 85% raised TSH, and 92% had an exaggerated TSH response to TRH. Thyroid function should be tested regularly.

GONADAL

Gonadal dysfunction, particularly menstrual irregularity, has been described, probably sec-

Table 2 Common radiological findings in AHO

General	Hands/feet	Limbs	Axial	Skull
Ectopic calcification/ossification	Metacarpal and phalangeal shortening	Shortened ulna Radial bowing	Caudal narrowing of lumbar interpedicular distances	Thickened calvarium with widened diploe
Advanced bone age	Cone shaped, absent, prematurely fused epiphyses	Tibial bowing Coxa vara		Hyperostosis frontalis interna
Osteoporosis		Coxa valga		Intracranial calcification
Exostoses	Metacarpal thickening and loss of diaphyseal constriction			Retained teeth
Osteomalacia/rickets				Pulp calcifications
Osteitis fibrosa cystica				

ondary to gonadotrophin resistance.^{40,47,48} In practice, investigation is probably only merited if problems arise.

GROWTH HORMONE

Growth hormone (GH) secretion and GH treatment for short stature in AHO has not been widely evaluated, possibly because short stature is also a feature of the hormone responsive form (PPHP).

Genetics

The human *Gsα* gene spanning 20 kb and 13 exons was cloned in 1988.⁴⁹ It has been localised to chromosome 20q13^{50,51} and is present in single copy per haploid genome. Heterozygous loss of function mutations of the *Gsα* gene have been found both in persons with PHP Ia and PPHP. These have predominantly been single base changes^{52–55} and small deletions^{53,54,56} with a 43 bp deletion being the largest to date.⁵⁷ So far mutations have been unique to each family and there are no obvious mutation hot spots. Furthermore in five reported families, related persons with PHP Ia and PPHP have been found to have the same *Gsα* mutation.^{52–54} Additional factors therefore appear to influence the phenotype resulting from such *Gsα* gene mutations and one of these may be the parental origin.

An excess of maternal transmissions has long been recognised in AHO and until recently X linked dominant inheritance was commonly counselled. In a recent review of reported familial cases, offspring with PHP I were found to result exclusively from maternal transmission and the few offspring with PPHP from paternal transmission.⁵⁸ We have noted a concordance of phenotype in nine affected sibs from four different families and in one family have observed a switch in phenotype from PPHP in a female with a paternally derived *Gsα* abnormality to PHP Ia in her own offspring.⁵⁵

This suggests that the *Gsα* gene may be subject to a parent of origin effect, perhaps through imprinting. Localisation of *GNAS*, the mouse homologue of *Gsα*, to an imprinted region of distal mouse chromosome 2 provides some support for this theory.^{59,60} However, since equivalent reductions in *Gsα* bioactivity have been found in PHP Ia and PPHP in all membranes so far tested, imprinting would necessarily be operating at a tissue or even cell specific level.⁵⁵

Differential diagnosis

Confusion often arises between McCune-Albright syndrome and AHO although the two are clinically entirely distinct. Remarkably, both result from mutations in the *Gsα* gene; however, in the McCune-Albright syndrome these are gain of function mutations and occur on a somatic mosaic basis.⁶¹

There is considerable phenotypic overlap between PPHP and acrodysostosis^{62,63} since short stature, brachymetaphalangism, advanced bone age, mental retardation, and many

radiological features are common to both. The generalised metacarpal and phalangeal shortening characteristic of acrodysostosis has also been observed in confirmed cases of AHO (Wilson and Trembath, unpublished observations).⁶⁴ Cutaneous ossification does not occur in acrodysostosis but is not universal in PPHP. Pronounced nasal hypoplasia is considered a distinguishing feature of acrodysostosis but has been observed in patients with PHP Ia.⁶⁴

Similar overlap occurs with the isolated brachydactyly syndromes E and D⁶⁵ since they may be radiologically indistinguishable from AHO³⁴ and have been observed to segregate in families where AHO is segregating.⁶⁶ Acrodysostosis and the brachydactylies may represent poles of the phenotypic spectrum of AHO and *Gsα* evaluation will help to resolve this issue.

Other syndromes including Turner's syndrome and multiple familial exostoses are associated with short stature and metacarpal shortening but rarely cause diagnostic confusion.

The soft tissue ossification of myositis ossificans is distinguishable by its location in muscle and frequently painful, disabling nature.⁶⁷ Furthermore, the characteristic single phalanx, or phalangeal fusion of the big toe and shortening of the first metacarpal or proximal phalanx of the thumb in myositis ossificans contrasts with the pattern of shortening in AHO.

AHO has been observed in association with chromosomal abnormalities, specifically 15q⁶⁸ and 2q⁶⁹ deletions, but coincidental pathology was not excluded by *Gsα* measurement.

Other forms of PHP are distinguishable on endocrine and metabolic grounds (table 1).

LCW is an Medical Research Council Clinical Training Fellow. RCT gratefully acknowledges grant support of the Research Trust for Metabolic Diseases of Childhood. The authors wish to thank the many clinicians who have notified us of patients with AHO.

- 1 Albright F, Burnett CH, Smith PH, Parson W. Pseudohypoparathyroidism—an example of 'Seabright-Bantam syndrome'. *Endocrinology* 1942;30:922–32.
- 2 Albright F, Forbes AP, Henneman PH. Pseudo-pseudohypoparathyroidism. *Trans Assoc Am Physicians* 1952;65:337–50.
- 3 Houslay MD. G-protein linked receptors: a family probed by molecular cloning and mutagenesis procedures. *Clin Endocrinol* 1992;36:525–34.
- 4 Spiegel AM, Weinstein LS, Shenker A. Abnormalities in G protein-coupled signal transduction pathways in human disease. *J Clin Invest* 1993;92:1119–25.
- 5 Spiegel AM, Shenker A, Weinstein LS. Receptor-effector coupling by G proteins: implications for normal and abnormal signal transduction. *Endocr Rev* 1992;13:536–65.
- 6 Simon MI, Strathmann MP, Gautam N. Diversity of G proteins in signal transduction. *Science* 1991;252:802–8.
- 7 Levine MA, Downs RW Jr, Marx SJ, Singer M, Aurbach GS, Spiegel AM. Deficient activity of guanine nucleotide regulatory protein in erythrocytes from patients with pseudohypoparathyroidism. *Biochem Biophys Res Commun* 1980;94:1319–24.
- 8 Farfel Z, Brickman AS, Kaslow HR, Brothers VM, Bourne HR. Defect of receptor-cyclase coupling protein in pseudohypoparathyroidism. *N Engl J Med* 1980;303:237–42.
- 9 Farfel Z, Bourne HR. Deficient activity of receptor-cyclase coupling protein in platelets of patients with pseudohypoparathyroidism. *J Clin Endocrinol Metab* 1980;51:1202–4.
- 10 Bourne HR, Kaslow HR, Brickman AS, Farfel Z. Fibroblast defect in pseudohypoparathyroidism, type I: reduced activity of receptor-cyclase coupling protein. *J Clin Endocrinol Metab* 1981;53:636–40.
- 11 Downs RW Jr, Levine MA, Drezner MK, Burch WM Jr, Spiegel AM. Deficient adenylate cyclase regulatory protein in renal membranes from a patient with pseudohypoparathyroidism. *J Clin Invest* 1983;71:231–5.

- 12 Levine MA, Jap TS, Mauseth RS, Downs RW, Spiegel AM. Activity of the stimulatory guanine nucleotide-protein is reduced in erythrocytes from patients with pseudohypoparathyroidism and pseudopseudohypoparathyroidism: biochemical, endocrine, and genetic analysis of Albright's hereditary osteodystrophy in six kindreds. *J Clin Endocrinol Metab* 1986;62:497-502.
- 13 Chase LR, Melson GL, Aurbach GD. Pseudohypoparathyroidism: defective excretion of 3',5'-AMP in response to parathyroid hormone. *J Clin Invest* 1969;48:1832-44.
- 14 Mallette LE, Kirkland JL, Gagel RF, Law WM Jr, Heath H. III. Synthetic human parathyroid hormone-(1-34) for the study of pseudohypoparathyroidism. *J Clin Endocrinol Metab* 1988;67:964-72.
- 15 Tomlinson S, Hendy GN, O'Riordan JLH. A simplified assessment of response to parathyroid hormone in hypoparathyroid patients. *Lancet* 1976;i:62-4.
- 16 Breslau NA. Pseudohypoparathyroidism: current concepts. *Am J Med Sci* 1989;298:130-40.
- 17 Abou Samra AB, Juppner H, Force T, et al. Expression cloning of a common receptor for parathyroid hormone and parathyroid hormone-related peptide from rat osteoblast-like cells: a single receptor stimulates intracellular accumulation of both cAMP and inositol triphosphates and increases intracellular free calcium. *Proc Natl Acad Sci USA* 1992;89:2732-6.
- 18 Spiegel AM. Pseudohypoparathyroidism. In: Scriver CR, Beaudet AL, Sly WS, Valle D, eds. *The metabolic basis of inherited disease*. 6th ed. New York: McGraw Hill, 1989: 2013-27.
- 19 Fitch N. Albright's hereditary osteodystrophy: a review. *Am J Med Genet* 1982;11:11-29.
- 20 de Wijn EM, Steendijk R. Growth and development in a girl with pseudohypoparathyroidism and hypoparathyroidism. *Acta Paediatr Scand* 1982;71:657-60.
- 21 de Wijn EM, Steendijk R. Growth and maturation in pseudo-hypoparathyroidism: a longitudinal study in 5 patients. *Acta Endocrinol* 1982;101:223-6.
- 22 Geominne L. Albright's hereditary poly-osteochondrodystrophy. *Acta Genet Med Gemellol* 1965;14:226-81.
- 23 Steinbach HL, Rudhe U, Jonsson M, Young DA. Evolution of skeletal lesions in pseudohypoparathyroidism. *Radiology* 1965;85:670-6.
- 24 Farfel Z, Friedman E. Mental deficiency in pseudo-hypoparathyroidism type I is associated with Ns-protein deficiency. *Ann Intern Med* 1986;105:197-9.
- 25 Spranger JW. Skeletal dysplasias and the eye: Albright's hereditary osteodystrophy. *Birth Defects* 1969;5:122-8.
- 26 Eyre WG, Reed WB. Albright's hereditary osteodystrophy with cutaneous bone formation. *Arch Dermatol* 1971;104: 635-42.
- 27 Brook CGD, Valman HB. Osteoma cutis and Albright's hereditary osteodystrophy. *Br J Dermatol* 1971;85:471-5.
- 28 Prendiville JS, Lucky AW, Mallory SB, Mughal Z, Mimouni F, Langman CB. Osteoma cutis as a presenting sign of pseudohypoparathyroidism. *Pediatr Dermatol* 1992;9: 11-8.
- 29 Steinbach HL, Young DA. The roentgen appearance of pseudohypoparathyroidism (PH) and pseudo-pseudohypoparathyroidism (PPH). Differentiation from other syndromes associated with short metacarpals, metatarsals, and phalanges. *Am J Roentgenol* 1966;97:49-66.
- 30 Illum F, Dupont E. Prevalences of CT-detected calcification in the basal ganglia in idiopathic hypoparathyroidism and pseudohypoparathyroidism. *Neuroradiology* 1985;27:32-7.
- 31 Wong S, Zakov ZN, Albert DM. Scleral and choroidal calcifications in a patient with pseudohypoparathyroidism. *Br J Ophthalmol* 1979;63:177-80.
- 32 Schuster V, Sandhage K. Intracardiac calcifications in a case of pseudohypoparathyroidism type Ia (PHP-Ia). *Pediatr Cardiol* 1992;13:237-9.
- 33 Hewitt M, Chambers TL. Early presentation of pseudohypoparathyroidism. *J R Soc Med* 1988;81:666-7.
- 34 Poznanski AK, Werder EA, Giedion A. The pattern of shortening of the bones of the hand in PHP and PPHP - a comparison with brachydactyly E, Turner syndrome, and acrodysostosis. *Radiology* 1977;123:707-18.
- 35 Ray EW, Gardner LI. Pseudo-pseudo-hypoparathyroidism as an allelic form of pseudo-hypoparathyroidism: report of the youngest case, with description of a diagnostic screening technique. *Am J Dis Child* 1959;96:599-601.
- 36 Drezner MK, Neelon FA. Pseudohypoparathyroidism. In: Stanbury JB, Wyngaarden JB, Fredrickson DS, Goldstein JL, Brown MS, eds. *The metabolic basis of inherited disease*. 5th ed. New York: McGraw Hill, 1983:1508-27.
- 37 Koch T, Lehnhardt E, Bottinger H, et al. Sensorineural hearing loss owing to deficient G proteins in patients with pseudohypoparathyroidism: results of a multicentre study. *Eur J Clin Invest* 1990;20:416-21.
- 38 Weinstock RS, Wright HN, Spiegel AM, Levine MA, Moses AM. Olfactory dysfunction in humans with deficient guanine nucleotide-binding protein. *Nature* 1986;322:635-6.
- 39 Ritchie GM. Dental manifestations of pseudohypoparathyroidism. *Arch Dis Child* 1965;40:565-72.
- 40 Faull CM, Welbury RR, Paul B, Kendall-Taylor P. Pseudohypoparathyroidism: its phenotypic variability and associated disorders in a large family. *Q J Med* 1991;78: 251-64.
- 41 Croft LK, Witkop CJ Jr, Glas JE. Pseudohypoparathyroidism. *Oral Surg* 1965;20:758-70.
- 42 Burnstein MI, Kottamasu SR, Pettifor JM, Sochett E, Ellis BI, Frame B. Metabolic bone disease in pseudohypoparathyroidism: radiologic features. *Radiology* 1985; 155:351-6.
- 43 Kerr D, Hosking DJ. Pseudohypoparathyroidism: clinical expression of PTH resistance. *Q J Med* 1987;65:889-94.
- 44 Levine MA, Jap TS, Hung W. Infantile hypothyroidism in two sibs: an unusual presentation of pseudohypoparathyroidism type Ia. *J Pediatr* 1985;107:919-22.
- 45 Weisman Y, Golander A, Spirer Z, Farfel Z. Pseudohypoparathyroidism type Ia presenting as congenital hypothyroidism. *J Pediatr* 1985;107:413-5.
- 46 Levine MA, Downs RW Jr, Moses AM, et al. Resistance to multiple hormones in patients with pseudohypoparathyroidism. Association with deficient activity of guanine nucleotide regulatory protein. *Am J Med* 1983;74: 545-56.
- 47 Shima M, Nose O, Shimizu K, Seino Y, Yabuuchi H, Saito T. Multiple associated endocrine abnormalities in a patient with pseudohypoparathyroidism type Ia. *Eur J Pediatr* 1988;147:536-8.
- 48 Wolfsdorf JL, Rosenfield RL, Fang VS, Kobayashi R, Razdan AK, Kim MH. Partial gonadotrophin-resistance in pseudohypoparathyroidism. *Acta Endocrinol* 1978;88:321-8.
- 49 Kozasa T, Itoh H, Tsukamoto T, Kaziro Y. Isolation and characterization of the human G α gene. *Proc Natl Acad Sci USA* 1988;85:2081-5.
- 50 Gejman PV, Weinstein LS, Martinez M, et al. Genetic mapping of the Gs- α subunit gene (GNAS 1) to the distal long arm of chromosome 20 using a polymorphism detected by denaturing gradient gel electrophoresis. *Genomics* 1991;9:782-3.
- 51 Rao VV, Schnitzler S, Hansmann I. G protein Gs alpha (GNAS 1), the probable candidate gene for Albright hereditary osteodystrophy, is assigned to human chromosome 20q12-q13.2. *Genomics* 1991;10:257-61.
- 52 Patten JL, Johns DR, Valle D, et al. Mutation in the gene encoding the stimulatory G protein of adenylate cyclase in Albright's hereditary osteodystrophy. *N Engl J Med* 1990;322:1412-9.
- 53 Weinstein LS, Gejman PV, Friedman E, et al. Mutations of the Gs alpha-subunit gene in Albright hereditary osteodystrophy detected by denaturing gradient gel electrophoresis. *Proc Natl Acad Sci USA* 1990;87:8287-90.
- 54 Miric A, Vecchio JD, Levine MA. Heterogeneous mutations in the gene encoding the alpha-subunit of the stimulatory G protein of adenylate cyclase in the Albright hereditary osteodystrophy. *J Clin Endocrinol Metab* 1993;76:1560-8.
- 55 Wilson LC, Oude Luttikhuis MEM, Fraser WD, Clayton PT, Trembath RC. Parental origin of Gs α gene mutations in Albright's hereditary osteodystrophy. *J Med Genet* (in press).
- 56 Weinstein LS, Gejman PV, de Mazancourt P, American N, Spiegel AM. A heterozygous 4-bp deletion mutation in the Gs alpha gene (GNAS1) in a patient with Albright hereditary osteodystrophy. *Genomics* 1992;13:1319-21.
- 57 Oude Luttikhuis MEM, Wilson LC, Leonard JV, Trembath RC. Characterization of a de novo 43 base pair deletion of the Gs α gene in Albright hereditary osteodystrophy. *Genomics* 1994;21:455-7.
- 58 Davies SJ, Hughes HE. Imprinting in Albright's hereditary osteodystrophy. *J Med Genet* 1993;30:101-3.
- 59 Ashley PL, Ellison J, Sullivan KA, Bourne HR, Cox DR. Chromosomal assignment of the murine G α and G α genes. Implications for the obese mouse. *J Biol Chem* 1987;262:15299-301.
- 60 Cattanch BM, Kirk M. Differential activity of maternally and paternally derived chromosome regions in mice. *Nature* 1985;315:496-8.
- 61 Weinstein LS, Shenker A, Gejman PV, Merino MJ, Friedman E, Spiegel AM. Activating mutations of the stimulatory G protein in the McCune-Albright syndrome. *N Engl J Med* 1991;325:1688-95.
- 62 Robinow M, Pfeiffer RA, Gorlin RJ, et al. Acrodysostosis: a syndrome of peripheral dysostosis, nasal hypoplasia and mental retardation. *Am J Dis Child* 1971;121:195-203.
- 63 Butler MG, Rames LJ, Wadlington WB. Acrodysostosis: report of a 13-year-old boy with review of literature and metacarpophalangeal pattern profile analysis. *Am J Med Genet* 1988;30:971-80.
- 64 Ablow RC, Hsia YE, Brandt IK. Acrodysostosis coinciding with pseudohypoparathyroidism and pseudo-pseudohypoparathyroidism. *Am J Roentgenol* 1977;128:95-9.
- 65 Fitch N. Classification and identification of inherited brachydactylies. *J Med Genet* 1979;16:36-44.
- 66 Graudal N, Milman N, Nielsen LS, Niebuhr E, Bonde J. Coexistent pseudohypoparathyroidism and D brachydactyly in a family. *Clin Genet* 1986;30:449-55.
- 67 Connor JM, Evans DAP. Fibrodysplasia ossificans progressiva: the clinical features and natural history of 34 patients. *J Bone Joint Surg (Br)* 1982;64:76-83.
- 68 Hedeland H, Berntorp K, Arheden K, Kristofferson U. Pseudohypoparathyroidism type I and Albright's hereditary osteodystrophy with a proximal 15q chromosomal deletion in mother and daughter. *Clin Genet* 1992;42: 129-34.
- 69 Phelan MC, Rogers RC, Byrd LK. Albright hereditary osteodystrophy and del(2)(q37) in two unrelated individuals. *Am J Hum Genet* 1993;53 (suppl): 484.