Syndrome of the month

Cleft hand/foot: clinical and developmental aspects

Paul W Buss

Abstract

Isolated limb reduction defects occur in approximately 1 in 2000 live births within which central ray anomalies are an important subgroup. Most affected persons have mild or moderate functional impairment. Considerable psychological morbidity may also occur. While there have been major strides forwards in our understanding of vertebrate limb development, the mechanisms responsible for central ray deformities remain poorly understood. Several case reports of central clefting anomalies associated with chromosomal rearrangements or interstitial deletions of 7q21.2-q21.3 suggest that this chromosomal region is important for limb development.

(J Med Genet 1994;31:726-730)

While the term ectrodactyly literally means "missing digits", it is now used as a descriptive term for a central ray anomaly that presents clinically as a central cleft of the hands or feet.¹ Historically, terms such as "lobster claw"² and split hand³ have been used to describe this defect which results from absence, or reduction, of one or more of the three central digits (fig 1). The earliest known description of cleft hand was that by Pare⁴ although Barsky⁵ suggested that this report represented two fingered hemimelia rather than true cleft hand. Central ray defects, according to Swanson's modified classification, represent terminal longitudinal limb anomalies.⁶ Cleft limb has a number of associations and as a result it can be

a useful major feature for syndrome diagnosis.

Large families with isolated cleft limbs have been described which usually show autosomal dominant inheritance and marked variability in expression and penetrance,⁷ although several autosomal recessive pedigrees have also been reported.⁸⁻¹⁰ Birch-Jensen¹¹ estimated the incidence of cleft hand deformity at 1 in 90 000 births. More recent epidemiological studies suggest the incidence of central ray anomalies to be closer to 1 in 10 000 (approximately 0.7 per 10 000 live births).^{12,13} The data of Czeizel *et al*¹⁴ indicate that three-quarters (77%) of cases present with unimelic involvement.

In 1960 Stevenson and Jennings¹⁵ suggested an abnormal segregation pattern in families with isolated ectrodactyly (pedigree analysis showed an excess of affected sons of affected fathers). David,¹⁶ when describing three pedigrees with isolated cleft limbs, suggested the possibility of germinal mosaicism. X linked inheritance has also been described in a seven generation Pakistani kindred.¹⁷

Ford¹⁸ hypothesised that "ectrodactyly" represented an anomaly that could be explained by a structural chromosomal defect although at that time it was acknowledged that techniques for chromosome analysis were poor. Since then a number of case reports have indicated that the 7q21.2-q21.3 region is a candidate site for cleft limbs.¹⁹⁻²⁰ These have resulted in considerable molecular interest in this region.

Clinical features

Clinical problems associated with isolated cleft hand and foot are usually cosmetic and psy-

Figure 1 Typical central clefts of hands and feet; wide central clefts with normal marginal (radial and ulnar) rays. In this case the child had EEC syndrome.

Department of
Neonatal Medicine,
University of Bristol,
Bristol BS2 8EG, UK
P W Buss

Correspondence to
Dr Buss, Department of
Paediatrics, Royal Gwent
Hospital, Newport, Gwent
NP9 2UB, UK.

Table 1 List of conditions in which cleft limb may be a feature (McKusick No)

EEC syndrome (129900)
Cornelia de Lange syndrome (122430)
Acrorenal syndrome (102520)
Focal dermal hypoplasia (305600)
Ectrodactyly and cleft palate syndrome (129830)
Ectrodactyly/mandibulofacial dysostosis (183700)
Ectrodactyly and macular dystrophy (185800)

Non-specific associations with several anomalies are noted by Czeizel *et al*¹⁴ and also with neural tube defects.

chological rather than functional. Only in the less common variant monodactylous ectrodactyly, where radial rays are also hypoplastic, is function severely compromised. Considerable anatomical variability occurs. Cleft hand has been described with pre- and postaxial duplication and reduction anomalies.¹ Absence of the central digits is the major feature with varying association with both bony and soft tissue syndactyly. Many cases will merely have one limb affected and others can have tetramelic involvement. Birch-Jensen¹¹ found that 58% of cases of cleft hands were bilateral with an equal sex distribution. These clinical findings were confirmed by Barsky.⁵ The majority of cases of cleft hand are sporadic. In Birch-Jensen's study,¹¹ 50% had a family history of limb defects while only 20% of Barsky's group⁵ were familial. The study of Czeizel *et al*¹⁴ showed a high percentage of presentation as atypical forms, though this is based on a "wide" interpretation as to what actually constitutes atypical split hand/foot. The definition of this varies between published reports but actually serves relatively little clinical purpose. It is also subject to wide differences in clinical interpretation. A consensus might be that atypical cleft limb occurs when there is not total absence of the long finger or when there are, in addition, reduction anomalies affecting the marginal rays. This tends to be (though not always) sporadic and unilateral, affecting mainly the upper limbs.

Clinically, few functional problems occur in the upper limbs provided opposition is possible. However when a preaxial anomaly coexists or when monodactyly (a single digit) occurs, marked limitation of function can result. Functional limitation can result when central clefts are particularly wide or deep and operative management to close these can often result in significant improvement in function. Limitation of function owing to syndactyly of the fingers may also occur and this also can be overcome surgically.²⁷ In the lower limbs the major practical difficulty encountered is one of obtaining adequately wide fitting shoes.

Many persons with "ectrodactyly" experience psychological problems. These often emerge in mid-childhood as a reluctance to use hands, even a need to hide them. Non-verbal communication may be affected by this and common important gestures, such as hand shaking, are often avoided. These problems have frequently not been overcome even in adult life (Buss, unpublished data). Providing adequate reassurance to parents of affected children may allow them to help their children develop a positive body image. Anecdotal reports of cases with tetramelic involvement who

are professional artists, speed typists, and even marathon runners may also be of considerable reassurance.

Radiology usually reflects the clinical findings but sometimes transverse metacarpals or proximal phalanges affect the appearance. Occasionally atypical cleft hand and monodactyly coexist (fig 2). In the EEC syndrome (the ectodermal dysplasia, ectrodactyly, and clefting syndrome), preaxial anomalies also may coexist with a central cleft. While careful examination of the parents' hands may be helpful, there is little extra information to be gained from routine radiography of the hands and feet of parents who are clinically normal.

Differential diagnosis: clinical approach

Recognition of a central ray deformity may be a useful clue for the diagnosis of a number of underlying disorders (EEC syndrome, Cornelia de Lange syndrome, acrorenal syndrome, etc) (table 1). When presented with a neonate with a central ray defect diagnostic difficulties may arise. In such cases magnification of nails and dental x rays may help in identifying involvement of ectodermal structures (difficult in the neonate/infant with oligosymptomatic EEC syndrome). It is also worthwhile reviewing these children during the first year to confirm normal development of ectodermal structures and, if uncertain of this, electron microscopic examination of the hair shaft may be helpful in diagnosing ectodermal dysplasia (Buss, unpublished data). In the newborn it is important to document hair distribution and birth weight. A renal ultrasound scan may help make the diagnosis of the EEC or acrorenal syndrome. In older children and adults with "isolated" ectrodactyly, one often notes minor associated ectodermal anomalies (of the teeth in particular).¹ Language milestones should be checked. Ectrodactyly associated with conductive deafness and mandibular hypoplasia has been described in a father and son by Patterson and Stevenson²⁸ and more recently with sensorineural deafness by Raas-Rothschild *et al*²⁹ although the latter report looks distinctly similar to oligosymptomatic EEC syndrome. The otic capsule is interestingly also of ectodermal origin.³⁰ Opitz and Frias³¹ described an association with cleft palate as a separate entity, the ECP syndrome. Karsch³² recognised

Figure 2 Combined monodactyly with atypical cleft deformity showing some preservation of marginal rays in right hand.

Table 2 Review of reported cases of ectromelia associated with anomalies of chromosome 7q21

Author	Clinical description	7q deletion/karyotype
Del Porto <i>et al</i> ¹⁹	Male infant, low set ears, hypotonia, beaked nose, bilateral cleft foot	del(7) (q11–q21 or q22)
Pfiefer <i>et al</i> ²⁰	Female aged 2 years, microcephaly, low set ears, mental retardation, cleft palate, R cleft hand and foot	del(7) (q11.2–q22.1)
Tajara <i>et al</i> ²¹	Male aged 3 years, microcephaly, anteverted nares, micrognathia, retardation, clefting of three limbs	del(7) (q11.2–q22)
Morey <i>et al</i> ²²	Premature male, R cleft hand, amelia of lower limbs, facial dysmorphism, ASD. Died at 7 months	del(7) (q21.3–q31.3)
Sharland <i>et al</i> ²³	Male infant, tetramelic ectrodactyly, low set ears, hypertelorism	Complex rearrangement t(5;7;9) with breaks at 7q21.2 and 7q31.3
Roberts <i>et al</i> ²⁴	Male, tetramelic ectrodactyly, mild facial dysmorphism and mild developmental delay	Complex rearrangement (intrachromosomal) del(7) (q21.3–22.1)
Naritomi <i>et al</i> ²⁵	Male, trimelic ectrodactyly, bifid uvula, high arched palate	ins(3;7) (q21;q34q22). Break at 7q22q34
Genuardi <i>et al</i> ²⁶	5 generation family, 7 affected, 5 male. Unimelic (3 cases), tetramelic (4 cases). Two with sensorineural deafness. One case with tibial hypoplasia	Balanced translocation t(2;7) (q21.1;q22.1)

an association with impaired vision owing to cataract and macular dystrophy.

For parents of isolated cases empirical recurrence risks are low but there is a potential risk to offspring of affected persons of 50%. With possible heterogeneity and altered penetrance risks may be lower.³³ Certainly isolated cleft limbs can produce some very difficult diagnostic as well as counselling situations. Constriction amputation of digits may be difficult to distinguish from unimelic atypical cleft limb. Clues as to a constrictive aetiology might include the observation of amniotic strands at birth or lymphoedema distal to a possible constriction ring. Soft tissue syndactyly also occurs in constriction anomalies but is usually distal.³⁴ In such cases no other bony anomalies are usually radiologically present and occasionally the astute neonatal resident will have asked for the placenta to be examined showing an amniotic tear. Unilateral limb reduction deficits may also have a vascular origin and the clinician should be aware of the relevant points in the history, for example, chorionic villus sampling, presence of twins (reversed arterial perfusion), and any occurrences that may have compromised fetal circulation. It can be difficult to separate some of these potential causes and in the atypical isolated unimelic case explanation of a number of potential aetiologies may occasionally be necessary.

Cleft hand and foot malformation and 7q abnormalities

Eight recent case reports of ectrodactyly and 7q21.2–q21.3 anomalies are summarised in table 2. It is of interest that six of seven isolated cases reported are male (note the preponderance of males observed by Stephenson and Jennings¹⁵ and in the familial cases of Czeizel *et al*¹⁴). The significance of this is unclear. Genuardi *et al*²⁶ recently reported a family segregating a balanced translocation with a breakpoint at 7q22.1, in which two out of nine translocation carriers had limb deficiencies (similar abnormalities were present in a further five persons in whom chromosome analysis was not possible). Overall there is a wide range of clinical involvement from apparently isolated cleft limbs to ectromelia associated with fatal anomalies. None of the isolated cases had uni-

lateral cleft limbs, although three members of a family²⁶ showed unimelic involvement. The majority of the reported cases have typical cleft limbs.

Few other reports exist that link central ray anomalies with chromosome 7. Klep-de Pater *et al*³⁵ described a female infant with cleft palate and digital reduction deformities of the second and fifth digits of the right hand (not typical central cleft) and no evidence of ectodermal dysplasia.³⁵ This case had a karyotype 46,XX, del(7) (q11→q22). A number of additional case reports of deletions involving the 7q22 band have not been associated with ectrodactyly.³⁶

Aetiology and developmental aspects

The general association of cleft limb with anomalies of ectodermal structures is not particularly surprising as there are similarities in limb development to that of several ectodermal structures. Wolpert's experiments which confirmed earlier observations of a crucial reliance of embryonic limb ectoderm on the underlying mesoderm are not merely of historical importance.³⁷ He postulated the presence of a signalling mechanism (probably a diffusible morphogen) which could be interpreted as a positional signal for regional development within the progress zone of the developing limb bud mesoderm. Historically it has seemed likely that the endogenous signalling morphogen in the limb bud of mammals may be related to retinoic acid (possibly 3'4'didehydroretinoic acid).³⁸ However an exciting recent report indicates that a newly identified gene encoding the signalling molecule, Sonic hedgehog, may be involved in establishing polarising activity in the developing limb bud and hence antero-posterior patterning.³⁹

Developmental digit anomalies such as those seen in central ray anomalies, however, are not easily explained by a diffusible morphogen model. Homeobox genes are consistently expressed in the developing limb bud and digits in a way that reflects their location within a homeobox cluster (temporospatial colinearity).⁴⁰ These genes are activated sequentially by varying concentrations of retinoic acid within the limb bud. The expression of genes of the homeobox 4 (HOX D) type is necessary for normal limb development in mice⁴¹ and chick.⁴² Transcript expression can be predictably

altered in experimental animal models by implantation of retinoic acid soaked beads in strategic positions within the developing limb bud with resultant anomalies⁴³ including mirror hands and polydactyly. Central ray digit reduction anomalies have not yet been observed in these experiments. It may be that the anomaly represents a localised disturbance of HOX expression at the ectodermal/mesodermal junction. Recent work has shown distinct expression of MSX1 (HOX 7) and MSX2 (HOX 8) at this site in limbs^{44,45} and teeth.⁴⁶ Additionally combined expression of both HOX D and of HOX A (localised to 7p11) appears necessary for organised cellular differentiation in the developing limb bud. The msh related MSX1 and MSX2 genes, although widely distributed in the early limb bud, are later organised at the distal part of the bud and are responsible for limb growth, possibly by maintaining a pool of undifferentiated cells at the progress zone.⁴⁷ Although great strides have been made in our knowledge of vertebrate limb patterning in the context of central ray anomalies, it must also be conceded that currently there are no obvious developmental genes which map to the 7q candidate region. In addition there are no ectrodactylous mouse mutants which map to the homologous region on mouse chromosome 5. It is interesting to note, however, that mutation of MSX2 (HOX 8) has been associated with craniosynostosis and limb defects (including triphalangeal thumb and metatarsal shortening).⁴⁸

Mice have been bred with dactylaplasia which closely resembles the human cleft limb. In these it has been hypothesised that the anomaly occurs as a result of effects from two loci.⁴⁹ Whether this could be extrapolated to human cleft limb is debatable but a similar explanation was put forward by David.¹⁶ The association of cleft limb with absence of long bones⁵⁰ suggests that the defect may occasionally be a result of a common defect in regional limb specification. A child exposed in utero to a retinoic acid derivative during the crucial time of limb development with absent digits and syndactyly has been reported.⁵¹ More recently two case reports of limb anomalies in a fetus and child with prenatal exposure to isotretinoin have been reported.⁵² One could postulate that the mechanism underlying this may be a failure of differentiation resulting from misexpression of homeobox genes. However, it may be worth noting that other teratogens such as ethanol have also been postulated to cause ectrodactyly,⁵³ in this particular case with no clear underlying mechanism.

Evidence is clearly accumulating to make the 7q21.2–q21.3 locus a prime candidate for isolated ectrodactyly. While it has also been postulated as a locus for investigation in syndromes such as the EEC syndrome⁵⁴ (contiguous genes), it is likely that cleft limb represents a genetically heterogeneous condition and the 7q genes represent just one potential defect affecting a common pathway for limb differentiation.

The author wishes to acknowledge that his work on cleft limbs was conducted as Action Research Training Fellow in Medical Genetics at the Institute of Medical Genetics, University of Wales College of Medicine, Cardiff.

- 1 Tentamy S, McKusick V. *The genetics of hand malformations*. New York: Alan A Liss, 1978:53–63.
- 2 Goldmann EE. Beitrag zur Lehre von der Misbildungen der Extremitäten. *Beitr Klin Chir* 1891;7:239–56.
- 3 Meller J. Ein Fall von angeborener Spaltbildung der Hände und Füße. *Berl Klin Wochenschr* 1893;30:232–3.
- 4 Pare A. *Les oeuvres de M Ambroise Pare*. Paris: Gabriel Boun, 1575:5–8.
- 5 Barsky A. Cleft hand, classification, incidence and treatment. *J Bone Hand Surg (Am)* 1964;46:1707–19.
- 6 Swanson A. A classification for congenital limb defects. *J Hand Surg* 1978;1:8–12.
- 7 Lewis T, Embleton D. Split hand and split foot deformities: their types, origin and transmission. *Biometrika* 1908;6:26–58.
- 8 Verma I, Joseph R, Bharijansa S, Sadanand M. Split hand and split foot deformity inherited as an autosomal recessive trait. *Clin Genet* 1976;9:8–14.
- 9 Zlotogora J, Nubani N. Is there an autosomal recessive form of split hand and split foot malformation? *J Med Genet* 1988;25:138–40.
- 10 Friere Maia N. Autosomal recessive form of ectrodactyly and its implications in genetic counselling. *J Hered* 1971;62:53.
- 11 Birch-Jensen A. *Congenital deformities of the upper extremities*. Copenhagen: Ejnar Munksgaard, 1949.
- 12 Froster UG, Baird P. Limb reduction defects in over 1 million consecutive live births. *Teratology* 1989;39:127–35.
- 13 Calzolari E, Manioguigi O, Garani GP, Cocchi G, Magnani C, Milan M. Limb reduction defects in Emilia Romagna Italy: epidemiological and genetic study in 173 109 consecutive live births. *J Med Genet* 1990;29:353–7.
- 14 Czeizel AE, Vitez M, Kodaj T, Lenz W. An epidemiological study of isolated split hand/foot in Hungary, 1975–1984. *J Med Genet* 1993;30:593–6.
- 15 Stevenson A, Jennings L. Ectrodactyly – evidence in favour of disturbed segregation in offspring of affected males. *Ann Hum Genet* 1960;24:89–96.
- 16 David TJ. Dominant ectrodactyly and possible germinal mosaicism. *J Med Genet* 1972;9:316–20.
- 17 Ahmad M, Abbass H, Haque S, Gebjard F. X-chromosomally inherited split-hand/split foot anomaly in a Pakistani kindred. *Hum Genet* 1987;75:169–73.
- 18 Ford CE. Autosomal anomalies. *Second Conference on Congenital Malformations*, NY Foundation, 1963:25.
- 19 Del Porto G, D'Alessandro E, De Matties C, LoRe M, DiFusco I. Delzione interstiziale del braccio lungo del cromosoma 7 e sue correlazioni cliniche. *Pathologica (Suppl)* 1983;75:268–71.
- 20 Pfeifer RA. Interstitial deletion of chromosome 7 (q11.2 q22.1) in a child with split hand/split foot malformation. *Ann Genet (Paris)* 1984;27:45–8.
- 21 Tajara EH, Varella-Garcia M, Gusson ACT. Interstitial long arm deletion of chromosome 7 and ectrodactyly. *Am J Med Genet* 1989;32:192–4.
- 22 Morey MA, Higgins RR. Ectro-amelia syndrome associated with an interstitial deletion of 7q. *Am J Med Genet* 1990;35:95–9.
- 23 Sharland M, Patton M, Hill L. Ectrodactyly of hands and feet in a child with a complex translocation including 7q21.2. *Am J Med Genet* 1991;39:413–14.
- 24 Roberts SH, Hughes HE, Davies SJ, Meredith AL. Bilateral split hand and foot malformation in a boy with a de novo interstitial deletion of 7q21.3. *J Med Genet* 1991;28:479–81.
- 25 Naritomi K, Izumikawa Y, Tohma T, Hirayama K. Inverted insertion of chromosome 7q and ectrodactyly. *Am J Med Genet* 1993;46:492–3.
- 26 Genuardi M, Maria GP, Vincenza S, Bellussi A, Zollini M, Neri G. Split hand/split foot anomaly in a family segregating a balanced translocation with a breakpoint on 7q22.1. *Am J Med Genet* 1993;47:823–31.
- 27 Nutt JN, Flatt AE. Congenital central hand deficit. *J Hand Surg* 1981;6:48–60.
- 28 Patterson TJS, Stevenson AC. Craniofacial dysostosis and malformation of the feet. *J Med Genet* 1964;1:112–14.
- 29 Raas-Rothschild A, Aviram A, Ben-Ami T, et al. Newly recognised ectrodactyly/deafness syndrome. *J Craniofac Genet Dev Biol* 1989;9:121–7.
- 30 Mikaelian DO, Der Kaloustian VM, Shahin NA. Congenital ectrodactyly with hearing loss. *Arch Otolaryngol* 1970;92:85–9.
- 31 Opitz J, Frias JL. The ECP syndrome, another autosomal dominant cause of monodactylous ectrodactyly. *Eur J Ped* 1980;133:217–20.
- 32 Karsch J. Erbliche augenmissbildung in verbindung mit spalthand und fuss. *Z Augenheilkd* 1936;89:274–9.
- 33 Emery AEH. A problem for genetic counselling – split hand deformity. *Clin Genet* 1977;12:125–7.
- 34 Gellis S. Constrictive bands in the human. *Birth Defects* 1977;13(1):259–68.
- 35 Klep-de Pater JM, Bijlsma JB, Bleeker-Wagemakers EM. Two cases with different deletions of chromosome 7. *J Med Genet* 1979;15:161–3.
- 36 Winter R, Tickle C. Syndactylies and polydactylies – embryological overview and suggested classification. *Eur J Hum Genet* 1993;1:96–104.
- 37 Wolpert L. Mechanisms of limb development and malformation. *Br Med Bull* 1976;32:65–70.

- 38 Ragsdale CW Jr, Brockes J. Retinoids and their targets in vertebrate development. *Curr Opin Cell Biol* 1992;**3**: 928–34.
- 39 Riddle RD, Johnson RL, Laufer E, Tabin C. Sonic hedgehog mediates the polarizing activity of the ZPA. *Cell* 1993;**75**: 1401–16.
- 40 Lewis T. A gene complex controlling segmentation in drosophila. *Nature* 1978;**276**:565.
- 41 Dolle P, Izpisua-Belmonte JC, Falkenstein H, Renucci A, Duboule D. Coordinate expression of murine Hox 5 complex homeobox containing genes during limb pattern formation. *Nature* 1989;**342**:767–9.
- 42 Izpisua-Belmonte JC, Tickle C, Dolle P, Wolpert L, Duboule D. Expression of the homeobox HOX 4 genes and the specification of position in chick wing development. *Nature* 1991;**350**:585–9.
- 43 Bryant SV, Gardiner DM. Retinoic acid, local cell–cell interactions and pattern formation in vertebrate limbs. *Dev Biol* 1992;**152**:1–25.
- 44 Benoit R, Lyons G, Simandl BK, Atshushi R, Buckingham M. The apical ectodermal ridge regulates Hox 7 and Hox 8 gene expression in developing chick limb buds. *Genes Dev* 1991;**5**:2363–74.
- 45 Davidson D, Crawley A, Hill RE, Tickle C. Position dependent expression of two related homeobox genes in developing vertebrate limbs. *Nature* 1991;**352**:429–31.
- 46 McKenzie A, Leeming GL, Tarrett, Jowett A, Sharpe PT. Human Hox 7 in regional and temporal expression in craniofacial and tooth development. *Development* 1991;**111**:269–85.
- 47 Tickle C. A tool for transgenesis. *Nature* 1992;**358**:188–9.
- 48 Wang Jabs E, Muller I, Li X, *et al.* A mutation in the homeodomain of the human MSX2 gene in a family affected with autosomal dominant craniosynostosis. *Cell* 1993;**75**:443–50.
- 49 Chai CK. Dactylaplasia in mice. *J Hered* 1981;**72**:234–7.
- 50 Hoyne AD, Lyons K, Nyhan W, Pauli R, Robinow M. Autosomal dominant ectrodactyly and abnormalities of long bones of upper and lower limbs. *J Pediatr* 1987;**111**: 538–43.
- 51 McBride WG. Limb reduction deformities in a child exposed to isotretinoin in utero at 26–40 days gestation. *Lancet* 1985;**i**:1276.
- 52 Rizzo R, Lammer E, Parsano E, Pavone L, Argyle JC. Limb reduction defects in humans associated with prenatal isotretinoin exposure. *Teratology* 1991;**44**:599–604.
- 53 Hermann J, Pallister PD, Opitz JM. Tetraectrodactyly and other skeletal manifestations in the foetal alcohol syndrome. *Eur J Pediatr* 1980;**133**:221–6.
- 54 Qumiseveh MB. EEC syndrome (ectodermal dysplasia, ectrodactyly, cleft lip and palate) is on 7q11.2–q21.3. *Clin Genet* 1992;**42**:101.