A familial Xp+ chromosome, dup (Xq26.3→qter)

A I Vasquez, H Rivera, L Bobadilla, J A Crolla

Abstract

A maternally transmitted Xp+ chromosome was associated with an abnormal phenotype, including developmental delay and short stature, in two male cousins and their 12 year old aunt. The respective mothers were not mentally impaired but had short stature. The G banding pattern identified the extra chromosome segment as a repeat of Xq26.3→qter attached to an apparently intact Xp22.3 sub-band, so the Xp+ chromosome may be described as rea(X)(Xqter→p22.3::Xq26.3→Xqter). The rearranged chromosome was late replicating in 97 to 100% of the metaphases in the mothers but it was early replicating in 43% of the lymphocytes in the mentally defective female (n=100 cells/subject). Fluorescence in situ hybridisation using X and Y chromosome paints, as well as cosmids A and 1A1 specific for loci within Xq28, confirmed both the identity of the extra segment and the entirety of the Xp pseudoautosomal region. Therefore, the phenotypic consequences in this family can be related to the Xq26.3→qter functional disomy allowing for the effects of X inactivation in the female carriers.

(J Med Genet 1995;32:891-893)

Partial Xq duplications usually involve bands proximal to q26 and are often ascertained through mentally retarded males in whom a distinct clinical picture has been associated with

Xq13.1→q21.1 disomy.¹⁻³ Four duplications extending into Xq27-28 have been described in phenotypically abnormal females,⁴ but only a brother-sister pair⁵ and a further girl⁶ have had duplications limited to such distal bands. In the present context, only intrachromosomal Xq duplications are considered, that is, the numerous isochromosomes/isodicentric Xq and Xp;Xq translocations ascertained through female patients⁷ as well as extra isochromosomes Xq in male patients⁸ are disregarded. Here we report on a three generation pedigree in which a maternally transmitted Xp+ chromosome implying duplication Xq26.3→qter resulted in two boys and one girl with significant clinical abnormalities.

Family report

The proband (III.4, fig 1) was the first and only child born to a 25 year old 153 cm tall mother and a 26 year old father. He was delivered at term by caesarian section because of fetal distress; Apgar scores were 4 at one minute and 6 at five minutes. Birth weight was 2900 g. Thereafter, he had hypotonia, delayed milestones, failure to thrive, and respiratory infections requiring hospital management. At 5 months of age he had poor social contact and no head control. His length was 55 cm, weight 3850 g, and OFC 38 cm (all below the 3rd centile). He also had a tented upper lip, downturned corners of mouth, inguinal hernias, hypoplastic genitalia with undescended testes, and slender fingers and toes. There was neither ichthyosis nor radiological evidence of chondrodysplasia punctata. Family investigations disclosed two phenotypically abnormal relatives on the maternal side: a male cousin (III.3) and an aunt (II.6). The cousin was the third child of a 31 year old 152 cm tall mother and an unrelated 27 year old father; both sibs are healthy children. Delivery was at term and spontaneous but the infant required respiratory assistance because of severe asphyxia. Birth weight was 2500 g. His clinical manifestations at 7 months of age were very similar to those of the proband and included hypotonia, developmental delay, failure to thrive (length 58 cm, weight 4700 g, OFC 37.5 cm), fish shaped mouth, inguinal hernias, hypoplastic genitalia, and cryptorchidism. Ichthyosis and chondrodysplasia punctata were absent. He also had recurrent bronchopneumonia which resulted in death at 13 months of age. The 12 year old aunt had obvious short stature and mental retardation; microphthalmia and linear skin defects were not apparent but a detailed physical examination was denied. The maternal

Division de Genetica,
Instituto Mexicano del
Seguro Social,
Ap Postal 1-3838,
Guadalajara, Jalisco,
Mexico
A I Vasquez
H Rivera
L Bobadilla

Wessex Regional
Genetics Laboratory,
Salisbury District
Hospital,
Odstock, Salisbury,
Wiltshire SP2 8BJ, UK
J A Crolla

Correspondence to:
Dr Rivera.

Received 25 November 1994
Revised version accepted for
publication 28 June 1995

Figure 1 Pedigree.

Figure 2 Sex chromosome pairs showing the Xp+ chromosome with duplication q26.3→qter (the Xp+ is on the right in each pair). (A) G bands. (B) Replication patterns in female subjects documenting the inactive (left hand pairs) or active (right hand pairs) status of the abnormal X chromosome.

grandmother was also short statured whereas the remaining maternal aunt (II.4) had a normal phenotype and a height of 167 cm.

CYTOGENETIC STUDIES

Chromosome preparations from standard whole blood cultures of the proband and several relatives were analysed after GTG banding.⁹ Subsequently, X chromosome replication in the female carriers (see below) was assessed using a terminal 5-bromo-2'-deoxyuridine pulse followed by FPG staining.¹⁰ Whole chromosome painting with X and Y specific libraries (Cambio, Cambridge) was done on metaphase spreads of the proband and subject II.1. The methodology was essentially that indicated by the manufacturer. The origin of the extra segment (see below) was further assessed in the proband by means of fluorescence in situ hybridisation¹¹ using the Xq28 cosmids A and 1A1 specific for the loci L1CAM and DXS305/374 respectively.

All three phenotypically abnormal subjects as well as their mothers were found to have an Xp+ chromosome within an otherwise normal (male or female) karyotype; in contrast, a 46, XX karyotype was found in subject II.4. The G banding pattern identified the extra segment as a repeat of Xq26.3→qter attached to an apparently entire Xp22.3 sub-band (fig 2). Therefore, the Xp+ chromosome may be described as a rea(X) (Xqter→Xp22.3::Xq26.3→qter). In all three mothers the Xp+ was predominantly inactive (97 to 100% of the metaphases) whereas in the mentally retarded girl it was either late (57%) or early (43%) replicating (n = 100 cells/subject) (fig 2). The X chromosome paint showed that the extra segment consisted only of X chromosome sequences whereas the hybridisation pattern of cosmids A and 1A1 confirmed the distal Xq duplication (fig 3). The Y chromosome paint showed an apparently normal pseudoautosomal region at p22.3 in the rearranged X

chromosome (fig 3). Cell lines are not available from any of these people.

Discussion

It is uncertain whether this abnormal chromosome resulted from a recombination within an untraced (either germinal or constitutional) pericentric inversion or from an inter-homologue translocation. Although telomeric sequences at the breakpoint junction were not looked for, neither their presence nor absence may unequivocally elucidate the origin as a pericentric inversion, which may result from a single break at Xq26.3 with transposition of the Xq distal segment onto the Xp telomere and healing of the broken end.¹² Regardless of the mechanism, the present Xp+ chromosome looks like an actual recombinant.⁵

The two male infants presented in this paper and one previously published boy with a similar imbalance⁵ did not show ichthyosis or other exclusive stigmata of Xp22 nullisomy, a finding consistent with the very distal localisation of the Xp breakpoint in both rearrangements. Otherwise, males with duplications for other Xq segments also exhibit perinatal asphyxia, severe mental retardation, marked short stature, inguinal hernias, hypoplastic genitalia, and cryptorchidism.^{1-3,13} Of particular relevance here are three severely affected boys with mental retardation, hypotonia, no speech development, seizures, short stature, and microcephaly secondary to disomy for 5–10 Mb of Xq28 resulting from aberrant Xq-Yq interchanges.¹⁴ Remarkably, males with partial Xp duplications also show mental and growth retardation as major components of a phenotype that features sex reversion whenever the relevant gene is involved.^{15,16} Taken together, the reported duplications cover the entire X chromosome with the exception of Xp11.1→q11 and confirm the assertion that partial constitutional disomies of this chromosome are compatible with postnatal survival

Figure 3 Image enhanced FISH results on the Xp+ observed in the proband. (A) Painting with a whole Y chromosome paint library gave complete coverage of the euchromatic region of the normal Y (not shown) and cross hybridisation to the interstitial pseudoautosomal region in distal Xp of the abnormal X and also to Xq21. (B) The cosmid 1A1 (DXS305 374), which maps to Xq28, showed signal both at Xq28 and at the distal tip of Xp.

of the affected males.² The common phenotypic consequences of duplications for different X chromosome segments can hardly be regarded as discrete effects of specific genes but may rather be ascribed to a supragenetic mechanism.

Most females with Xq duplications appear to be phenotypically normal but at least six previous cases have presented with developmental delay, congenital anomalies, short stature, or ovarian dysfunction.⁴ The constant inactivation of the rearranged X chromosome documented in normal carriers as well as the opposite pattern found in one abnormal girl with a de novo duplication⁶ and the data in our family argue for a simple correlation between gene expression of the duplicated segment and phenotype; furthermore, the preferential inactivation of the duplicated Xq chromosome also observed in most phenotypically abnormal carriers⁴ may still agree with such a relationship as the inactivation pattern would vary among different tissues. Incidentally, the occurrence of short stature in some "normal" carriers^{4,13} (present study) points to a wide clinical spectrum and is consistent with the aforementioned correlation. Additional support for the crucial role of the partial X disomy in the female phenotype comes from the severe clinical consequences of non-inactivated small ring X chromosomes¹⁷⁻¹⁹ as well as from balanced X; autosome translocations whose significant association with mental retardation and congenital anomalies in female carriers has been related to functional disomy rather than to spreading of inactivation into the autosomal segment.²⁰

We thank Professor Patricia A Jacobs for helpful criticisms and advice and Dr Mark Hirst of the Institute of Molecular Medicine, John Radcliffe Hospital, Oxford, for the gift of cosmids A and 1A1.

- 1 Thode A, Partington MW, Yip MY, Chapman C, Richardson VF, Turner G. A new syndrome with mental retardation, short stature and an Xq duplication. *Am J Med Genet* 1988;30:239-50.

- 2 Schmidt M, Du Sart D, Kalitsis P, et al. Duplications of the X chromosome in males: evidence that most parts of the X chromosome can be active in two copies. *Hum Genet* 1991;86:519-21.
- 3 Yokoyama Y, Narahara K, Tsuji K, et al. Growth hormone deficiency and empty sella syndrome in a boy with dup(X)(q13.3→q21.2). *Am J Med Genet* 1992;42:660-4.
- 4 Aughton DJ, AlSaadi AA, Johnson JA, Transue DJ, Trock GL. Dir dup(X)(q13-qter) in a girl with growth retardation, microcephaly, developmental delay, seizures, and minor anomalies. *Am J Med Genet* 1993;46:159-64.
- 5 Mohandas T, Geller RL, Yen PH, et al. Cytogenetic and molecular studies on a recombinant human X chromosome: implications for the spreading of X chromosome inactivation. *Proc Natl Acad Sci USA* 1987;84:4954-8.
- 6 Magenis RE, Donlon TA, Canfield WM, Koler R, Bigley R. Partial duplication of X long arm, q26.3-qter: gene dose effects of G6PD, HPRT, factor VIII:C and factor IX. *Cytogenet Cell Genet* 1984;37:528.
- 7 Bjerglund Nielsen L, Boczkowski K, Mikkelsen M, Dahl G, Andersen E. Partial Turner's syndrome in four girls with Xq duplication and Xp deficiency. *Hum Genet* 1982;61:12-17.
- 8 Richer CL, Bleau G, Chapdelaine A, Murer-Orlando M, Lemieux N, Cadotte M. A man with isochromosome Xq Klinefelter syndrome with lack of height increase and normal androgenization. *Am J Med Genet* 1989;32:42-4.
- 9 Seabright M. A rapid banding technique for human chromosomes. *Lancet* 1971;ii:971-2.
- 10 Perry P, Wolff S. New Giemsa method for the differential staining of sister chromatids. *Nature* 1974;251:156-8.
- 11 Pinkel D, Landegent J, Collins C, et al. Fluorescence in situ hybridisation with human chromosome-specific libraries: detection of trisomy 21 and translocations of chromosome 4. *Proc Natl Acad Sci USA* 1988;85:9138-42.
- 12 Dobbs M, Overhauser J, Wasmuth JJ. Molecular analysis of a case of meiotic recombination leading to cri-du-chat syndrome. *Cytogenet Cell Genet* 1988;47:5-7.
- 13 Miller PR, Williams LJ, McKeown C. A case of Xq duplication in a growth retarded male infant. *Clin Cytogenet Bull* 1993;2:46.
- 14 Lahn BT, Ma N, Breg WR, Stratton R, Surti U, Page DC. Xq-Yq interchange resulting in supernormal X-linked gene expression in severely retarded males with 46,XYq- karyotype. *Nature Genet* 1994;8:243-50.
- 15 Arn P, Chen H, Tuck-Muller CM, et al. SRVX, a sex reversing locus in Xp21.2→p22.11. *Hum Genet* 1994;93:389-93.
- 16 Ogata T, Matsuo N. Testis determining gene(s) on the X chromosome short arm: chromosomal localisation and possible role in testis determination. *J Med Genet* 1994;31:349-50.
- 17 Van Dyke DL, Wiktor A, Palmer CG, et al. Ullrich-Turner syndrome with a small ring X chromosome and presence of mental retardation. *Am J Med Genet* 1992;43:996-1005.
- 18 Dennis NR, Collins AL, Crolla JA, Cockwell AE, Fisher AM, Jacobs PA. Three patients with ring(X) chromosomes and a severe phenotype. *J Med Genet* 1993;30:482-6.
- 19 Migeon BR, Luo S, Stasiowski BA, et al. Deficient transcription of XIST from tiny ring X chromosomes in females with severe phenotypes. *Proc Natl Acad Sci USA* 1993;90:12025-9.
- 20 Schmidt M, Du Sart D. Functional disomies of the X chromosome influence the cell selection and hence the X inactivation pattern in females with balanced X-autosome translocations: a review of 122 cases. *Am J Med Genet* 1992;42:161-9.