LETTERS TO THE EDITOR

The Gene Bomb

I would like to comment on Tom Wilkie's comments on *The Gene Bomb*¹ (*J Med Genet* 1997;34:438-9), since I feel he missed the whole point of the book and presented to the readers of this journal a distorted view of the issues I attempted to raise. They have implications for all of us.

First, the subtitle, which somewhat more accurately portrays the issues, *Does Higher Education and Advanced Technology Accelerate the Selection of Genes for Learning Disorders, ADHD, Addictive and Disruptive Behaviors?* was left out.

Second, no mention was made of part I of the book which reviews data showing the following.

(1) Epidemiological studies using structured diagnostic instruments have shown a highly significant increase in frequency, and decrease in the age of onset, in younger cohorts for all the major psychiatric disorders including depression, manic depressive disorder, schizoaffective disorder, suicide, obsessive-compulsive disorders, panic disorder, phobias, alcoholism, and drug abuse.

(2) Using the Achenbach Child Behavior Checklist, Achenbach performed epidemiological studies in 1976 and 1989. Of 112 behavioural scores, 42 were worse in 1989 while none was worse in 1976. In addition, behaviours relating to specific diagnoses, such as ADHD and conduct disorder, increased from 10% in 1976 to 18.2% only 13 years later. In 1976, 8.2% of children had to repeat a grade while in 1989 17.8% had to repeat a grade.

(3) Studies of the same localised region in Sweden by Gillberg have shown an increase in the frequency of autism from 4/10 000 in 1980, to 7.5/10 000 in 1984, to 11.5/10 000 in 1988.

(4) US Department of Education studies have shown a progressive increase in the frequency of learning disabilities from 890 000 cases in 1976 to 2 714 000 cases in 1993. At the same time the population has only increased from 210 000 000 to 240 000 000.

To date there have been no adequate explanations for these trends and observations, very few professionals, including geneticists, have either been aware of or concerned about these trends, and no one has offered any satisfactory explanation for them. I work with children with hereditary behavioural disorders (ADHD, Tourette syndrome, learning disabilities, autism, Asperger syndrome, conduct disorder, and others) on a daily basis and I and their parents are concerned. I am particularly concerned about the trend in learning disabilities. These have increased from 2 000 000 in 1988 to 2 700 000 in 1993. At this rate the frequency of learning disorders is doubling every 15 years.

In attempting to understand these trends, I noticed the following. First, all of these disorders have a strong genetic component. Second, there is a strong inverse correlation between the frequency of these disorders,

especially learning disorders, and years of education. Not surprisingly, those with learning disorders tend to drop out of school, while those without learning disorders are more likely to go to college and possibly graduate school. Third, there is an extraordinarily high inverse correlation between the number of years of education and the age of initiation of child birth. At the beginning of this century only 2% of the population attended college. The rate dramatically increased after the second world war and now 38% attend college. All of these trends in behaviour disorders have occurred in the latter half of this century. At the present time the spread in years of the age of onset of childbearing, for those who drop out of school before completing a high school education versus those who go to college, is 8.9 years. Fourth, like compound interest, anything that turns over more rapidly will more rapidly increase in frequency. This can provide a very strong selection for the genes associated with learning disability, totally independent of the number of children people have. Comparing 1920 to 1990 shows that the magnitude of the selective force (spread in years by percent attending college) has increased 40 fold.

How does one respond to these observations or concerns? There are several possibilities. One is to shoot the messenger, as Wilkie has chosen to do. A second is to provide alternative explanations for these trends. I would prefer this so I can then stop worrying about the subject. The third is to accept it as a legitimate concern and either disprove the hypothesis or continue research into the genes involved in these disorders so that their frequencies in the population by age can be monitored.

In the book I suggested two approaches. The first were "motherhood and apple pie" approaches that no one should object to, such as special programmes to encourage children with ADHD and learning and related disorders to remain in school, and programmes for day care for children of college students so that if they wished, they could start a family before completing their education. The second approach is to continue (or accelerate) the research towards providing a better understanding of the genes involved. Among other things this would allow couples to participate in totally voluntary programmes by which they could make informed choices about reproductive decisions, as we now encourage people with a family history of cystic fibrosis or Huntington's disease to do.

Finally, I would like to point out that when the various behavioural disorders, most of which appear to be polygenic, are combined they occur in over 35% of the general population. This is 35 times the frequency of all the single gene disorders combined. I would suggest that those who propose that geneticists should not be involved in studies of the genetics of human behaviour, and prefer that we do not engage in responsible talk about some of the social implications of these studies, because it may give geneticists a bad name, are guilty of sticking their head in the sand. If geneticists don't become involved and play some role in attempting to understand the cause of these trends, I suspect no one will. In his book, *And the Band Played On*, Randy Shilts severely chastised the professional community for failing to mount an early response to the AIDS epidemic, because they preferred not to become involved or felt uncomfortable with the subject. Are we going

to be similarly chastised in the next century because we also failed to care, or felt uncomfortable with the subject, and made no attempt to understand these trends?

DAVID E COMINGS

Department of Medical Genetics, City of Hope Medical Center, Duarte, California 91010-0269, USA

1 Comings DE. *The gene bomb. Does higher education and advanced technology accelerate the selection of genes for learning disorders, ADHD, addictive, and disruptive behaviors?* Duarte, CA: Hope Press, 1996.

Gaucher disease plus

In a recent letter, Uyama *et al*¹ discussed a Japanese sibship of three subjects with glucocerebrosidase deficiency. These patients had oculomotor abnormalities characteristic of neuronopathic Gaucher disease along with several more unusual clinical manifestations described in other populations, including valvular calcifications, communicating hydrocephalus, and corneal opacities.²⁻⁵ Yet although the patients are reported to have glucocerebrosidase deficiency and point mutations on both glucocerebrosidase alleles, the authors suggest that the patients have a "Gaucher-like" disorder.¹

Gaucher disease is characterised by a vast phenotypic heterogeneity that is continually expanding.⁶ Recently, there has been a greater appreciation that diverse symptoms including congenital ichthyosis,⁷ hydrops fetalis,⁷ primary pulmonary hypertension,⁸ valvular calcifications,³⁻⁵ and even Parkinsonian symptoms⁹ can be seen in patients with Gaucher disease. Thus, we would suggest that the symptoms observed in the patients described by Uyama *et al*¹ are just part of the broader Gaucher phenotype.

Although the authors mention other genes near to glucocerebrosidase, they discount the likelihood of a contiguous gene effect.¹ Since the discovery of human metaxin,¹⁰ a gene located immediately downstream and transcribed convergently to the glucocerebrosidase pseudogene, we have characterised the locus more completely and have found that the region encompassing glucocerebrosidase is particularly gene rich.^{11,12} The physical relationship of metaxin and glucocerebrosidase on human chromosome 1q21 described by Uyama *et al*¹ is incorrect, for actually the pseudogene for human metaxin and not the gene is immediately downstream to the glucocerebrosidase gene.¹⁰ We suggest that patients with these more unusual Gaucher phenotypes may indeed have "Gaucher disease plus", an interruption in a second contiguous gene, rather than a "Gaucher-like" illness. In Hunter syndrome, it has recently been shown that some patients with atypical phenotypes have deletions including segments of nearby genes.¹³ Further characterisations of all the genes around the glucocerebrosidase locus may provide a more complete explanation for the phenotypic variation observed in Gaucher disease.

ELLEN SIDRANSKY

EDWARD I GINNS

Clinical Neuroscience Branch, National Institute of Mental Health, National Institutes of Health, Bethesda, MD, USA

1 Uyama E, Uchino M, Ida H, Eto Y, Owada M. D409H/D409H genotype in Gaucher-like disease. *J Med Genet* 1997;34:175.

2 Uyama E, Takahashi K, Owada M, *et al*. Hydrocephalus, corneal opacities, deafness, valvular

- heart disease, deformed toes and leptomenin-
geal fibrous thickening in adult siblings: a new
syndrome associated with β -glucocereb-
rosidase deficiency and a mosaic population of
storage cells. *Acta Neurol Scand* 1992;86:407-
20.
- 3 Chabas A, Cormand B, Grinberg D, et al. Unu-
sual expression of Gaucher's disease: cardio-
vascular calcifications in three sibs ho-
mozygous for the D409H mutation. *J Med
Genet* 1995;32:740-2.
 - 4 Abrahamov A, Elstein D, Gross-Tsur V, et al.
Gaucher's disease variant characterized by pro-
gressive calcification of heart valves and unique
genotype. *Lancet* 1995;346:1000-3.
 - 5 Beutler E, Kattamis C, Sipe J, Lipson M. 1342C
mutation in Gaucher's disease. *Lancet* 1995;
346:1637.
 - 6 Sidransky E, Ginns EI. Clinical heterogeneity
among patients with Gaucher's disease (clinical
conference). *JAMA* 1993;269:1154-7.
 - 7 Sidransky E, Sherer DM, Ginns EI. Gaucher
disease in the neonate: a distinct Gaucher phe-
notype is analogous to a mouse model created
by targeted disruption of the glucocerebrosi-
dase gene. *Pediatr Res* 1992;32:494-8.
 - 8 Dawson A, Elias DJ, Rubenson D, et al. Pulmo-
nary hypertension developing after Alglucerase
therapy in two patients with type 1 Gaucher
disease complicated by the hepatopulmonary
syndrome. *Ann Intern Med* 1996;125:901-4.
 - 9 Neudorfer O, Giladi N, Elstein D, et al. Occur-
rence of Parkinson's syndrome in type 1
Gaucher disease. *Q J Med* 1996;89:691-4.
 - 10 Long GL, Winfield S, Adolph KW, Ginns EI,
Bornstein P. Structure and organization of the
human metaxin gene (MTX) and pseudogene.
Genomics 1996;33:177-84.
 - 11 Sidransky E. New perspectives in type 2
Gaucher disease. *Adv Pediatr* (in press).
 - 12 Sidransky E, Ginns EI. Gaucher disease: the
best laid schemes of mice and men. *Baillière
Clin Haematol* (in press).
 - 13 Timms KM, Bondeson ML, Ansari-Lari MA, et
al. Molecular and phenotypic variation in
patients with severe Hunter syndrome. *Hum
Mol Genet* 1997;6:479-86.

Molecular characterisation of cystic fibrosis patients in the state of São Paulo (Brazil)

To characterise the CF patient population in
the state of São Paulo (Brazil), we estimated
the frequency of mutations $\Delta F508$, G542X,
N1303K, G551D, and R553X, which are
some of the most frequent mutations
worldwide. In a sample of 120 CF chromo-
somes, the frequencies we obtained were as
follows: $\Delta F508=31.7\%$, G542X=8.3%,
N1303K=2.5%. The G551D and R553X
mutations were not detected in this sample.
The mutation detection rate was 42.5%
based on the analysis of these five mutations.
We conclude that further screening for more
mutations will be necessary for molecular
diagnosis and carrier detection because of the
ethnic heterogeneity of this population. The
present study represents a step towards the
molecular characterisation of CF patients in
the state of São Paulo.

Cystic fibrosis is the most common auto-
somal recessive disease, with an incidence of
1 in 2000-3000 births in various groups.¹ The
CF gene was cloned in 1989.^{2,4} The major
mutation causing CF is a three base pair
deletion which results in the loss of phenyl-
alanine 508 ($\Delta F508$).⁴ This mutation is found
on about 70% of CF chromosomes world-
wide and varies considerably between popu-
lations. The CF Genetic Analysis Consor-
tium has identified over 620 other sequence
alterations in the CF gene. Most of these
mutations are rare. Only a few mutant alleles
have a worldwide frequency of 1% or greater,
although they may occur at higher frequen-
cies in selected populations.⁵

Knowledge of the spectrum of CF gene
mutations in different populations allows for
the possibility of genetic testing. In Brazil,
Raskin *et al*⁶ and Martins *et al*⁷ evaluated the
frequency of $\Delta F508$ in the state of São Paulo.
Their results showed frequencies of 52% (60/
116) and 31% (15/48), respectively. In view
of these differing results and in order to con-
tribute to the molecular characterisation of
CF patients in the state of São Paulo, we have
analysed the frequencies of CF mutations
 $\Delta F508$, G542X,⁸ N1303K,⁹ G551D,¹⁰ and
R553X.¹⁰

Sixty unrelated CF patients regularly
followed at the University (UNICAMP)
Hospital in Campinas were studied. Blood
samples were obtained from these patients,
who had been diagnosed with CF on the basis
of typical clinical manifestations or a positive
sweat test (>60 mEq/l) or both. Genomic
DNA was extracted from peripheral blood
leucocytes according to standard protocols.

DNA amplification was performed accord-
ing to a general procedure.¹¹ The primers and
annealing temperatures used for each ampli-
fied region have been reported before.^{4,12}

Amplified samples from each subject were
screened for the five previously mentioned
mutations. The mutation $\Delta F508$ was de-
tected by heteroduplex DNA formation.¹³
The mutations G542X and N1303K were
detected by digestion with *Bst*NI¹² and mu-
tations G551D and R553X were detected by
digestion with *Hinc*II followed by ASO
hybridisation. This combination of methods
was used because digestion with *Hinc*II by
itself does not distinguish mutations G551D
and R553X.¹²

The frequencies of mutations $\Delta F508$,
G542X, and N1303K in the sample studied
were 31.7%, 8.3%, and 2.5%, respectively.
Mutations G551D and R553X were not
detected.

The main objective of this study was to
contribute to the molecular characterisation
of CF patients in the state of São Paulo. The
observed frequency of the $\Delta F508$ mutation
(31.7%) is consistent with the ethnic com-
position of the white population in this state,
which consists predominantly of immigrants
from Italy, Portugal, and Spain where the fre-
quency of this mutation is lower than in the
rest of the world (70%).¹⁴⁻¹⁶ Our result for
 $\Delta F508$ differed significantly from that re-
ported by Raskin *et al*⁶ ($\chi^2_{(1)}=9.77$, $p<0.01$)
but is in accord with that reported by Martins
*et al*⁷ ($\chi^2_{(1)}=0.003$, $p>0.95$). The sample stud-
ied by Raskin *et al*⁶ might be composed
essentially of patients with more severe
phenotypes, which are correlated with the
 $\Delta F508$ mutation.

The frequencies of the next two most com-
mon mutations in the sample studied are also
consistent with the above ethnic composition.
The G542X mutation is fairly common
throughout Europe, particularly southern
Europe,¹⁷ and the N1303K mutation is com-
mon throughout Europe. Mutations G551D
and R553X have higher frequencies in north-
ern European countries, and for this reason
they may be rare in the population of the state
of São Paulo.

The most immediate application which
results from determination of the spectrum of
CF gene mutations in different populations is
genetic testing. In the sample of CF chromo-
somes that we studied, 42.5% of them could
be identified by the analysis of five mutations.
Reaching a 90% detection rate in this
population may require testing many addi-
tional low frequency mutations since São

Paulo, like the whole of Brazil, is genetically
very heterogeneous.

In conclusion, population molecular char-
acterisation is an important aspect in the
study of CF but, as expected for any
non-isolated population, it depends on the
analysis of a great number of mutations. Our
data should contribute further to the charac-
terisation of CF in the state of São Paulo.

We thank Maria del Pilar D Espinoza for her
participation in the clinical trial. This work was
supported by grants from CNPq and FAEP.

ENEIDA A PARIZOTTO
CARMEN S BERTUZZO

Departamento de Genética Médica, Faculdade de
Ciências Médicas, Universidade Estadual de
Campinas, CP 6111, 13083-970 Campinas, SP, Brazil

ANTONIO F RIBEIRO

Departamento de Pediatria, Faculdade de Ciências
Médicas, Universidade Estadual de Campinas, CP
6111, 13083-970 Campinas, SP, Brazil

- 1 Welsh MJ, Tsui LC, Boat TF, Beaudet AL.
Cystic fibrosis. In: Scriver CR, Beaudet AL, Sly
WS, Valle D, eds. *The metabolic and molecular
basis of inherited disease*. New York: McGraw-
Hill, 1995:3799-876.
- 2 Rommens JM, Iannuzzi MC, Kerem B, et al.
Identification of the cystic fibrosis gene:
chromosome walking and jumping. *Science*
1989;245:1059-65.
- 3 Riordan JR, Rommens JM, Kerem B, et al.
Identification of the cystic fibrosis gene:
cloning and characterization of complementary
DNA. *Science* 1989;245:1066-73.
- 4 Kerem B, Rommens JM, Buchanan JA, et al.
Identification of the cystic fibrosis gene: genetic
analysis. *Science* 1989;245:1073-80.
- 5 The CF Genetic Analysis Consortium. Popula-
tion variation of common cystic fibrosis mu-
tations. *Hum Mutat* 1994;4:167-77.
- 6 Raskin S, Phillips JA III, Krishnamani MR, et al.
DNA analysis of cystic fibrosis in Brazil by
direct PCR amplification from Guthrie cards.
Am J Med Genet 1993;46:665-9.
- 7 Martins CS, Ribeiro F, Costa FF. Frequency of
the cystic fibrosis delta F508 mutation in a
population from São Paulo State, Brazil. *Braz J
Med Biol Res* 1993;26:1037-40.
- 8 Kerem B, Zielenski J, Markiewicz D, et al. Iden-
tification of mutations in regions correspond-
ing to the two putative nucleotide (ATP)-
binding folds of the cystic fibrosis gene. *Proc
Natl Acad Sci USA* 1990;87:8447-51.
- 9 Osborne L, Knight R, Santis G, Hodson M. A
mutation in the second nucleotide binding fold
of the cystic fibrosis gene. *Am J Hum Genet*
1991;48:608-12.
- 10 Cutting GR, Kasch LM, Rosenstein BJ, et al. A
cluster of cystic fibrosis mutations in the first
nucleotide-binding fold of the cystic fibrosis
conductance regulator protein. *Nature* 1990;
346:366-9.
- 11 Saiki RK, Gelfand DH, Stoffel S, et al. Primer-
directed enzymatic amplification of DNA with
a thermostable DNA polymerase. *Science* 1989;
239:487-91.
- 12 Ng ISL, Pace R, Richard MV, et al. Methods for
analysis of multiple cystic fibrosis mutations.
Hum Genet 1991;87:613-17.
- 13 Rommens JM, Kerem B, Greer W, Chang P, Tsui
LC, Ray P. Rapid nonradioactive detection of
the major cystic fibrosis mutation. *Am J Hum
Genet* 1990;46:395-6.
- 14 Novelli G, Gasparini P, Savoia PF, Pignatti PF,
Sangiulio F, Dallapiccola B. Polymorphic
DNA haplotypes and delta F508 deletion in
212 Italian CF families. *Hum Genet* 1990;85:
420-1.
- 15 Duarte A, Barreto C, Marques-Pinto L, et al.
Cystic fibrosis in the Portuguese population:
haplotype distribution and molecular pathol-
ogy. *Hum Genet* 1990;85:404-5.
- 16 Chillón M, Nunes V, Casals T, et al. Distribu-
tion of the delta F508 mutation in 194 cystic
fibrosis families. *Hum Genet* 1990;85:396-7.
- 17 Nunes V, Gasparini P, Novelli G, et al. Analysis
of 14 cystic fibrosis mutations in five South
European populations. *Hum Genet* 1991;87:
737-8.