Original articles

Localisation of a gene causing endocrine neoplasia to a 4 cM region on chromosome 1p35-p36

C Williamson, A A J Pannett, J T Pang, C Wooding, M McCarthy, M N Sheppard, J Monson, R N Clayton, R V Thakker

Abstract

The development of some endocrine tumours, such as medullary thyroid carcinomas, pheochromocytomas, anterior pituitary adenomas, and parathyroid adenomas involve a putative tumour suppressor gene located on chromosome 1p32-pter, a region that represents 111 cM. In order to refine the location of this gene, 93 endocrine tumours (39 parathyroid adenomas, 40 anterior pituitary adenomas, seven pancreatic islet cell adenomas, and seven carcinoids) were investigated for loss of tumour heterozygosity (LOH) using the seven polymorphic loci 1pter - D1S228 - D1S507 - D1S234 - D1S476 - D1S220 - D1S207 - D1S206 - 1cen. LOH was detected in 27% of the parathyroid tumours and in 7.5% of the pituitary tumours, but in none of the pancreatic islet cell or carcinoid tumours. In addition, seven of the 10 parathyroid tumours that showed LOH of chromosome 1p facilitated a more precise mapping of this putative tumour suppressor gene; five tumours involved a loss only of the telomeric locus D1S228, whereas two other tumours showed LOH at the centromeric loci D1S507, D1S234, D1S476, and D1S220, but not D1S228. Thus, our results have mapped this tumour suppressor gene implicated in endocrine tumours to a 4 cM region flanked by D1S228 and D1S507 on chromosome 1p35-p36.

(*J Med Genet* 1997;34:617-619)

Keywords: endocrine tumours; loss of heterozygosity; chromosome 1p35-p36

The involvement of a tumour suppressor gene located on chromosome 1p in the aetiology of some endocrine tumours, for example, medullary thyroid carcinoma,¹⁻³ pheochromocytomas,^{2,3} anterior pituitary adenomas,⁴⁻⁶ and parathyroid adenomas,^{4,7} has been established by the demonstration of a loss of tumour heterozygosity (LOH). This gene has been mapped to 1p32-1pter,⁷ a region that represents 111 cM,⁸ and in order to facilitate positional cloning studies of this gene, we

undertook investigations to refine this location. In addition, we investigated the potential role of this gene on chromosome 1p in the aetiology of other endocrine tumours, such as pancreatic islet cell adenomas and carcinoid tumours, which may also occur in association with the multiple endocrine neoplasia type I (MENI) syndrome located on chromosome 11q13.⁹⁻¹¹

Materials and methods

TISSUE SPECIMENS

The 93 endocrine tumours (39 parathyroid tumours (30 from MENI patients and nine from non-MENI patients), seven pancreatic islet cell tumours (all from MENI patients), 40 anterior pituitary adenomas (all from non-MENI patients), and seven carcinoids (one intestinal from a MENI patient and six bronchial from non-MENI patients)) were obtained and stored at -70°C with a matching blood sample. The seven pancreatic islet cell tumours consisted of three insulinomas, three glucagonomas, and one non-functioning adenoma and the subtypes of the 40 anterior pituitary adenomas (14 somatotrophinomas, 15 non-functioning tumours, nine prolactinomas, and two corticotrophinomas) have been previously reported.^{6,12-14}

ALLELE LOSS STUDIES

Genomic DNA was extracted from tumours and leucocytes, as previously described.¹⁵ The seven microsatellite polymorphisms from chromosome 1p (D1S228, D1S507, D1S234, D1S476, D1S220, D1S207, and D1S206) were detected by the amplification of genomic DNA from each patient's paired tumour and leucocyte sample with specific oligonucleotide primers flanking the repeat sequence using the polymerase chain reaction (PCR) and LOH was scored as previously described.^{5,13,16} All the parathyroid, pancreatic islet cell, and carcinoid tumours were investigated at each of the seven chromosome 1p loci. However, owing to the limited quantities of DNA available from the pituitary tumours, the investigation of these tumours was restricted and only the polymorphisms at D1S228 and D1S507, which were found to define the critical interval from the parathyroid tumour data (fig 1), were investi-

MRC Molecular Endocrinology Group, MRC Clinical Sciences Centre, Royal Postgraduate Medical School, Du Cane Road, London W12 0NN, UK
C Williamson
A A J Pannett
J T Pang
C Wooding
R V Thakker

Department of Endocrinology, The Royal London Hospital, Whitechapel Road, London E1 1BB, UK
M McCarthy
J Monson

The Royal Brompton Heart Hospital, Sydney Street, London SW3 6NP, UK
M N Sheppard

Department of Medicine, North Staffordshire Royal Infirmary, Princes Road, Hartshill, Stoke-on-Trent ST4 7LN, UK
R N Clayton

Correspondence to: Professor Thakker.

Received 30 July 1996
Revised version accepted for publication 6 May 1997

Figure 1 Analysis of alleles on chromosome 1p in DNA from leucocytes and parathyroid tumours from seven patients. Patients 4, 5, and 7 suffered from MENI. The leucocyte and parathyroid tumour alleles were determined as shown in fig 2, and 35 of the 49 leucocyte genotypes determined were heterozygous. The corresponding parathyroid tumour genotypes indicated retention of heterozygosity (+) at 24 loci and allelic deletions (-) at 11 loci. The remaining 14 homozygous loci (black circles) in the leucocytes were not studied in the tumours, as the small amounts of DNA obtained from some tumours limited the loci which could be examined. Allelic deletions of chromosomes 11q13 and 13q are indicated (Y=yes, N=no, U=unknown), and those previously reported^{13, 14} are indicated by an asterisk. The results indicate that a putative tumour suppressor gene involved in the development of endocrine tumours is located in the interval flanked by D1S228 and D1S507 on chromosome 1p35-p36.

gated in all the pituitary tumours. An assessment of LOH at the chromosome 11q13 loci (D11S480, PYGM, D11S913, D11S1917, and D11S970) and the chromosome 13q12-q14 loci (Rb1.20 and D13S260) in the majority of these tumours has been previously reported,^{5 6 13 14} and tumours obtained since these studies were similarly characterised and the results pooled.

Results

Thirty-seven of the 39 patients with parathyroid tumours, 33 of the 40 patients with anterior pituitary tumours, all of the seven patients with pancreatic tumours, and all of the seven patients with carcinoid tumours were informative for the 1p loci that were examined. However, LOH was detected only in the parathyroid and anterior pituitary tumours and occurred in 27% and 7.5% of the respective tumours. Interestingly, six of the 28 (21%) MENI parathyroid tumours and four of the nine (44%) non-MENI parathyroid tumours showed chromosome 1p allele loss and this contrasted to the chromosome 11q13 allele loss in these tumours which were 80% (24 out

of 30) and 33% (three out of nine), respectively. Seven of the 37 (19%) parathyroid tumours (six MENI and one non-MENI) showed allele loss involving both chromosomes 1p and 11q13 (figs 1 and 2). The clinical and histological features, for example, age and sex of the patient and weight of the gland, of these patients were similar to those of patients who showed allele loss of only one chromosome. LOH involving chromosome 1p was observed in only three (two of 13 somatotrophinomas and one of eight prolactinomas) of the 40 anterior pituitary adenomas. Chromosome 1p allele loss was not observed in the pancreatic islet cell tumours or the carcinoid tumours. Seven of the 10 parathyroid tumours that showed LOH of chromosome 1p facilitated a more precise mapping of this putative tumour suppressor gene (fig 1). Thus, in five tumours allele loss involving only the telomeric locus D1S228 was observed whereas in two other tumours LOH at the centromeric loci D1S507, D1S234, D1S476, and D1S220, but not at D1S228, was observed. The combined results from these seven tumours located this putative tumour suppressor gene to an interval flanked by D1S228 and D1S507. The genetic distance between D1S228 and D1S507 has been estimated previously to be 4 cM,⁸ and thus our results have helped to define the map location of this gene causing endocrine neoplasia to chromosome 1p35-p36.

Discussion

Our results which show allele loss of chromosome 1p in 21% of MENI and 44% of non-MENI parathyroid adenomas are consistent with those of a previous study⁷ that located this putative tumour suppressor gene to chromosome 1p32-pter. Our observations of chromosome 1p allele loss in somatotrophinomas and prolactinomas are also in keeping with those previously reported in these subtypes and one corticotrophinoma.^{4, 6} In addition, our studies have defined a more precise location for this putative endocrine tumour suppressor gene to a 4 cM region, flanked by D1S228 and D1S507, in chromosome 1p35-p36. However, a previous report investigating chromosome 1p LOH in medullary thyroid carcinomas and phaeochromocytomas, which are associated with multiple endocrine neoplasia type 2, indicated that the location of a tumour suppressor gene in these tumours was likely to be in the vicinity of 1cen.¹⁷ Thus, it is possible that there may be two tumour suppressor genes located on chromosome 1p that are involved in the aetiology of some endocrine tumours. The absence of LOH involving chromosome 1p loci in the pancreatic islet cell and carcinoid tumours may either reflect a relatively low level of LOH, which would not have been detected in the small numbers of tumours studied, or a lack of any involvement of chromosome 1p tumour suppressor genes in the development of these tumours. Our definition of a 4 cM region located on chromosome 1p32-pter that is involved in the aetiology of some parathyroid tumours represents an important advance in identifying this putative tumour suppressor

Figure 2 Allelic deletions on chromosome 1p in six parathyroid tumours (patients 1, 7, and 8), a somatotrophinoma (patient 9), and a prolactinoma (patient 10), none of which has previously been reported.^{5,6} Patients 7 and 8 have MEN1. The panels show some of the autoradiographs obtained with the microsatellite polymorphisms at the D1S228, D1S507, D1S234, D1S476, and D11S913 loci. At each represented locus, the leucocytes (L) are heterozygous and the tumour (T) cells have either retained this heterozygosity (+) or developed a loss of heterozygosity (-). The combined LOH studies of chromosome 1p and 11q13 loci from the four parathyroid tumours of patient 8 show that T₁ and T₂ have allelic deletions of chromosome 1p and 11q13, whereas T₃ and T₄ have only an 11q13 loss. These results show that allelic deletions on chromosome 1p are involved in the development of parathyroid and pituitary tumours.

gene. A number of potential candidate genes are located in this region and these include the zinc finger ZFM60,¹⁸ the helix-loop-helix protein Heir-1,¹⁹ the developmental regulator gene Pax7,²⁰⁻²² the cell cycle regulator p58,²³ the transcription factor DAN that has homology to a mouse tumour suppressor gene.²⁵ Thus our studies, which have identified the region to search for such potential candidate genes, will help in further positional cloning studies aimed at identifying this tumour suppressor gene causing endocrine neoplasia.

This work was supported by the Medical Research Council (UK) (CW, AAJP, JTP, CW, and RVT). CW is an MRC Training Fellow and AAJP and JTP are MRC PhD students.

- 1 Mathew CG, Smith BA, Thorpe K, *et al.* Deletion of genes on chromosome 1 in endocrine neoplasia. *Nature* 1987;328:524-6.
- 2 Khosia S, Patel VM, Hay ID, *et al.* Loss of heterozygosity suggests multiple genetic alterations in pheochromocytomas and medullary thyroid carcinomas. *J Clin Invest* 1991; 87:1691-9.

- 3 Moley JF, Brother MB, Fong CT, *et al.* Consistent association of 1p loss of heterozygosity with pheochromocytomas from patients with multiple endocrine neoplasia type 2 syndromes. *Cancer Res* 1992;52:770-4.
- 4 Bystrom C, Larsson C, Blomberg C, *et al.* Localization of the MEN1 gene to a small region within chromosome 11q13 by deletion mapping in tumors. *Proc Natl Acad Sci USA* 1990;87:1968-72.
- 5 Thakker RV, Pook MA, Wooding C, Boscaro M, Scanarini M, Clayton RN. Association of somatotrophinomas with loss of alleles on chromosome 11 and with gsp mutations. *J Clin Invest* 1993;91:2815-21.
- 6 Boggild MD, Jenkinson S, Pistorello M, *et al.* Molecular genetic studies of sporadic pituitary tumors. *J Clin Endocrinol Metab* 1994;78:387-92.
- 7 Cryns VL, Yi SM, Tahara H, Gaz RD, Arnold A. Frequent loss of chromosome arm 1p DNA in parathyroid adenomas. *Genes Chrom Cancer* 1995;13:9-17.
- 8 Dracopoli NC, Bruns GA, Brodeur GM, *et al.* Report and abstracts of the First International Workshop of Human Chromosome 1 Mapping 1994, Bethesda, Maryland, 25-27 March 1994. *Cytogenet Cell Genet* 1994;67:144-65.
- 9 Larsson C, Skogseid B, Oberg K, Nakamura Y, Nordenskold M. Multiple endocrine neoplasia type 1 maps to chromosome 11 and is lost in insulinoma. *Nature* 1988;332:85-7.
- 10 Friedman E, Sakaguchi K, Bale AE, *et al.* Clonality of parathyroid tumors in familial multiple endocrine neoplasia type 1. *N Engl J Med* 1989;321:213-18.
- 11 Thakker RV, Bouloux P, Wooding C, *et al.* Association of parathyroid tumors in multiple endocrine neoplasia type 1 with loss of alleles on chromosome 11. *N Engl J Med* 1989; 321:218-24.
- 12 Thakker RV. The molecular genetics of the multiple endocrine neoplasia syndromes. *Clin Endocrinol (Oxf)* 1993;38:1-14.
- 13 Pang JT, Lloyd SE, Wooding C, *et al.* Genetic mapping studies of 40 loci and 23 cosmids in chromosome 11p13-11q13, and exclusion of μ -calpain as the multiple neoplasia type I gene. *Hum Genet* 1996;97:732-41.
- 14 Pearce SH, Trump D, Wooding C, Sheppard MN, Clayton RN, Thakker RV. Loss of heterozygosity studies at the retinoblastoma and breast cancer susceptibility (BRCA2) loci in pituitary, parathyroid, pancreatic and carcinoid tumours. *Clin Endocrinol (Oxf)* 1996;45:195-200.
- 15 Thakker RV, Davies KE, Whyte MP, Wooding C, O'Riordan JL. Mapping the gene causing X-linked recessive idiopathic hypoparathyroidism to Xq26-Xq27 by linkage studies. *J Clin Invest* 1990;86:40-5.
- 16 Scheinman SJ, Pook MA, Wooding C, Pang JT, Frymoyer PA, Thakker RV. Mapping the gene causing X-linked recessive nephrolithiasis to Xp11.22 by linkage studies. *J Clin Invest* 1993;91:2351-7.
- 17 Mulligan LM, Gardner E, Smith BA, Mathew CGP, Ponder BAJ. Genetic events in tumour initiation and progression in multiple endocrine neoplasia type 2. *Genes Chrom Cancer* 1993;6:166-77.
- 18 Lichter P, Bray P, Ried T, Dawid IB, Ward DC. Clustering of C2-H2 zinc finger motif sequences within telomeric and fragile site regions of human chromosomes. *Genomics* 1992;13:999-1007.
- 19 Ellmeier W, Aguzzi A, Kleiner E, Kurzbauer R, Weith A. Mutually exclusive expression of a helix-loop-helix gene and N-myc in human neuroblastomas and in normal development. *EMBO J* 1992;11:2563-71.
- 20 Schafer BW, Mattei MG. The human paired domain gene PAX7 (Hup1) maps to chromosome 1p35-1p36.2. *Genomics* 1993;17:249-51.
- 21 Shapiro DN, Sublett JE, Li B, Valentine MB, Morris SW, Noll M. The gene for PAX7, a member of the paired-box-containing genes, is localized on human chromosome arm 1p36. *Genomics* 1993;17:767-9.
- 22 Stapleton P, Weith A, Urbanek P, Kozmik Z, Busslinger M. Chromosomal localization of seven PAX genes and cloning of a novel family member, PAX-9. *Nat Genet* 1993;3:292-8.
- 23 Eipers PG, Barnoski BL, Han J, Carroll AJ, Kidd VJ. Localization of the expressed human p58 protein kinase chromosomal gene to chromosome 1p36 and a highly related sequence to chromosome 15. *Genomics* 1991;11: 621-9.
- 24 Lees JA, Saito, Vidal M, *et al.* The retinoblastoma protein binds to a family of E2F transcription factors. *Mol Cell Biol* 1993;13:7813-25.
- 25 Enomoto H, Ozaki T, Takahashi E, *et al.* Identification of human DAN gene, mapping to the putative neuroblastoma tumor suppressor locus. *Oncogene* 1994;9:2785-91.