Linkage analysis in Usher syndrome type I (USH1) families from Spain

Carmen Espinós, Carmen Nájera, José Maria Millán, Carmen Ayuso, Montserrat Baiget, Herminio Pérez-Garrigues, Onofre Rodrigo, Concepción Vilela, Magdalena Beneyto

Departamento de Genética, Fac C C Biológicas, Universidad de Valencia, C/ Dr Moliner 50, 46100 Burjassot, Valencia, Spain
C Espinós
C Nájera

Unidad de Genética y Diagnóstico Prenatal, Hospital La Fe, Valencia, Spain
J M Millán
M Beneyto

Departamento de Genética, Fundación Jiménez Díaz, Madrid, Spain
C Ayuso

Unitat de Genética, Hospital de la Santa Creu i Sant Pau, Barcelona, Spain
M Baiget

Servicio de Otorrinolaringología, Hospital La Fe, Valencia, Spain
H Pérez-Garrigues
O Rodrigo

Unidad de Neurofisiología, Hospital La Fe, Valencia, Spain
C Vilela

Correspondence to: Professor Nájera.

Received 21 January 1997
Revised version accepted for publication 22 October 1997

Table 1 Characteristics of polymorphic markers on chromosome 11q used in the present study

Locus	Primers	Alleles	Size (bp)	Annealing temp	References
ROMI*	ROM1/2			55°C	26
INT2	FGF3	9	161-177	55°C	27
D11S916	AFM185yal	9	135-153	60°C	29
D11S527	JG4-A/JG4-C	12	146-166	55°C	30
OMP	OMP	4	244-254	55°C	28
D11S911	AFM155xh10	13	159-203	50°C	29
D11S35	780/781	6	152-162	40°C	31
D11S528	42026/42027	6	188-208	50°C	32

*ROM1/2 marker was studied by means of SSCP.

Table 2 Characteristics of microsatellite polymorphisms on chromosome 3q used in the present study

Locus	Primers	Alleles	Size (bp)	Annealing temp	References
RHO	RHO	4	120	57°C	33
D3S1555	AFM056xf12	8	221-237	55°C	29
D3S1308	AFM240ve1	5	104-112	50°C	29
D3S3705	AFMa043tf5	5	228-236	55°C	29
D3S1299	AFM220yh12	6	218-228	53°C	29
D3S1279	AFM164xg9	10	264-282	55°C	29

Abstract

Usher syndrome (USH) is an autosomal recessive hereditary disorder characterised by congenital sensorineural hearing loss and gradual visual impairment secondary to retinitis pigmentosa (RP). The disorder is clinically and genetically heterogeneous. With regard to Usher type I (USH1), several subtypes have been described, the most frequent being USH1B located on chromosome 11q13.5. Of 18 USH1 families studied by linkage analysis, 12 (67%) showed significant lod score values for locus D11S527 ($Z_{\max}=14.032$, $\theta=0.000$) situated on chromosome 11q. Our findings suggest considerable genetic heterogeneity in the Spanish USH1 population. It is important to note that one of our families linked to the USH1B locus shows interesting intrafamilial clinical variability. As regards the remaining six USH1 families, the linkage analysis did not provide conclusive data, although two of them show slight linkage to markers located on chromosome 3q ($Z_{\max}=1.880$, $\theta=0.000$ for D3S1279), the same location that had previously been assigned to some USH3 families.

(J Med Genet 1998;35:391-398)

Keywords: Usher syndrome; linkage analysis; genetic heterogeneity

Usher syndrome (USH), which is recognised as the most common cause of combined deafness and blindness in adults, is an autosomal

recessive hereditary disorder characterised by congenital sensorineural hearing loss and gradual visual impairment secondary to retinitis pigmentosa (RP).^{1,2}

The disorder is clinically heterogeneous and, at present, three types can be distinguished on the basis of clinical findings principally related to audiological and vestibular parameters.³⁻⁷ Usher type I (USH1) patients suffer severe to profound congenital hearing loss and abnormal or absent vestibular response, whereas Usher type II (USH2) is characterised by moderate to severe congenital hearing loss and normal vestibular function. Usher type III (USH3), which is clinically similar to USH2, is characterised by the progressive nature of its hearing loss and by the mild vestibular dysfunction. Although retinitis pigmentosa has been accepted as occurring earliest in type I, the ocular data are insufficient for distinguishing between types. It is interesting to note that a number of heterozygous carriers have shown significant audiological and ophthalmological abnormalities.^{8,9}

Genetic heterogeneity for USH1 and USH2 has been verified by linkage analysis. At least six different loci have been located: USH1A on 14q,^{10,11} USH1B on 11q,^{12,13} USH1C on 11p,^{14,15} USH1D on 10q,¹⁶ USH1E on 21q21,¹⁷ and USH2A on 1q.^{18,19} There are, however, families of both types which fail to show linkage to any of these regions.^{11,17} As far as USH3 is concerned, a notable clustering has been observed in eastern Finland. Sankila *et al*²⁰ assigned an USH3 locus to chromosome 3q. Furthermore, some USH2 families have been linked to the USH3 locus.²¹ To date, only one gene related to USH is known, myosin VIIA, an unconventional myosin, involved in the USH1B phenotype.²²⁻²⁴

In the present study, the results of linkage analysis in 18 Spanish USH1 families are reported.

Material and methods

FAMILIES AND CLINICAL STUDIES

Eighteen Spanish families clinically diagnosed as having USH1 with a total of 34 affected patients (23 males and 11 females) have been studied. Most of these families have multiple affected children and four show consanguinity. Patient II.4 of family 11 (fig 1) shows peculiar clinical features, since he suffers prelingual profound deafness and vestibular dysfunction; however, to date (39 years old), he has not shown any symptoms of RP. Moreover, he is also severely mentally retarded. In the remaining patients, mental retardation or physical dysmorphism were not noted.

Figure 1 Pedigrees and haplotypes of the 12 Usher syndrome type I families linked to 11q. Haplotypes were constructed based on the minimum number of recombinations between these markers.

Clinical tests were performed on all affected subjects. The ophthalmological studies included determination of visual acuity, fundus ophthalmoscopy, visual field examination, electroretinography, and visually evoked potentials. The diagnostic criteria used to recognise RP were night blindness, loss of visual field and acuity, and altered or unrecordable electroretinogram. The otolaryngological evalua-

tion included audiological examination, neurological exploration, electronystagmography, and posturography.

DNA STUDIES

DNA was extracted from peripheral blood leucocytes from each person.²⁵ PCR amplifications were performed in a total volume of 50 µl containing 200 ng of human genomic DNA, 50

Table 3 Lod scores of chromosome 11 markers versus Usher type I in the 12 linked families

	Recombination (θ)						
	0.000	0.010	0.05	0.10	0.20	0.30	0.40
ROMI	—∞	2.440	3.356	3.355	2.732	1.823	0.789
INT2	—∞	4.344	5.902	5.953	4.788	3.105	1.314
D11S916	—∞	1.522	4.784	3.970	4.725	3.192	1.369
D11S527	14.032	13.998	12.770	11.224	8.120	5.008	2.001
OMP	7.259	7.114	6.520	5.755	4.168	2.556	1.000
D11S911	13.524	13.253	12.151	10.740	7.845	4.883	1.970
D11S35	—∞	—8.456	—4.792	0.459	1.508	1.122	0.511
D11S528	—∞	—0.909	1.931	2.606	2.418	1.611	0.653

Figure 2 Genetic distances (KcM: Kosambi centimorgans) and locus order of the 11q markers used in the study. Data were obtained from Genome Data Bank (GDB) and van Heyningen and Little.³⁷

mmol/l KCl, 10 mmol/l Tris-HCl, pH 8.3, 1.5 mmol/l MgCl₂, 100 µg/ml gelatin, 200 µmol/l dNTP, 1 µmol/l of each primer (50 pm), and 1 unit of *Taq* DNA polymerase. The polymorphic microsatellite markers and their PCR conditions are given in tables 1 and 2.²⁶⁻³³ The amplified DNA was electrophoresed on a 8-12% polyacrylamide gel and stained with silver. The marker ROM1/2 was analysed by SSCP with electrophoresis on a polyacrylamide gel containing 5% glycerol.

Figure 3 Genetic distances (KcM: Kosambi centimorgans) and locus order of the 3q markers used. Data were obtained from Genome Data Bank (GDB) and Sankila et al.²⁰

LINKAGE ANALYSIS
All the markers used in the present study were microsatellite markers, except for ROM1/2 which detects a polymorphism within intron 1 of the ROMI gene.^{34 35} Pairwise and multipoint linkage analysis was performed using the MLINK and LINKMAP subroutines of version 5.1 of the LINKAGE program.³⁶ The linear order of markers on chromosomes 11q³⁷ and 3q,²⁰ and distances between them, are shown in figs 2 and 3. Penetrance was assumed to be 0.99 and the mode of inheritance was assumed to be autosomal recessive.

Results
Haplotype analysis (fig 1) suggested that in 12 of the 18 families (67%), linkage for the USH1B locus existed. Pairwise lod scores for chromosome 11 marker loci are presented in table 3. Taking all 12 families together, pairwise linkage analysis gives lod scores of 14.032, 7.259, and 13.524 for D11S527, OMP, and D11S911 respectively, at zero recombination distance.

Table 4 Lod scores of chromosome 3 markers versus Usher type I in the two linked families

	Recombination (θ)						
	0.000	0.010	0.05	0.10	0.20	0.30	0.40
RHO	1.454	1.428	1.318	1.172	0.847	0.492	0.158
D3S1555	0.125	0.116	0.086	0.056	0.018	0.004	0.000
D3S1308	1.153	1.127	1.020	0.883	0.599	0.320	0.245
D3S3705	0.505	0.487	0.416	0.332	0.185	0.079	0.019
D3S1299	0.852	0.830	0.742	0.630	0.408	0.206	0.056
D3S1279	1.880	1.838	1.669	1.450	0.992	0.536	0.157

Figure 4 Pedigrees and haplotypes of the two Usher syndrome type I families linked to 3q. Haplotypes were constructed based on the minimum number of recombinations between these markers.

Family 15

Figure 5 Pedigrees and haplotypes of the families unlinked to any known Usher locus. Haplotypes were constructed based on the minimum number of recombinations between these markers.

Family 11 shows linkage to markers placed on chromosome 11q if we assume that patient II.4 is affected with USH1. If only the two females are considered as affected, this family fails to show linkage to any of the USH loci described.

In the 12 families linked to 11q, none of the alleles that closely flank USH1B displayed significant differences between gene frequencies

of the parental USH1B chromosomes and the normal parental chromosomes, which indicates that linkage disequilibrium is not present in this Spanish sample.

The remaining six USH1 families were studied with markers located in other Usher candidate regions: 14q32, 11p15.1, 10q, 21q21, 3q21-q25, and 1q42. For two of the families (fig 4), slightly positive lod scores were

Figure 5 (cont).

observed for five markers located on chromosome 3q ($Z_{\max}=1.880$, $\theta=0.000$ for D3S1279) (table 4). For the remaining four USH1 families (fig 5), no conclusive evidence of linkage to any loci previously described was obtained.

With respect to family 15, which includes three sibs with clinical symptoms of USH1, it is important to note that the mother suffers profound bilateral sensorineural deafness, difficult but intelligible speech, normal vestibular response, and absence of RP. The father had no ocular or otological symptoms. The dead grandmother, who was not investigated, also had a hearing impairment of unknown degree. Taking this family as USH1, segregation of the disease with markers of the loci described was not observed.

Discussion

Even though a high percentage of the 18 USH1 Spanish families studied are linked to chromo-

some 11q (67%), there are an important number of cases (33%) that fail to show linkage to either 11q or other known regions for USH1 (11p, 14q, 10q, or 21q). The results indicate considerable genetic heterogeneity in these USH1 families. For the 11q linked cases, lod scores obtained here are in agreement with those previously reported elsewhere with respect to the physical mapping and localisation of USH1B. These new data, thus, place the candidate gene in the region between D11S527 and D11S911, which is compatible with other reports^{12 13} and which, although we have not yet analysed mutations in the myosin gene, supports the role of this gene in the pathogenesis of USH1B families.^{22-24 38 39} These results also discount the possible involvement in the USH1 syndrome of other genes formerly considered candidates, such as ROMI and INT2,^{12 35} since recombinant events have been found with these loci.

Figure 5 (cont).

It is important to emphasise that at least two of the USH1 families in this study show positive lod scores for markers located on chromosome 3q. This chromosomal region is exactly the same as that in which Sankila *et al*²⁰ and, to a lesser extent, Kimberling *et al*²¹ reported linkage for USH3 families from Finland and for some Swedish USH2 families, respectively. To date, families classified as USH1 have not been found to be related to this locus, although the possibility that some USH3

patients show a phenotype similar to USH1 or USH2 has already been suggested.⁴⁰ Lod scores are not high, but they contrast with the negative results for markers located on chromosome 3q shown by the other USH1 families linked to 11q or unlinked to the other loci described, and also with the data obtained for markers placed in other candidate regions in these two families.

Vestibular function is a reliable and consistent discriminator between Usher types I and II

since most, if not all, type I patients lack vestibular function.⁴¹ There are, nevertheless, cases of USH2 and USH3 in which some abnormal vestibular response is reported.^{6, 42} In this study, families 1 to 18 could not be diagnosed as USH2 or USH3 since they had severely abnormal vestibular function and profound congenital prelingual hearing loss leading to total deaf-mutism. Similarly, the onset of RP is not conclusive for differentiation of the three types.⁴³ Therefore, according to the present results and to others which are similar,^{20, 21} families classified as USH1, USH2, and USH3 have all shown linkage to 3q. It would be interesting to re-examine the clinical parameters, owing to the wide range of clinical symptoms in every Usher type, and to perform genotype-phenotype correlations in order to discriminate between the different types and to investigate whether some of them may be the result of allelic manifestations of a single gene.

It is important to stress the special clinical aspects of family 11, which has in the same sibship two affected females who show a typical USH1 phenotype and one male who suffers profound hearing impairment and absence of RP. This family has been considered as USH1, mainly because of linkage to 11q markers. Moreover, since allelic variants of the myosin VIIA gene may be involved in both USH1B syndrome and an isolated form of deafness (DFNB2),^{44, 45} the possibility that a mutation in this gene is responsible for the phenotypic variability within this family cannot be discarded. It is also interesting to note that the same genomic region is involved in a dominant form of non-syndromic deafness (DFNA11).⁴⁶ This phenomenon has already been observed in other chromosome regions (DFNA3 and DFNB1 to 13q11,^{47, 48} DFNA9 and DFNB5 to 14q12⁴⁹).

Family 15 may represent a similar example to family 11, since it also presents intrafamilial clinical variability if we consider members of different generations. This family remains unlinked to any Usher locus described to date. It could be possible that one mutation in an unknown gene is responsible for syndromic and non-syndromic forms of hearing loss within this family. If this supposition were correct, each family would represent an example of a single mutation responsible for syndromic and isolated forms of deafness.

The authors would like to thank the patients and their family members for their help and cooperation. We acknowledge the Asociación de retinosis pigmentaria de la Comunidad Valenciana, the FAARPEE, ONCE, Fundación ONCE, and the Fondo de Investigaciones Sanitarias (FISS No 95/1814) for financial support.

- 1 Usher CH. Bowman's lecture: on a few hereditary eye affections. *Trans Ophthalm Soc UK* 1935;55:164-245.
- 2 Hallgren B. Retinitis pigmentosa combined with congenital deafness; with vestibulo-cerebellar ataxia and mental abnormality in a proportion of cases. A clinical and genetic-statistical study. *Acta Psychiatr Scand* 1959;34(suppl 138):5-101.
- 3 Kimberling WJ, Möller CG, Davenport SLH, et al. Usher syndrome: clinical findings and gene localization studies. *Laryngoscope* 1989;99:66-72.
- 4 Tamayo ML, Bernal JE, Tamayo GE, et al. Usher syndrome: results of a screening program in Colombia. *Clin Genet* 1991;40:304-11.
- 5 van Aarem A, Cremers CWRJ, Pinckers AJLG, Huygen PLM, Hombergen GCJH, Kimberling BJ. The Usher syn-

- drome type 2A: clinical findings in obligate carriers. *Int J Pediatr Otorhinolaryngol* 1995;31:159-74.
- 6 Gorlin RJ, Tilsner TJ, Feinstein S, Duvall AJ. Usher's syndrome type III. *Arch Otolaryngol* 1979;105:353-4.
- 7 Karjalainen S, Pakarinen L, Terasvirta M, Kaariainen H, Vartiainen E. Progressive hearing loss in Usher's syndrome. *Ann Otol Rhinol Laryngol* 1989;98:863-6.
- 8 Karjalainen S, Terasvirta M, Kärjä J, Kääriäinen H. Usher's syndrome type III: ENG findings in four affected and six unaffected siblings. *J Laryngol Otol* 1985;99:43-8.
- 9 Wagenaar M, ter Rahe B, van Aarem A, et al. Clinical findings in obligate carriers of type I Usher syndrome. *Am J Med Genet* 1995;59:375-9.
- 10 Kaplan J, Gerber S, Bonneau D, et al. A gene for Usher syndrome type I (USH1A) maps to chromosome 14q. *Genomics* 1992;14:979-87.
- 11 Larget-Piet D, Gerber S, Bonneau D, et al. Genetic heterogeneity of Usher syndrome type I in French families. *Genomics* 1994;21:138-43.
- 12 Kimberling WJ, Weston MD, Möller CG, et al. Linkage of Usher syndrome type I gene (USH1B) to the long arm of chromosome 11. *Genomics* 1992;14:988-94.
- 13 Bonné-Tamir B, Korostishevsky M, Kalinsky H, et al. Genetic mapping of the gene for Usher syndrome: linkage analysis in a large Samaritan kindred. *Genomics* 1994;20:36-42.
- 14 Keats BJB, Nouri N, Pelias MZ, Deininger PL, Litt M. Tightly linked flanking microsatellite markers for the Usher syndrome type I locus on the short arm of chromosome 11. *Am J Hum Genet* 1994;54:681-6.
- 15 Smith RJH, Lee EC, Kimberling WJ, et al. Localization of two genes for Usher syndrome type I to chromosome 11. *Genomics* 1992;14:995-1002.
- 16 Wayne S, Der Kaloustian VM, Schloss M, et al. Localization of Usher syndrome type ID gene (USH1D) to chromosome 10. *Hum Mol Genet* 1996;5:1689-92.
- 17 Chaib H, Kaplan J, Gerber S, et al. A newly identified locus for Usher syndrome type I, USH1E, maps to chromosome 21q21. *Hum Mol Genet* 1997;6:27-31.
- 18 Kimberling WJ, Weston MD, Möller CG, et al. Localization of Usher syndrome type II to chromosome 1q. *Genomics* 1990;7:245-9.
- 19 Lewis RA, Otterud B, Stauffer D, Lalouel JM, Leppert M. Mapping recessive ophthalmic diseases: linkage of the locus for Usher syndrome type II to a DNA marker on chromosome 1q. *Genomics* 1990;7:250-6.
- 20 Sankila EM, Pakarinen L, Kääriäinen H, et al. Assignment of an Usher syndrome type III (USH3) gene to chromosome 3q. *Hum Mol Genet* 1995;4:93-8.
- 21 Kimberling WJ, Weston MD, Möller C, et al. Gene mapping of Usher syndrome type IIa: localization of the gene to a 2.1-cM segment on chromosome 1q41. *Am J Hum Genet* 1995;56:216-23.
- 22 Weil D, Blanchard S, Kaplan J, et al. Defective myosin VIIA gene responsible for Usher syndrome type 1B. *Nature* 1995;374:60-1.
- 23 Weston MD, Kelley PM, Overbeck LD, et al. Myosin VIIA mutation screening in 189 Usher syndrome type I patients. *Am J Hum Genet* 1996;59:1074-83.
- 24 Levy G, Levi-Acobas F, Blanchard S, et al. Myosin VIIA gene: heterogeneity of the mutations responsible for Usher syndrome type IB. *Hum Mol Genet* 1997;6:111-16.
- 25 Kunkel LM, Tantravah U, Eisenhard M, Latt SA. Regional localization on the human X of DNA sequences cloned from flow sorted chromosomes. *Nucleic Acids Res* 1982;10:1557-71.
- 26 Nichols BE, Bascom R, Litt M, McInnes R, Sheffield VC, Stone EM. Refining the locus for Best vitelliform macular dystrophy and mutation analysis of the candidate gene ROM1. *Am J Hum Genet* 1994;54:95-103.
- 27 Polymeropoulos MH, Xiao H, Rath DS, Merrill CR. Dinucleotide repeat polymorphism at the int-2 proto-oncogene locus. *Nucleic Acids Res* 1990;18:7468.
- 28 Piekke-Dahl S, Weston MD, Overbeck LD, Evans KL, Porteous DJ, Kimberling WJ. Dinucleotide repeat polymorphism at the human olfactory marker protein (OMP) locus on chromosome 11q13.5 near tyrosinase (TYR). *Hum Mol Genet* 1993;2:822.
- 29 Gyapay G, Morissette J, Vignal A, et al. The 1993-94 Génethon human genetic linkage map. *Nat Genet* 1994;7:246-339.
- 30 Brown DL, Gault J, Thompson MB, Hauge XY, Evans GA, Litt M. Dinucleotide repeat polymorphisms at the D11S527 locus. *Nucleic Acids Res* 1991;19:4790.
- 31 Litt M, Sharman V, Luty JA. Dinucleotide repeat polymorphism at the D11S35 locus. *Nucleic Acids Res* 1990;18:5921.
- 32 Hauge XY, Evans GA, Litt M. Dinucleotide repeat polymorphisms at the D11S528 locus. *Nucleic Acids Res* 1991;19:1964.
- 33 Weber JL, May PE. Abundant class of human DNA polymorphisms which can be typed using the polymerase chain reaction. *Am J Hum Genet* 1989;44:388-96.
- 34 Bascom RA, Garcia-Heras J, Hsieh GL, et al. Localization of the photoreceptor gene ROM1 to human chromosome 11 and mouse chromosome 19: sublocalization to human 11q13 between PGA and PYGM. *Am J Hum Genet* 1992;51:1028-35.
- 35 Bascom RA, Liu L, Humphries P, Fishman GA, Murray JC, McInnes RR. Polymorphisms and rare sequence variants at the ROM1 locus. *Hum Mol Genet* 1993;2:1975-7.
- 36 Lathrop GM, Lalouel JM, Julier C, Ott J. Multilocus linkage analysis in humans: detection of linkage and estimation of recombinations. *Am J Hum Genet* 1985;37:482-98.

- 37 van Heyningen V, Little PFR. Report of the fourth international workshop on human chromosome 11 mapping 1994. *Cytogenet Cell Genet* 1995;69:128-58.
- 38 Hasson T, Heintzelman MB, Santos-Sacchi J, Corey DP, Mooseker MS. Expression in cochlea and retina of myosin VIIa, the gene product defective in Usher syndrome type 1B. *Proc Natl Acad Sci USA* 1995;92:9815-19.
- 39 El-Amraoui A, Sahly I, Picaud S, Sahed J, Abitbol M, Petit C. Human Usher 1B/mouse *shaker-1*: the retinal phenotype discrepancy explained by the presence/absence of myosin VIIA in the photoreceptor cells. *Hum Mol Genet* 1996;5:1171-8.
- 40 Pakarinen L, Sankila EM, Tuppurainen K, Karjalainen S, Kääriäinen H. Usher syndrome type III (USH3): the clinical manifestations in 42 patients. *Scand J Log Phon* 1995;20:141-50.
- 41 Kimberling WJ, Möller C. Clinical and molecular genetics of Usher syndrome. *J Am Acad Audiol* 1995;6:63-72.
- 42 Smith RJH, Berlin CI, Hejtmancik JF, et al. Clinical diagnosis of the Usher syndromes. *Am J Med Genet* 1994;50:32-8.
- 43 Pakarinen L, Tuppurainen K, Laippala S, Mäntyjärvi M, Puhakka H. The ophthalmological course of Usher syndrome type III. *Int Ophthalmol* 1996;19:307-11.
- 44 Liu XZ, Walsh J, Mburu P, et al. Mutations in the myosin VIIA gene cause non-syndromic recessive deafness. *Nat Genet* 1997;16:188-90.
- 45 Weil D, Küssel P, Blanchard S, et al. The autosomal recessive isolated deafness, DFNB2, and the Usher 1b syndrome are allelic defects of the myosin-VIIA gene. *Nat Genet* 1997;16:1991-3.
- 46 Tamagawa Y, Kitamura K, Ishida T, et al. A gene for a dominant form of non-syndromic sensorineural deafness (DFNA11) maps within the region containing the DFNB2 recessive deafness gene. *Hum Mol Genet* 1996;5:849-52.
- 47 Chaib H, Lina-Granade G, Guilford P, et al. A gene responsible for a dominant form of neurosensory non-syndromic deafness maps to the NSRD1 recessive deafness gene interval. *Hum Mol Genet* 1994;3:2219-22.
- 48 Kelsell DP, Dunlop J, Stevens HP, et al. Connexin 26 mutations in hereditary non-syndromic sensorineural deafness. *Nature* 1997;387:80-3.
- 49 Manolis EN, Yandavi N, Nadol JB Jr, et al. A gene for non-syndromic autosomal dominant progressive postlingual sensorineural hearing loss maps to chromosome 14q12-12. *Hum Mol Genet* 1996;5:1047-50.