CHRIS F INGLEHEARN
EMMA E TARTTELIN
T JEFFERY KEEN
Molecular Medicine Unit, St James's University
Hospital, Leeds, UK

SHOMI S BHATTACHARYA
Department of Molecular Genetics, Institute of
Ophthalmology, London, UK

ANTHONY T MOORE
Ophthalmology Department, Addenbrooke's Hospital,
Cambridge, UK

RACHEL TAYLOR
ALAN C BIRD
Department of Clinical Ophthalmology, Institute of
Ophthalmology, London, UK

- 1 Boughman JA, Conneally PM, Nance WE. Population genetic studies of retinitis pigmentosa. *Am J Hum Genet* 1980;32:223-35.
- 2 Inglehearn CF. Molecular genetics of inherited retinal dystrophies. *Eye* 1998;12:571-9.
- 3 Sung CH, Davenport CM, Hennessey JC, et al. Rhodopsin mutations in autosomal dominant retinitis pigmentosa. *Proc Natl Acad Sci USA* 1991;88:6481-5.
- 4 Inglehearn CF, Tarttelin EE, Plant C, et al. A genetic survey of twenty dominant retinitis pigmentosa families: frequencies of the eight known loci and evidence for further heterogeneity. *J Med Genet* 1998;35:1-5.
- 5 Souied EH, Rozet JM, Gerber S, et al. Screening for mutations within the rhodopsin, peripherin-RDS and ROM1 genes in autosomal dominant retinitis pigmentosa (ADRP) in pedigrees from France. *Invest Ophthalmol Vis Sci* 1995;36(suppl):890.
- 6 Xu S, Schwartz M, Rosenberg T, Gal A. A ninth locus (RP18) for autosomal dominant retinitis pigmentosa maps in the pericentric region of chromosome 1. *Hum Mol Genet* 1996;8:1193-7.
- 7 Xu S, Rosenberg T, Gal A. Refined genetic mapping of autosomal dominant retinitis pigmentosa locus RP18 reduces the critical region to 2 cM between D1S442 and D1S2858 on chromosome 1q. *Hum Genet* 1998;102:493-4.
- 8 Lyness AL, Ernst W, Quinlan MP, et al. A clinical, psychophysical and electroretinographic survey of patients with autosomal dominant retinitis pigmentosa. *Br J Ophthalmol* 1985;69:326-39.

Frequency of inherited deletions of 22q11

We read with interest the report by Ryan *et al* on the spectrum of clinical features associated with 22q11 deletions, in which they aim "to increase the clinical information on which management and counselling can be based". They report a frequency of inherited deletions of 28% in their study. We have looked at the frequency of familial cases of 22q11 deletions in patients from all of Wales, none of whom was included in the European collaborative study, and have found a much lower frequency.

Since starting investigations for 22q11 deletions, it has been our policy to test both parents of all deletion cases irrespective of the reason for referral. Fifty cases with a deletion have now been detected, and these have been referred by clinicians from a range of disciplines, including medical genetics, paediatrics, cardiology, psychiatry, and nephrology. In 41 cases both parents were studied and in two cases only maternal blood was available. Most of the families who have not been studied are recent referrals. The deletion was inherited in only four out of the 41 cases, giving a frequency of approximately 10%. The four cases of parental transmission were all maternal, and all parents with the microdeletion showed facial features associated with 22q11 deletion. One patient had a corrected double outlet right ventricle, an-

other had a ventral septal defect which closed spontaneously in childhood.

The frequency of familial transmission in our group of patients of 10% is nearer the figure of 8% reported by Driscoll *et al*¹ in the only other large study than the 28% reported by the European collaborative study.¹ It is difficult to explain the large discrepancy between these figures; however, we would stress that where possible we follow up all cases irrespective of parental phenotypes, and although the collaborative study is a far larger one, the authors recognise that patient selection for testing because of suggestive clinical features may have inflated their frequency. If an assumption is made that no further deletions were present in the parents not tested by Ryan *et al*,¹ then the minimal estimate of inherited deletion from their study is 81/558 or 15%, a value closer to our frequency.

An accurate figure for parental transmission of 22q11 microdeletions is important when counselling parents of a child recently diagnosed, helping to allay anxiety, guilt, and also fears of recurrence. A risk of 28% coming as it does from such a comprehensive and well respected study is the one most likely to be used for counselling purposes; however, in view of our findings and those of Driscoll *et al*,² we feel this may be too high and that more clinical data on non-selected patients are needed before a true frequency of familial transmission is known.

P W THOMPSON
S J DAVIES

Institute of Medical Genetics, University Hospital of
Wales, Heath Park, Cardiff CF4 4XW, UK

- 1 Ryan AK, Goodship JA, Wilson DI, et al. Spectrum of clinical features associated with interstitial chromosome 22q11 deletions: a European collaborative study. *J Med Genet* 1997;34:798-804.
- 2 Driscoll DA, Salvin J, Sellinger B, et al. Prevalence of 22q11 microdeletions in DiGeorge and velocardiofacial syndromes: implications for genetic counselling and prenatal diagnosis. *J Med Genet* 1993;30:813-817.

The annual incidence of DiGeorge/velocardiofacial syndrome

The majority of cases with either DiGeorge syndrome or velocardiofacial syndrome are caused by a submicroscopic deletion in chromosome 22q11 (del22q11).¹⁻³ Since the advent of a routine diagnostic test for this microdeletion, the number of patients diagnosed has increased dramatically, including many patients with either mild features or non-specific presenting symptoms. The del22q11 occurs much more frequently than previously thought, but precise incidence figures are not known. Wilson *et al*¹ found a del22q11 in approximately 5% of children with a congenital heart defect (CHD), and therefore an estimated incidence of at least 1/4000 live births. A more direct way is to

determine the annual incidence of cases with a del22q11, in a well defined region, with a known number of births. In a region in southern France, with an annual birth rate of approximately 23 000, Du Montcel *et al*⁴ found 1/9700 as a minimum incidence of the del22q11 associated with the typical clinical picture.

In the Flemish region of Belgium, all genetic tests are performed in four genetic centres. The number of births was extracted from the annual reports of the Study Centre for Perinatal Epidemiology (Studiecentrum voor perinatale epidemiologie, SPE), which registers in Flanders over 99% of all live and stillborn children with a birth weight of over 500 g. Routine genetic testing for a del22q11 by means of FISH became available during the years 1992-1993. A total of 151 Flemish cases have been diagnosed since and, of these, 94 were born before 1992, six in 1997, and 51 in the five year period between January 1992 and December 1996. The annual birth rate in Flanders ranges from 68 613 in 1992 to 63 550 in 1996, with a total of 326 166 births during the five year study period of 1992-1996. Therefore, the estimated annual incidence of a del22q11 in 1992-1996 is 15.3/100 000 newborns (95% confidence intervals 13.3-17.2), or 1/6395 (table 1).

However, as in the study of Du Montcel *et al*,⁴ it is evident that this represents a minimum estimate, since many cases with mild features remain undiagnosed. In the total group of patients with a del22q11, 81 of the 151 patients (54%) have a symptomatic congenital heart defect (CHD), compared to 37 of the 51 patients (72.5%) born during the last five years. This confirms the clinical experience that the diagnosis is delayed in patients without a heart defect. In our series of patients from 1992-1996, mean age at diagnosis of those with a heart defect was 8.3 months (range: day of birth-39 months). During the last three years, the majority of infants with a conotruncal heart defect and a del22q11 were diagnosed during the first weeks of life. In contrast, patients without a heart defect were diagnosed at a mean age of 25.9 months (SD 17 months), when developmental delay or speech delay becomes evident. It can be estimated, therefore, that a large proportion of the children born during the last five years with a del22q11 but without a heart defect remain to be diagnosed.

Taken together, our observation is in good agreement with a maximal annual incidence of 1/4500 found in children born in 1993 in the study of Du Montcel *et al*⁴ and with an estimated incidence of 1/4000 as suggested by Wilson *et al*.³ We conclude that a del22q11 is among the most frequent causes of genetic syndromes.

K DEVRIENDT
J-P FRYNS

Centre for Human Genetics, University of Leuven,
Herestraat 49, B-3000 Leuven, Belgium

G MORTIER
Department of Medical Genetics, University of Ghent,
Belgium

Table 1 The minimum annual incidence of del22q11 at birth in Flanders

Year	No of cases	Annual incidence per 100 000	95% CI	Total births
1992	11	16	6.5-25.4	68 613
1993	10	15	5.2-24.7	66 780
1994	11	17.1	7-27.3	63 851
1995	11	17.2	7-27.4	63 372
1996	8	12.6	2.3-22.7	63 550
Total	51	15.3	13.3-17.2	326 166

M-N VAN THIENEN
Department of Medical Genetics, University of
Antwerp, Belgium

K KEYMOLEN
Centre for Medical Genetics, University of Brussels,
Belgium

- 1 Scambler PJ, Kelly D, Lindsay E, *et al.* Velocardio-facial syndrome associated with chromosome 22 deletions encompassing the Di-George locus. *Lancet* 1992;339:1138-9.
- 2 Driscoll DA, Salvin J, Sellinger B, *et al.* Prevalence of 22q11 microdeletions in Di-George and velocardiofacial syndromes: implications for genetic counselling and prenatal diagnosis. *J Med Genet* 1993;30:813-17.
- 3 Wilson DI, Cross IE, Wren C, *et al.* Minimum prevalence of chromosome 22q11 deletions. *Am J Hum Genet* 1994;55:A169.
- 4 Du Montcel ST, Mendizabal H, Aymé S, Lévy A, Philip N. Prevalence of 22q11 microdeletion. *J Med Genet* 1996;33:719.

Sharing of PPT mutations between distinct clinical forms of neuronal ceroid lipofuscinoses in patients from Scotland

The neuronal ceroid lipofuscinoses (NCLs, Batten disease) are a group of rare inherited neurodegenerative diseases of childhood classified according to their age of onset and ultrastructural appearance of storage material. A very unusual cluster of NCL cases is found in the west of Scotland.¹ Some patients have early infantile onset, INCL, a disease almost entirely confined to Finland and characterised on ultrastructure by granular osmiophilic deposits (GROD). Other patients have a later, juvenile, onset but are found to have GROD (vJNCL/GROD) rather than the fingerprint profiles usually found in juvenile onset cases. INCL in Finland is caused by mutations in the gene encoding palmitoyl protein thioesterase, PPT.² It was recently reported that cases of vJNCL/GROD from Scotland (and elsewhere) also result from mutations in this gene.³ We set out to determine the disease causing mutations in PPT in Scottish INCL patients to establish whether there was any sharing of mutations with vJNCL/GROD.

Four patients with a diagnosis of INCL were analysed. Exons of PPT were amplified from genomic DNA by PCR and then sequenced in forward and reverse directions.³ We found mutations in PPT in all four patients and of the eight chromosomes analysed three different nonsense mutations (Leu10STOP, Lys55STOP, and Arg151STOP, table 1) were present. Therefore, all the INCL patients are homozygous for mutations predicted to result in truncation of PPT. For each patient we amplified parental DNA and were able to show Mendelian inheritance of mutations.

Two of these mutations (Leu10STOP and Arg151STOP) are found in patients with vJNCL/GROD.³ However, in these cases

nonsense mutations do not occur in homozygous form and are only found in combination with a missense mutation. Therefore, we can now show that two clinically distinct forms of NCL are caused by shared mutations in PPT. The clinical significance of these findings is that the severity of the disease in these patients is dependent on the combination and type of mutations present.

There are several diseases in which different mutations in the same gene cause dissimilar clinical phenotypes, for example, CFTR (cystic fibrosis and congenital bilateral absence of the vas deferens⁴). Types A and B Niemann-Pick disease, like the NCLs, are lysosomal storage disorders which are both caused by mutations in the acid sphingomyelinase gene. As in this study, the same mutation has been found in both forms and the age of onset and severity of the phenotype is dependent on the other allele.⁵

Geographical clustering of a rare autosomal recessive genetic disease suggests a founder effect with patients inheriting the same ancestral disease chromosomes. Detailed genealogical information is not available on these patients. However, all but one of the 18 disease chromosomes in INCL and vJNCL/GROD cases are accounted for by two nonsense and one missense mutations (table 1) and it is likely that these are derived from individual ancestral chromosomes. High resolution haplotype analysis and population studies to determine carrier rates will be required to resolve the issue.

We thank the Medical Research Council (PBM, NDEG) and Action Research (KYL) for financial support.

PATRICIA B MUNROE
NICHOLAS D E GREENE
KIT-YI LEUNG
SARA E MOLE
R MARK GARDINER
HANNAH M MITCHISON

Department of Paediatrics, University College London
Medical School, Rayne Institute, 5 University Street,
London WC1E 6JJ, UK

JOHN B P STEPHENSON
YANICK J CROW

Departments of Paediatric Neurology and Medical
Genetics, Yorkhill Hospitals NHS Trust,
Glasgow G3 8SJ, UK

- 1 Crow YJ, Tolmie JL, Howatson AG, Patrick WJA, Stephenson JBP. Batten disease in the West of Scotland 1974-1995 including five cases of the juvenile form with granular osmiophilic deposits. *Neuropediatrics* 1997;28:140-4.
- 2 Vesa J, Hellsten E, Verkruyse LA, *et al.* Mutations in the palmitoyl-protein thioesterase gene causing infantile neuronal ceroid lipofuscinosis. *Nature* 1995;376:584-7.
- 3 Mitchison HM, Hofmann SL, Becerra CHR, *et al.* Mutations in palmitoyl-protein thioesterase gene (PPT; CLN1) causing juvenile neuronal ceroid lipofuscinosis with granular osmiophilic deposits. *Hum Mol Genet* 1998;7:291-7.
- 4 Chillon M, Casals I, Mercier B, *et al.* Mutations in the cystic-fibrosis gene in patients with congenital absence of the vas-deferens. *N Engl J Med* 1995;332:1475-80.

- 5 Levran O, Desnick RJ, Schuchman EH. Niemann-Pick type B disease. Identification of a single codon deletion in the acid sphingomyelinase gene and genotype/phenotype correlations in type A and B patients. *J Clin Invest* 1991;88:806-10.

PTEN and prostate cancer

The PTEN gene (phosphate and tensin homologue), located on 10q23,^{1,2} has been reported to be the Cowden disease susceptibility gene. Germline mutations in PTEN have been found in patients with this syndrome.³ This disorder is characterised by the development of hamartomas at various sites, as well as an increased predisposition for thyroid cancer and for breast cancer in women.⁴⁻⁶ PTEN has also been reported as being altered in other types of cancer including glioblastoma,^{1,2} endometrial carcinoma,^{7,8} and kidney carcinoma.² This gene has additionally been suggested to play a role in prostate cancer as PTEN alterations have been found in multiple prostate cancer cell lines.¹

In order to investigate the role of mutations in the PTEN gene in primary prostate cancer, we analysed microdissected prostate adenocarcinoma tissue from 28 patients with histopathologically confirmed cancer. All nine exons of PTEN were PCR amplified and screened for mutations by single strand conformational polymorphism analysis (SSCP). Samples displaying mobility shifts were subjected to DNA sequence analysis. This analysis failed to detect homozygous deletions of the PTEN gene in any sample. A heterozygous mutation was identified in only one of the prostate tumour samples and was characterised as a single base deletion in codon 68 (TAC→AC). Additionally, an A→G polymorphism 96 bp upstream of the beginning of exon 2 was found in nine of 28 samples (32.1%). Based on this analysis, we conclude that mutations of the PTEN gene are rare in primary prostate cancers.

EVAN A FACHER

Department of Epidemiology and Biostatistics, Case
Western Reserve University, 2500 MetroHealth Drive,
Rammelkamp-R459, Cleveland, OH 44109, USA

JOHN C LAW

Department of Human Genetics, University of
Pittsburgh, USA

- 1 Li J, Yen C, Liaw D, *et al.* PTEN, a putative protein tyrosine phosphatase gene mutated in human brain, breast, and prostate cancer. *Science* 1997;275:1943-7.
- 2 Steck PA, Pershouse MA, Jasser SA, *et al.* Identification of a candidate tumour suppressor gene, MMAC1, at chromosome 10q23.3 that is mutated in multiple advanced cancers. *Nat Genet* 1997;15:356-62.
- 3 Liaw D, Marsh DJ, Li J, *et al.* Germline mutations of the PTEN gene in Cowden disease, an inherited breast and thyroid cancer syndrome. *Nat Genet* 1997;16:64-7.
- 4 Carlson HE, Burns TW, Davenport SL, *et al.* Cowden disease: gene marker studies and measurements of epidermal growth factor. *Am J Hum Genet* 1986;38:908-17.
- 5 Starink TM, van der Veen JP, Arwert F, *et al.* The Cowden syndrome: a clinical and genetic study in 21 patients. *Clin Genet* 1986;29:222-33.
- 6 Eng C, Murday V, Seal S, *et al.* Cowden syndrome and Lhermitte-Duclos disease in a family: a single genetic syndrome with pleiotropy? *J Med Genet* 1994;31:458-61.
- 7 Kong D, Suzuki A, Zou TT, *et al.* PTEN1 is frequently mutated in primary endometrial carcinomas. *Nat Genet* 1997;17:143-4.
- 8 Tashiro H, Blazes MS, Wu R, *et al.* Mutations in PTEN are frequent in endometrial carcinoma but rare in other common gynecological malignancies. *Cancer Res* 1997;57:3935-40.

Table 1 PPT mutations in Scottish INCL and vJNCL/GROD

Case	Disease	Mutations
389	INCL	Arg151STOP/Arg151STOP
390	INCL	Arg151STOP/Lys55STOP
392	INCL	Arg151STOP/Leu10STOP
391	INCL	Leu10STOP/Leu10STOP
105, 341, 346	vJNCL/GROD ³	Arg151STOP/Thr75Pro
325, 345	vJNCL/GROD ³	Leu10STOP/Thr75Pro

Bold type indicates mutations present in both INCL and vJNCL/GROD.