Short reports

A novel mutation in the mitochondrial tRNA^{Ser(UCN)} gene in a family with non-syndromic sensorineural hearing impairment

Tim P Hutchin, Mick J Parker, Ian D Young, Adrian C Davis, Louise J Pulleyn, Jayne Deeble, Nicholas J Lench, Alex F Markham, Robert F Mueller

Abstract

We describe a family with non-syndromic sensorineural hearing impairment inherited in a manner consistent with maternal transmission. Affected members were found to have a novel heteroplasmic mtDNA mutation, T7510C, in the tRNA^{Ser(UCN)} gene. This mutation was not found in 661 controls, is well conserved between species, and disrupts base pairing in the acceptor stem of the tRNA, making it the probable cause of hearing impairment in this family. Sequencing of the other mitochondrial tRNA genes did not show any other pathogenic mutations. Four other mutations causing hearing impairment have been reported in the tRNA^{Ser(UCN)} gene, two having been shown to affect tRNA^{Ser(UCN)} levels. With increasing numbers of reports of mtDNA mutations causing hearing impairment, screening for such mutations should be considered in all cases unless mitochondrial inheritance can be excluded for certain.

(J Med Genet 2000;37:692–694)

Keywords: hearing impairment; mtDNA mutation; tRNA^{Ser(UCN)}

Sensorineural hearing loss is the most common phenotype of mitochondrial diseases and to date four point mutations of the mitochondrial DNA (mtDNA) have been reported in families with non-syndromic, maternally inherited hearing impairment (MIHI). The most common, the A1555G mutation in the 12S rRNA gene (MIM 561000), also causes increased sensitivity to the ototoxic side effects of aminoglycosides.^{1 2} The other three, A7445G, 7472insC, and T7511C, all lie in the tRNA^{Ser(UCN)} gene (MIM 590080) and in some cases hearing impairment can also be associated with other features, such as palmoplantar keratoderma or ataxia.^{3–6}

We describe a white family with non-syndromic sensorineural hearing impairment transmitted in a manner consistent with maternal inheritance (fig 1A), that is, there was no paternal transmission and II.2 had hearing impaired children by two different fathers. The proband (IV.6) was formally diagnosed as having sensorineural hearing impairment aged 15

months. His hearing loss is profound but asymmetrical, with an average loss in the better ear of 103 dB HL (associated with a gently sloping audiogram profile) and 115 dB HL in the other ear. His younger sister (IV.7) was not formally diagnosed as having a hearing impairment until 5 years of age. Her sensorineural hearing loss is severe but asymmetrical with an average loss in the better ear of 66 dB HL (with a gently sloping audiogram profile) and 75 dB HL in the other ear.

Their mother (III.9) had not previously been diagnosed as having a hearing loss but she was aware that her hearing was “dull at times”. Portable audiometry showed her to have a moderate sensorineural hearing loss, with an average loss in the better ear of 41 dB HL and 48 dB HL in the other ear.

The mothers' four sibs all have hearing impairment to varying degrees. The only other clinical feature of note in the family is mental handicap in a maternal aunt (III.7), which has previously been attributed to birth asphyxia. The maternal grandmother (II.2) died at the age of 80 and had suffered a loss of hearing for her last 10 years. With the exception of an uncle (III.5), who was only related by marriage, there was no history of exposure to ototoxic drugs or other illness in the family.

Following informed consent, DNA was extracted from blood taken from the proband, his sister, and both parents. No other family members were available for study nor were we able to carry out any further analysis of those available to us. Analysis by PCR and restriction enzyme digestion showed the absence of the A1555G, A7445G, T7511C, and 7472insC mutations. Further analysis, however, showed the gain of a *HinfI* site around base pair (bp) 7510 in the tRNA^{Ser(UCN)} gene of the proband, his sister, and mother (fig 1B). Sequencing showed this to be a T to C transition at bp 7510 (fig 1C). The T7510C mutation was heteroplasmic in all three affected family members tested, that is, >95% mutant in the two sibs and 90% in the mother. The following points provide further support for the T7510C mutation as the most probable cause of the hearing impairment in this family. (1) The *HinfI* site gain at bp 7510 is extremely rare. We did not find it in 141 white controls here, nor was it reported in 520 other controls in published

Molecular Medicine Unit, St James's University Hospital, Leeds LS9 7TF, UK
T P Hutchin
L J Pulleyn
J Deeble
N J Lench
A F Markham

MRC Institute of Hearing Research, University Park, Nottingham NG7 2RD, UK
M J Parker
A C Davis

Department of Clinical Genetics, City Hospital, Nottingham NG5 1PB, UK
M J Parker
I D Young

Department of Clinical Genetics, St James's University Hospital, Leeds LS9 7TF, UK
R F Mueller

Correspondence to:
Dr Hutchin, Molecular Medicine Unit, Clinical Sciences Building, St James's University Hospital, Beckett Street, Leeds LS9 7TF, UK, t.p.hutchin@leeds.ac.uk

Revised version received
22 March 2000
Accepted for publication
10 April 2000

Figure 1 (A) Pedigree of the family. Solid symbols denote those with hearing impairment. (B) Gain of *HinfI* restriction enzyme site in affected subjects. mtDNA was amplified with primers corresponding to bp 7392-7410 (forward) and 7608-7588 (reverse) and the PCR fragment digested to completion with 10 units of *HinfI* and fragments separated on a 2% agarose gel. The normal product gives fragments of 150 bp and 66 bp, the *HinfI* gain cuts the 150 bp fragment into 101 bp and 49 bp. Patient ID numbers correspond to those in (A), lanes 5 and 6 are hearing controls, and M is pBR322 *HaeIII* marker. (C) DNA sequence of the mtDNA heavy strand showing the T to C base change at bp 7510 in the proband compared to a hearing control. The proportion of mutant was estimated by densitometry of both the agarose and sequencing gels.

reports.^{7,8} (2) The base change disrupts a hydrogen bond in the acceptor stem of the tRNA^{Ser(UCN)} (fig 2A), which may affect tRNA levels or function. Similar mutations in this acceptor stem at bp 7511 and 7512 have previously been shown to cause hearing loss.^{4,9,10} (3) The 7510 residue is highly conserved in a wide range of species (fig 2B) forming an A-U base pair in all except bovine.

Sequencing of the other mitochondrial tRNA genes from this family did not show any other changes except for a homoplasmic A to G base change at bp 4336 in the tRNA^{Gln} gene. This base change is present in about 1% of the white population and although it has been reported in patients with Alzheimer's or Parkinson's disease in some studies,⁷ any specific disease association remains uncertain.¹¹ Hearing loss has not been reported in any of these persons, although such information was not actively sought. The A4336G mutation is therefore unlikely to be the primary cause of hearing loss in this family although it may contribute to the high penetrance of the T7510C mutation in this family.

Several mtDNA point mutations have been associated with non-syndromic sensorineural

hearing loss. Generally these mutations are homoplasmic or present at very high levels (>95%) and presumably exert only very mildly deleterious effects which are sufficient to affect the inner ear but with no noticeable effect on other tissues. The reason for this tissue specificity is not known. Fischel-Ghodsian¹² has suggested it could be because of different processing of the mitochondrial genes in the cochlea. Such a difference between tissues has been shown to exist, for example the A3302G mutation in the tRNA^{Leu(UUR)} gene¹³ causes a defect in RNA processing in skeletal muscle but not fibroblasts where there is no phenotypic defect.

It is interesting that five mutations in or affecting the tRNA^{Ser(UCN)} gene (T7510C, T7511C, T7512C, A7445G, and 7472insC) have all been associated with hearing loss. Two of these, A7445G and 7472insC, have been shown to cause a significant reduction in tRNA^{Ser(UCN)} levels and a mild mitochondrial defect in lymphoblasts or osteosarcoma cells.^{14,15} Whether or not the T7510C mutation has any effect on the processing of mitochondrial genes in the cochlea can only be answered by studying cells from the inner ear.

The increasing number of reports of families with hearing impairment resulting from

A

B

	Aminoacyl stem		D stem		D loop	
Human	GAAA	A	AG-	T-	CATG	GAGGC-----
Orang-utan	GAAA	A	AG-	T-	ACTG	GGGGT-----
Rat	GAGA	A	AG-	A-	CATA	ATGGT-----
Mouse	GAGA	A	AG-	A-	CATA	TAGGA-----
Cow	GAGA	G	AG-	A-	CATA	GAGGT-----
Chicken	AAGA	A	AG-	AA	GCAT	TAAGTGGTTTG
Fruit fly	AGTT	A	ATG	A-	GCTT	GAAT-----
Atlantic cod	AAGA	A	AG-	T-	GACA	GAGTGGT----
Frog	AAGA	A	AA-	TG	GCA-	GAGTGGTGA--
Sea anemone	GGAG	A	AG-	TG	GCAC	GATAGGA----

Figure 2 (A) Proposed secondary structure of tRNA^{Ser(UCN)}. The gene is encoded on the light strand and thus the base change is shown as A to G in the tRNA. (B) Interspecies homology of mitochondrial tRNA^{Ser(UCN)} genes. Aligned sequences were obtained from the tRNA compilation manual.¹⁷ Position 7510 is indicated by the boxed area.

mtDNA mutations, and the relatively high frequency of the A1555G mutation in Spain found by Estivill *et al*,¹⁶ suggest that mtDNA mutations may be a more common cause of familial non-syndromic hearing impairment than previously estimated. Sensorineural hearing impairment caused by mtDNA mutations is usually of childhood onset and progressive, although some persons may not be affected until much later in life. Given that some families with mitochondrial inherited hearing impairment may not be large enough to estab-

lish a pattern of maternal inheritance with certainty, screening for mtDNA mutations should always be considered in any family which does not exhibit paternal transmission of the hearing impairment.

This work was funded in part by a grant from Defeating Deafness. Work in the authors' laboratories is also supported by the Medical Research Council, European Community, Wellcome Trust, and Yorkshire Cancer Research. We are grateful to the family for their cooperation.

- Hutchin T, Haworth I, Higashi K, Fischel-Ghodsian N, Stoneking M, Saha N, Arnos C, Cortopassi G. A molecular basis for human hypersensitivity to aminoglycoside antibiotics. *Nucleic Acids Res* 1993;21:4174-9.
- Prezant TR, Agopian JV, Bohlman MC, Bu X, Oztas S, Qiu WQ, Arnos KS, Cortopassi GA, Jaber L, Rotter JL, Shohat M, Fischel-Ghodsian N. Mitochondrial ribosomal RNA mutation associated with both antibiotic-induced and non-syndromic deafness. *Nat Genet* 1993;4:289-94.
- Reid FM, Vernham GA, Jacobs HT. A novel mitochondrial point mutation in a maternal pedigree with sensorineural deafness. *Hum Mutat* 1994;3:243-7.
- Sue CM, Tanji K, Hadjigeorgiou GM, Andreu AL, Nishino I, Krishna S, Bruno C, Hirano M, Shanske S, Bonilla E, Fischel-Ghodsian N, DiMauro S, Friedman R. Maternally inherited isolated hearing loss in a large kindred with a novel T7511C mutation in the mitochondrial DNA tRNA(Ser(UCN)) gene. *Neurology* 1999;52:1905-8.
- Tiranti V, Chariot P, Carella F, Toscano A, Soliveri P, Girlanda P, Carrara F, Fratta GM, Reid FM, Mariotti C, Zeviani M. Maternally inherited hearing loss, ataxia and myoclonus associated with a novel point mutation in mitochondrial tRNA^{Ser(UCN)} gene. *Hum Mol Genet* 1995;8:1421-7.
- Verhoeven K, Ensink RJH, Tiranti V, Huygen PLM, Johnson DF, Chatteman IS, Van Laer L, Verstreken M, Van de Heyning P, Fischel-Ghodsian N, Zeviani M, Cremers CW, Willems PJ, Van Camp G. Hearing impairment and neurological dysfunction associated with a mutation in the mitochondrial tRNA^{Ser(UCN)} gene. *Eur J Hum Genet* 1999;7:45-51.
- Shoffner JM, Brown MD, Torroni A, Lott MT, Cabell MF, Mirra SS, Beal MF, Yang CC, Gearing M, Salvo R, Watts RL, Juncos JL, Hansen LA, Crain BJ, Fayad M, Reckord CL, Wallace DC. Mitochondrial DNA variants observed in Alzheimer disease and Parkinson disease patients. *Genomics* 1993;17:171-84.
- Kogelnik AM, Lott MT, Brown MD, Navathe SB, Wallace DC. MITOMAP: a human mitochondrial genome database - 1998 update. *Nucleic Acids Res* 1998;26:112-15.
- Nakamura M, Nakano S, Goto Y, Ozawa M, Nagahama Y, Fukuyama H, Akiuchi I, Kaji R, Kimura J. A novel point mutation in the mitochondrial tRNA(Ser(UCN)) gene detected in a family with MERRF/MELAS overlap syndrome. *Biochem Biophys Res Commun* 1995;214:86-93.
- Jaksch M, Klopstock T, Kurlmann G, Dörner M, Hofman S, Kleinle S, Hegemann S, Weissert M, Müller-Höcker J, Pongratz D, Gerbitz KD. Progressive myoclonus epilepsy and mitochondrial myopathy associated with mutations in the tRNA(Ser(UCN)) gene. *Ann Neurol* 1998;44:635-40.
- Hutchin TP, Heath PR, Pearson RCA, Sinclair AJ. Mitochondrial DNA mutations in Alzheimer's disease. *Biochem Biophys Res Commun* 1997;241:221-5.
- Fischel-Ghodsian N. Mitochondrial deafness mutations reviewed. *Hum Mutat* 1999;13:261-70.
- Bindoff LA, Howell N, Poulton J, McCullough DA, Morten KJ, Lightowlers RN, Turnbull DM, Weber K. Abnormal RNA processing associated with a novel tRNA mutation in mitochondrial DNA. A potential disease mechanism. *J Biol Chem* 1993;268:19559-64.
- Guan MX, Enriquez JA, Fischel-Ghodsian N, Puranam RS, Lin CP, Maw MA, Attardi G. The deafness-associated mitochondrial DNA mutation at position 7445, which affects tRNA^{Ser(UCN)} precursor processing, has long-range effects on NADH dehydrogenase subunit ND6 gene expression. *Mol Cell Biol* 1988;18:5868-79.
- Toompuu M, Tiranti V, Zeviani M, Jacobs HT. Molecular phenotype of the nt 7472 deafness-associated mitochondrial mutation in osteosarcoma cell hybrids. *Hum Mol Genet* 1999;8:2275-83.
- Estivill X, Govea N, Barcelo A, Perello E, Badenas C, Romero E, Moral L, Scozzari R, D'Urbano L, Zeviani M, Torroni A. Familial progressive sensorineural deafness is mainly due to the mitochondrial DNA A1555G mutation and is enhanced by treatment with aminoglycosides. *Am J Hum Genet* 1998;62:27-35.
- Sprinzi M, Horn C, Brown M, Ioudovitch A, Steinberg S. tRNA compilation manual, 1999; <http://www.uni-bayreuth.de/departments/biochemie/trna/>