Electronic letter

Autosomal recessive non-syndromic hearing loss in the Lebanese population: prevalence of the 30delG mutation and report of two novel mutations in the connexin 26 (*GJB2*) gene

Myrna Mustapha, Nabiha Salem, Valérie Delague, Eliane Chouery, Michella Ghassibeh, Myriam Rai, Jacques Loiselet, Christine Petit, André Mégarbané

Unité de Génétique Médicale, Faculté de Médecine, Université Saint-Joseph, Beirut, Lebanon

M Mustapha
N Salem
V Delague
E Chouery
M Ghassibeh
M Rai
J Loiselet
A Mégarbané

Unité de Génétique des Déficiences Sensoriels, CNRS-URA 1968, Institut Pasteur, Paris, France

M Mustapha
C Petit

Correspondence to:
Dr Mégarbané, Unité de Génétique Médicale, Faculté de Médecine, Université Saint-Joseph, 42 rue de Grenelle, 75007 Paris, France,
megarban@dm.net.lb

Table 1 Different loci found in 48 Lebanese families with autosomal recessive non-syndromic hearing loss

Designation	Locus	Gene	Encoded product	No of families
DFNB1	13q12	<i>GJB2</i>	Connexin 26	16
DFNB3	17p11.2	<i>MYO15</i>	Myosin XV	4
DFNB4,14,17	7q31	<i>PDS</i> (DFNB4)		3
DFNB5	14q12			1
DFNB6	3p14-p21			1
DFNB7	9q13-q21			1
DFNB9	2p22-p23	<i>OTOF</i>		4+1?
DFNB10	21q22.3	<i>TMPRSS3</i>		1
DFNB12	10q21-q22	<i>CDH23</i>	Cadherin	2+2?
DFNB13	7q34-q36			1
DFNB14	7q31			1
DFNB21	11q23-q25	<i>TECTA</i>	α -tectorin	2
DFNB22	16p11-q21			1
DFNB ?	—			7

EDITOR—The most common sensory deficit in humans is hearing loss, affecting 1 in 1000 children, with approximately half of the cases having a genetic cause. The majority of these genetic causes are non-syndromic, of which approximately 75% have an autosomal recessive mode of inheritance.¹ So far, nearly 30 genes that cause non-syndromic recessive deafness (NSRD) have been located (for review see <http://dnalab-www.uia.ac.be/dnalab/hhh>). The loci corresponding to NSRD are designated DFNB, with a number corresponding to the chronology of their localisation. The first locus, DFNB1 (MIM 220290), located on chromosome 13q11-12,² has been shown to be responsible for nearly half of NSRD owing to mutations in the gene encoding the gap junction protein connexin 26 (*GJB2*) (MIM 121011).^{3,4} One mutation, 30delG (also referred to as 35delG), accounts for the majority of mutations in this gene in some ethnic groups,³⁻⁸ while it is rarer in others.⁹⁻¹⁵

The geographical position of Lebanon, a small country of 10 500 km² on the eastern shores of the Mediterranean sea, has made it a historical crossroads between Asia, Africa, and Europe. As a consequence, the Lebanese population shows a wide genetic diversity, with no less than 17 ethno-religious communities. Today, the population is approximately 4 million people, with a world wide diaspora

estimated at 15 million. In Lebanon, consanguineous marriages are still frequent (from 10 to 30%), favouring the incidence of autosomal recessive diseases,¹⁶ such as haemoglobinopathies, sickle cell anaemia, familial Mediterranean fever, congenital hypothyroidism, cystic fibrosis, and deafness.

The purpose of this study is to summarise the different NSRD loci found in a case series of Lebanese families and to determine the carrier frequency of the 30delG mutation in selected Lebanese subjects.

Material and methods

Forty eight multiplex Lebanese families with non-syndromic congenital moderate to profound deafness, from various regions of the country, were included in this analysis. Informed consent was obtained from each family member before clinical investigation, blood sampling, and DNA analyses. Hearing loss was documented by audiological testing and a thorough examination was conducted to ensure the absence of any clinical sign suggesting another diagnosis. The pattern of inheritance was autosomal recessive for all the families studied.

The methods used for DNA extraction and linkage analyses were the same as those presented elsewhere.¹⁷⁻²³ In families linked to the DFNB1 locus, affected subjects were first tested for the common 30delG mutation according to Storm *et al.*⁸ In the absence of this mutation, amplification of the coding exon of the *GJB2* gene and DNA sequencing was performed according to Denoyelle *et al.*³

Additionally, 300 unrelated Lebanese subjects with no familial history of hearing problems, selected from our laboratory database to represent all ethno-religious Lebanese communities, were analysed to assess the prevalence of the 30delG mutation. Informed consent was obtained from each person involved and DNA was extracted from peripheral blood leucocytes following standard methods.²⁴

Results

Of the 48 Lebanese pedigrees investigated with non-syndromic recessive moderate to profound

Figure 1 DNA sequencing of the *GJB2* coding exon showing (A) a missense mutation, replacing arginine by histidine at codon 32 (R32H), and (B) an insertion of an adenine in position 291 (291insA) creating a stop codon.

deafness, 37 were Muslims, 10 Christians, and one Druze. Nearly 92% (44) of them were consanguineous. Table 1 summarises the different loci found in our population. Sixteen of these families were linked to the *DFNB1* locus (33%). In all of them, the affected subjects had a severe to profound hearing loss. In 15 of them, affected subjects were homozygous for the 30delG mutation in the *GJB2* gene. In the last one, a non-consanguineous Christian Maronite family, affected subjects were compound heterozygotes for two novel mutations, a missense mutation, replacing arginine by histidine at codon 32 (R32H), and an insertion of an adenine in position 291 (291insA) creating a stop codon (fig 1A, B).

In the remaining 32 families, at least 13 implicated loci were found (table 1). In three of these families, it was not possible to determine exactly which of three loci, *DFNB4*, 14, or 17, were implicated. They are all localised on the long arm of chromosome 7 at q31, and these three families were not sufficiently informative to allow a precise localisation. In four families, linkage to *DFNB9*, 10, and 12 was highly likely but could not be proven since the lod score was below 3 at $\theta=0.00$. Finally, seven families have not been linked to any of the already reported loci so far and are still undergoing more elaborate analysis.

In the 300 subjects tested for the 30delG mutation, we found seven heterozygotes, four Christians and three Muslims. Four were men and three women. This mutation was found with an allele frequency of 1/86, giving a prevalence of 1/7400 newborns affected with deafness secondary to a 30delG mutation.

Discussion

The majority of the cases of hereditary deafness are non-syndromic. So far, at least 77 chromosomal loci have been identified, 39 for autosomal dominant DFNA, 30 for autosomal recessive DFNB, and eight for X linked DFN¹ (<http://dnalab-www.uia.ac.be/dnalab/hhh>).

The contribution of the *GJB2* gene in NSRD in Lebanon (33%) is lower than in other studies around the world: 49% in Italy and Spain,⁶ 40% in the USA,²⁵ and 41.3% in Israel.²⁶ However, the allele frequency of the 30delG mutation among our familial DFNB1 cases was ~94%. Although, our sample size is relatively small, we suggest that this allele frequency is higher than in European populations where it has been estimated at 60-80% of all known *GJB2* mutations.¹⁴ The haplotype of affected subjects from most of these families was different (data not shown) suggesting that this mutation arose independently in each family. Furthermore, in order to determine the rate of 30delG mutation, we screened for the latter, by a PSDM method (PCR-site directed mutagenesis), 300 unrelated Lebanese subjects equally distributed by sex and community for the 30delG mutation. This mutation was found with a carrier frequency of 1/43, which is lower than the carrier frequency of 1/31-35 found in white populations.^{6 27 28} Also, we did not find an excess in any particular ethno-religious community. Thus, we suggest that the 30delG

mutation is common in our familial cases linked to DFNB1, most likely because of the highly mutable characteristic of a stretch of six identical nucleotides³ and not because of a particular ancestral mutation. Random DNA samples from 100 males and 100 females (400 chromosomes) were subsequently tested for the two novel mutations found in one family and none was found, confirming the absence of ancestral mutation in the *GJB2* gene in the Lebanese population.

For the remaining families not linked to DFNB1 (64%), 12 already reported loci were implicated (table 1). Apart from DFNB1, the most frequent loci in our population were DFNB3 and DFNB9. DFNB3 was initially reported in Benkala, an Indonesian village,²⁹ and also in India.³⁰ The haplotypes of the affected subjects from these families and ours were different. The DFNB9 locus was found in Lebanon,¹⁷ but also in Turkey³¹ and India.³² In the Lebanese families, the same nonsense mutation, tyr703 to ter, was found in the *OTOF* gene,²³ raising the possibility of an ancestral mutation at this locus in Lebanon. Finally, it is worth remembering that five different DFNB loci, DFNB9, 12, 13, 14, 21, and 22, were found in Lebanese families.¹⁷⁻²² All these results confirm the genetic diversity of the Lebanese population.

Our strategy in investigating families with NSRD has changed since the report of the mutations in the *GJB2* gene. Affected patients are first tested for the 30delG mutation and the gene is sequenced if it is absent. If the results are negative, elimination of the known loci and a genome wide search is performed. In one family, several affected subjects with 30delG present on one allele were detected. No other mutation was found after sequencing. The genome wide search allowed us to link this family to DFNB12. This finding agrees with the suggested possibility⁶ that some subjects might be carriers of the 30delG mutation because of its high frequency in the population and have mutations leading to hearing loss in a different gene. Amazingly, this family also contained subjects with Usher syndrome, linked to the Usher 2A locus.

In conclusion, the precise number of people affected with early and profound hearing loss in Lebanon is unknown. Nevertheless, because of the high rate of consanguineous marriages, the incidence may be higher than has been reported world wide. Our findings indicate that screening for one mutation, 30delG, would allow the early detection of about one third of NSRD cases in Lebanon. Further analysis of the different genes implicated in hearing loss in the Lebanese population will improve the performance of clinicians providing genetic counselling, especially in consanguineous Lebanese families.

The first two authors contributed equally to this work. We are indebted to the participating families for their kind cooperation. Special thanks go to Professor G Lefranc for his continuous support and advice and to Dominique Weil for her help. This work was supported by grants from the Saint Joseph University for Scientific Research.

- Robertson NG, Morton CC. Beginning of a molecular era in hearing and deafness. *Clin Genet* 1999;55:149-59.
- Guilford P, Ben Arab S, Blanchard S, Levilliers J, Weissenbach J, Belkahlia A, Petit C. A non-syndrome form of neurosensory, recessive deafness maps to the pericentromeric region of chromosome 13q. *Nat Genet* 1994;6:24-8.
- Denoyelle F, Weil D, Maw MA, Wilcox SA, Lench NJ, Allen-Powell DR, Osborn AH, Dahl HH, Middleton A, Houseman MJ, Dode C, Marlin S, Boullila-ElGaid A, Grati M, Ayadi H, BenArab S, Bitoun P, Lina-Granade G, Godet J, Mustapha M, Loiselet J, El-Zir E, Aubois A, Joannard A, Petit C. Prelingual deafness: high prevalence of a 30delG mutation in the connexin 26 gene. *Hum Mol Genet* 1997;6:2173-7.
- Kelsell DP, Dunlop J, Stevens HP, Lench NJ, Liang JN, Parry G, Mueller RF, Leigh IM. Connexin 26 mutations in hereditary non-syndromic sensorineural deafness. *Nature* 1997;387:80-3.
- Antoniadi T, Gronskov K, Sand A, Pampanos A, Brondum-Nielsen K, Petersen MB. Mutation analysis of the *GJB2* (connexin 26) gene by DGGE in Greek patients with sensorineural deafness. *Hum Mutat* 2000;16:7-12.
- Estivill X, Fortina P, Surrey S, Rabionet R, Melchionda S, D'Agruma L, Mansfield E, Rappaport E, Govea N, Mila M, Zelante L, Gasparini P. Connexin-26 mutations in sporadic and inherited sensorineural deafness. *Lancet* 1998;351:394-8.
- Green GE, Scott DA, McDonald JM, Woodworth GG, Sheffield VC, Smith RJ. Carrier rates in the midwestern United States for *GJB2* mutations causing inherited deafness. *JAMA* 1999;281:2211-16.
- Storm K, Wilcox S, Flothmann K, Van Camp G. Determination of the carrier frequency of the common *GJB2* (connexin-26) 35delG mutation in the Belgian population using an easy and reliable screening method. *Hum Mutat* 1999;14:263-6.
- Abe S, Usami S, Shinkawa H, Kelley PM, Kimberling WJ. Prevalent connexin 26 gene (*GJB2*) mutations in Japanese. *J Med Genet* 2000;37:41-3.
- Adato A, Raskin L, Petit C, Bonne-Tamir B. Deafness heterogeneity in a Druze isolate from the Middle East: novel *OTOF* and *PDS* mutations, low prevalence of *GJB2* 35delG mutation and indication for a new DFNB locus. *Eur J Hum Genet* 2000;8:437-42.
- Brobbly GW, Muller-Myhsok B, Horstmann RD. Connexin 26 R143W mutation associated with recessive nonsyndromic sensorineural deafness in Africa. *N Engl J Med* 1998;338:548-50.
- Fuse Y, Doi K, Hasegawa T, Sugii A, Hibino H, Kubo T. Three novel connexin26 gene mutations in autosomal recessive non-syndromic deafness. *Neuroreport* 1999;10:1853-7.
- Morell RJ, Kim HJ, Hood LJ, Goforth L, Friderici K, Fisher R, Van Camp G, Berlin CI, Oddoux C, Ostrer H, Keats B, Friedman TB. Mutations in the connexin 26 gene (*GJB2*) among Ashkenazi Jews with nonsyndromic recessive deafness. *N Engl J Med* 1998;339:1500-5.
- Wilcox SA, Osborn AH, Dahl HH. A simple PCR test to detect the common 35delG mutation in the connexin 26 gene. *Mol Diagn* 2000;5:75-8.
- Zelante L, Gasparini P, Estivill X, Melchionda S, D'Agruma L, Govea N, Mila M, Monica MD, Lutfi J, Shohat M, Mansfield E, Delgrosso K, Rappaport E, Surrey S, Fortina P. Connexin26 mutations associated with the most common form of non-syndromic neurosensory autosomal recessive deafness (DFNB1) in Mediterraneans. *Hum Mol Genet* 1997;6:1605-9.
- Der Kaloustian VM, Naffah J, Loiselet J. Genetic diseases in Lebanon. *Am J Hum Genet* 1980;7:187-203.
- Chaib H, Place C, Salem N, Chardenoux S, Vincent C, Weissenbach J, El-Zir E, Loiselet J, Petit C. A gene responsible for a sensorineural nonsyndromic recessive deafness maps to chromosome 2p22-23. *Hum Mol Genet* 1996;5:155-8.
- Chaib H, Place C, Salem N, Dodé C, Chardenoux S, Weissenbach J, el Zir E, Loiselet J, Petit C. Mapping of DFNB12, a gene for a non-syndromal autosomal recessive deafness, to chromosome 10q21-22. *Hum Mol Genet* 1996;5:1061-4.
- Mustapha M, Chardenoux S, Nieder A, Salem N, Weissenbach J, el-Zir E, Loiselet J, Petit C. A sensorineural progressive autosomal recessive form of isolated deafness, DFNB13, maps to chromosome 7q34-q36. *Eur J Hum Genet* 1998;6:245-50.
- Mustapha M, Salem N, Weil D, el-Zir E, Loiselet J, Petit C. Identification of a locus on chromosome 7q31, DFNB14, responsible for prelingual sensorineural non-syndromic deafness. *Eur J Hum Genet* 1998;6:548-51.
- Mustapha M, Weil D, Chardenoux S, Elias S, El-Zir E, Beckmann JS, Loiselet J, Petit C. An alpha-tectorin gene defect causes a newly identified autosomal recessive form of sensorineural pre-lingual non-syndromic deafness, DFNB21. *Hum Mol Genet* 1999;8:409-12.
- Mustapha M, Chardenoux S, Salem N, El-Zir E, Beckman JS, Loiselet J, Petit C. DFNB22, a recessive form of sensorineural hearing loss, maps to chromosome 16p11-q21. *Eur J Hum Genet* (in press).
- Yasunaga S, Grati M, Cohen-Salmon M, El-Amraoui A, Mustapha M, Salem N, El-Zir E, Loiselet J, Petit C. A mutation in *OTOF*, encoding otoferlin, a FER-1-like protein, causes DFNB9, a nonsyndromic form of deafness. *Nat Genet* 1999;21:363-9.

- 24 Miller SA, Dykes DD, Polesky HF. A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Res* 1988; **16**:1215.
- 25 Kelley PM, Harris DJ, Comer BC, Askew JW, Fowler T, Smith SD, Kimberling WJ. Novel mutations in the connexin 26 gene (GJB2) that cause autosomal recessive (DFNB1) hearing loss. *Am J Hum Genet* 1998; **62**:792-9.
- 26 Sobe T, Vreugde S, Shahin H, Berlin M, Davis N, Kanaan M, Yaron Y, Orr-Urtreger A, Frydman M, Shohat M, Avraham KB. The prevalence and expression of inherited connexin 26 mutations associated with nonsyndromic hearing loss in the Israeli population. *Hum Genet* 2000; **106**: 50-7.
- 27 Lench N, Houseman M, Newton V, Van Camp G, Mueller R. Connexin-26 mutations in sporadic non-syndromal sensorineural deafness. *Lancet* 1998; **351**:415.
- 28 Gasparini P, Rabionet R, Barbujani G, Melchionda S, Petersen M, Brondum-Nielsen K, Metspalu A, Oitmaa E, Pisano M, Fortina P, Zelante L, Estivill X. High carrier frequency of the 35delG deafness mutation in European populations. Genetic Analysis Consortium of GJB2 35delG. *Eur J Hum Genet* 2000; **8**:19-23.
- 29 Friedman TB, Liang Y, Weber JL, Hinnant JT, Barber TD, Winata S, Arhya IN, Asher JH Jr. A gene for congenital, recessive deafness DFNB3 maps to the pericentromeric region of chromosome 17. *Nat Genet* 1995; **9**:86-91.
- 30 Liang Y, Wang A, Probst FJ, Arhya IN, Barber TD, Chen KS, Deshmukh D, Dolan DF, Hinnant JT, Carter LE, Jain PK, Lalwani AK, Li XC, Lupski JR, Moeljopawiro S, Morell R, Negrini C, Wilcox ER, Winata S, Camper SA, Friedman TB. Genetic mapping refines DFNB3 to 17p11.2, suggests multiple alleles of DFNB3, and supports homology to the mouse model shaker-2. *Am J Hum Genet* 1998; **62**:904-15.
- 31 Leal SM, Apaydin F, Barnwell C, Iber M, Kandogan T, Pfister M, Braendle U, Cura O, Schwalb M, Zenner HP, Vitale E. A second middle eastern kindred with autosomal recessive non-syndromic hearing loss segregates DFNB9. *Eur J Hum Genet* 1998; **6**:341-4.
- 32 Yasunaga S, Grati M, Chardenoux S, Smith TN, Friedman TB, Lalwani AK, Wilcox ER, Petit C. OTOF encodes multiple long and short isoforms: genetic evidence that the long ones underlie recessive deafness DFNB9. *Am J Hum Genet* 2000; **67**:591-600.