Genotype-phenotype relationships in Berardinelli-Seip congenital lipodystrophy

L Van Maldergem, J Magré, T E Khallouf, T Gedde-Dahl Jr, M Delépine, O Trygstad, E Seemanova, T Stephenson, C S Albott, F Bonnici, V R Panz, J-L Medina, P Bogalho, F Huet, S Savasta, A Verloes, J-J Robert, H Loret, M de Kerdanet, N Tubiana-Rufi, A Mégarbané, J Maassen, M Polak, D Lacombe, C R Kahn, E L Silveira, F H D'Abronzio, F Grigorescu, M Lathrop, J Capeau, S O'Rahilly

J Med Genet 2002;**39**:722–733

See end of article for authors' affiliations

Correspondence to:
Dr L Van Maldergem,
Centre de Génétique
Humaine, Institut de
Pathologie et de
Génétique, Allée des
Templiers 41, B-6280
Loverval, Belgium;
vmald@skypro.be

Revised version received
10 May 2002
Accepted for publication
20 June 2002

Generalised lipodystrophy of the Berardinelli-Seip type (BSCL) is a rare autosomal recessive human disorder with severe adverse metabolic consequences. A gene on chromosome 9 (*BSCL1*) has recently been identified, predominantly in African-American families. More recently, mutations in a previously undescribed gene of unknown function (*BSCL2*) on chromosome 11, termed *seipin*, have been found to be responsible for this disorder in a number of European and Middle Eastern families. We have studied the genotype/phenotype relationships in 70 affected subjects from 44 apparently unrelated pedigrees of diverse ethnic origin. In all subjects, hepatic dysfunction, hyperlipidaemia, diabetes mellitus, and hypertrophic cardiomyopathy were significant contributors to morbidity with no clear differences in their prevalence between subjects with *BSCL1* or *BSCL2* and those with evidence against cosegregation with either chromosome 9 or 11 (designated *BSCLX*). *BSCL2* appears to be a more severe disorder than *BSCL1* with a higher incidence of premature death and a lower prevalence of partial and/or delayed onset of lipodystrophy. Notably, subjects with *BSCL2* had a significantly higher prevalence of intellectual impairment than those with *BSCL1* or *BSCLX* ($p < 0.0001$, OR 17.0, CI 3.6 to 79.0). The higher prevalence of intellectual impairment and the increased risk of premature death in *BSCL2* compared to *BSCL1* emphasise the importance of molecular diagnosis of this syndrome and have clear implications for genetic counselling.

Congenital lipodystrophy of the Berardinelli-Seip type (BSCL, OMIM 269700) is a rare condition characterised by a near total absence of adipose tissue.^{1–3} Because of the absence of functional adipocytes, dietary and endogenously synthesised lipid is stored aberrantly in metabolically important tissues such as muscle and liver. This leads to severe insulin resistance and, ultimately, diabetes mellitus, which is often difficult to control. Other metabolic consequences include hypertriglyceridaemia, which can be severe, and hepatic steatosis, which can progress to cirrhosis and death from hepatic failure. For reasons as yet unexplained, affected subjects also frequently develop a hypertrophic cardiomyopathy that can lead to death from cardiac failure. Acanthosis nigricans, acromegaly features, and muscle hypertrophy are also found.^{4–7}

The high incidence of parental consanguinity and high recurrence rate among sibs has long suggested that BSCL is usually an autosomal recessive condition. Studies in Norwegian patients suggested a founder effect traced back to the fourteenth century.⁸ An initial genome wide scan performed in a multi-ethnic set of consanguineous families failed to identify a locus displaying significant homozygosity by descent.⁹ Subsequently, Garg *et al*¹⁰ identified a gene on chromosome 9q34 (*BSCL1*) that was linked to the disease in 17 of 19 families, mostly of African-American origin. Magré *et al*¹¹ reported that this locus was unlikely to be responsible for the majority of affected families in other ethnic groups and, using extensive family material from Norway and the Lebanon, identified a second locus on 11q13 (*BSCL2*).¹¹ In the 11q13 linked families, multiple different homozygous and compound heterozygous mutations were found in a previously undescribed gene, termed *seipin*.

The *seipin* gene encodes a protein of unknown function with little homology to known proteins. Surprisingly, this gene is

widely expressed, with particularly high expression levels in the brain. Intellectual impairment has frequently been mentioned in more than 100 published clinical (case) reports of BSCL, but has often been attributed to factors such as the non-specific effects of chronic illness in childhood and/or the adverse social consequences of the very abnormal appearance of these children. The large number of affected subjects in the international consortium, which collaborated to identify the *BSCL2* gene, provided us with an opportunity to examine the genotype/phenotype relationship in this disease.

MATERIALS AND METHODS

Description of cohort

All studies were performed with the informed consent of the subjects and/or their recognised guardians and with the formal approval of local institutional ethics committees. Patients included in the cohort presented with lipodystrophy, hyperlipidaemia (in the vast majority of cases, hypertriglyceridaemia), hepatomegaly, acromegaly features, and insulin resistance. Seventy affected subjects from 44 independent pedigrees were studied. The families originated from a wide range of ethnicities and geographical locations (Appendix 2).

Genotyping

BSCL families were classified as *BSCL1*, *BSCL2*, or *BSCLX* on the basis of mutational analysis and haplotype analysis. All subjects classified as *BSCL2* had homozygous or compound heterozygous mutations in the *seipin* gene. Subjects were classified as *BSCL1* if there was evidence for cosegregation with chromosome 9q, but not with 11q. Two families showed evidence against cosegregation with either 9q or 11q and were termed *BSCLX*. One family was unclassifiable as no *seipin*

Table 1 Clinical features of patients with lipodystrophy. Group 1 (BSCL2) was defined by the presence of mutations in the *seipin* gene, while group 2 contains all other types of BSCL. Putative linkage on chromosome 9 defined group 3 (BSCL1). Nominal variables are expressed as prevalence rate (%) and were analysed by chi-square test. Numerical variables are expressed as mean (SEM) and significance was calculated by Mann-Whitney test. Statistical significance was considered at $p < 0.05$ comparing group 1 with groups 2 or 3.

	Group 1 BSCL2		Group 2 BSCL1/X/non-informative		Group 3 BSCL1	p 1/2	P 1/2
	(Index) (n=24)	Relatives G1 (n=21)	(Index) (n=20)	Relatives G2 (n=5)	(Index) (n=17)		
Gender (F/M)	10/14	8/13	15/5	3/2	13/4	0.02	0.02
Mean age (y)	20.8 (2.6)	18.5 (3.3)	20.3 (2.8)	24.0 (4.6)	21.6 (2.8)	NS	NS
Death (%)	16.7	14.3	10.0	0.0	0.0	NS	NS
Age at death (y)	26.2 (4.7)	20.7 (10.2)	4.5	–	–	NS	NS
Congenital lipodystrophy (%)	75.0	85.0	52.6	100.0	56.2	NS	NS
Age of delayed lipodystrophy (mth)	3.0 (1.6)	1.9 (1.8)	29.5 (17.5)	–	34.3 (20.6)	NS	NS
Average birth weight (g)	3022 (150)	3076 (120)	2845 (130)	2910 (340)	3007 (115)	NS	NS
Defect in birth weight (10th)	23.8	33.3	27.7	50.0	12.5	NS	NS
Diabetes mellitus (%)	45.8	23.8	35.0	0.0	35.3	NS	NS
Age of onset of diabetes (y)	14.7 ± 1.9	24.8 ± 2.7	14.3 ± 3.5	–	16.3 ± 3.4	NS	NS
Hypertrophic cardiomyopathy (%)	33.3	14.2	25.0	0.0	23.5	NS	NS
Bone cysts (%)	8.3	4.8	25.0	40.0	23.5	NS	NS
Intellectual impairment (%)	75.0	85.0	15.0	0.0	11.8	0.0001	0.0001
Parental consanguinity (%)	70.8	ND	45.0	ND	52.9	NS	NS
African ancestry (%)	0.0	ND	35.0	ND	58.2	0.005	0.002

ND, Not determined. NS, Not statistically significant.

mutation was found and haplotype analysis at chromosome 9 was uninformative.

Phenotypic evaluation

Each participant was evaluated by history, physical examination, and review of medical records. Participating investigators completed a standardised 98 item questionnaire. A formal assessment of intelligence quotient (IQ) was available only for a minority. Therefore, a simplified scale, based largely on educational performance, was used to assess whether there was mild, moderate, or severe intellectual impairment (Appendix 1).

Statistical analysis

From all 70 affected subjects, we considered only index cases (n=44) for statistical analysis while the remaining patients (n=26) were included in a group of relatives. Patients genotyped as BSCL2 (n=24) displayed evidence for mutations in the *seipin* gene on chromosome 11q. All other index cases with lipodystrophy (n=20) were considered as a unique group (BSCLs). Evidence provided for cosegregation with chromosome 9q allowed further classification as BSCL1 (n=17). Index patients lacking evidence for cosegregation with chromosomes 11q or 9q were classified as BSCLX (n=3). One patient was unclassifiable and, as such, uninformative. Congenital lipodystrophy (presence of lipodystrophy at birth) was introduced as a nominal (bimodal) variable whereas delayed lipodystrophy was introduced as a numerical variable as a function of age (month). Similarly, diabetes mellitus was introduced as a nominal variable and age of onset, when specified, was considered as a numerical variable (years). Birth weight was considered as a numerical variable (g). However, hypotheses were tested for the presence of defect in birth weight and, in this case, nominal variables were considered as defects defined under the 25th or 10th centile. Mental impairment was introduced either as a bimodal nominal variable or as a scoring index with four levels (no defect, mild, moderate, and severe). Ethnic origin (African/white) and consanguinity were introduced as bimodal nominal variables. Nominal and numerical variables were analysed with chi-square and non-parametric Mann-Whitney tests, respectively. Statistically significant values were considered at $p < 0.05$. For the multivariate analysis, logistic regression was performed with a plurimodal equation as provided by StatView 5.0 software (Abacus Concepts, Berkeley, CA). Significant values

($p < 0.1$) were introduced into the model equation, except data for genotyping (used for classification of groups). In the first step we compared the BSCL2 group against all other lipodystrophies but distinct groups of BSCL1 and BSCLX were also tested. Data in tables are expressed as prevalence rates (%) or as mean (SEM) in different groups of index cases and relatives.

RESULTS

Genotype

The molecular genetic analyses established that, of the total 44 families, the majority (24 families) represented BSCL2 (45 affected subjects). BSCL1 was the next largest subgroup with 17 families (21 affected subjects). One family was classified as BSCLX (one affected subject) and two families (three subjects) were unclassifiable. Appendix 2 shows the ethnic origin (country) of families within the three subtypes. Consistent with the previous findings of Garg *et al.*,¹⁰ all families of African ancestry were in the BSCL1 group. *Seipin* mutations (BSCL2) were found in predominantly white subjects from Lebanon, Portugal, and Norway. The two BSCLX families as well as the non-informative family were of white European extraction. The prevalence of parental consanguinity was greater, but not significantly so, in BSCL2 than in BSCL1 or other types of lipodystrophy (table 1).

Statistically significant differences were found when the BSCL2 group was compared to the putative BSCL1 genotype. The clinical profile of relatives of the two major BSCL types was very similar to the index cases. Ethnic differences were evident since 100% of BSCL2 cases were white, compared to only 65% in other lipodystrophies ($p < 0.0016$). Interestingly, all (35%) subjects of African ancestry were genotyped as putative BSCL1. Further detailed study of ethnic origin indicated that, among BSCL2 subjects, 24% were from Lebanon and 56% were of Portuguese origin ($p < 0.005$).

Phenotype

Major phenotypic features of all three subgroups taking into account index cases (used for statistical analysis) and relatives are summarised in table 1. Several major features appeared to characterise the BSCL2 group either compared to all cases of lipodystrophy or particularly to BSCL1 index cases. Thus, BSCL2 was characterised by a greater number of affected males than females, white ancestry, and highly significantly

Table 2 *Seipin* mutations in mentally retarded subjects and non-mentally retarded subjects

Mutation		No of patients n=45	Ethnic background	MR
Nonsense 32/45 (71%)	F63fsX75	2	Norwegian	Yes
	F101fsX111	1	Italian (Sicily)	Yes
	F105fsX111	21	Portuguese-Amazonian	No
	F108fsX113	2	Lebanese	Yes
			Portuguese (SA)	Yes
	R138X	5	Portuguese (SA)	No
	F213fsX231	1	Portuguese	Yes
			Indian	No
Deletion 2/45 (4.5%)	del E4-E6	2	Portuguese	Yes
Splice site 2/45 (4.5%)	639/+1G-A	1	Turkish	Yes
	deltaV99-S146			
	1016/-3C>G	1	Turkish	Yes.
	F224deltaY225-Q271fsX288			Severe developmental delay
Missense 5/45 (11%)	Ala112 Pro	5	Norwegian	Yes
Compound heterozygosity 4/45 (8.8%)	Ala 212 Pro/F 63 fsX 75	1	Norwegian	Yes
	F105fsX112/deletion E4-E6	1	Portuguese (France)	Yes
	F63 fsX75/1015/+5G→A or splice	2	British	No
	(F224fsX225)			

more frequent intellectual impairment. These features were also found when one considered all studied cases (index and relatives) of BSCL2 (n=45) compared to other lipodystrophies (n=25). Thus, 60% of BSCL2 patients were males compared to only 28% of other lipodystrophy cases.

Lipodystrophy and metabolic dysregulation

The majority of subjects were noted to have complete lipodystrophy either at or close to the time of birth. The prevalence of congenital onset of lipodystrophy was 79.5% in BSCL2 compared to 61% in other cases. Four subjects had a reliable history of a delayed onset of lipodystrophy and all of these were in the BSCL1 group. Although the congenital lipodystrophy was not significantly different among groups, it should be noted that the occurrence of lipodystrophy within the first year of life was 91% in BSCL2 patients compared to only 83% in other types of BSCL. There was a history of relative sparing of the face from the lipodystrophy in BSCL1 subjects of African origin, at least during the first decade of life. Hypertriglyceridaemia was reported in all subjects with marked variation in severity, even within individual families (for example, between 3 and 90 g/dl). The prevalence of diabetes mellitus was similar between BSCL2 (35.5%), BSCL1 (28.6%), and BSCLX (25%). The mean age at onset of clinical diabetes was also similar in BSCL2 and BSCL1 (17.9 (SD 1.9) and 16.3 (SD 3.4) years, respectively), whereas the age of onset for the single BSCLX patient with diabetes was 2 years. Hepatomegaly was noted in all subjects in this study.

Premature mortality

Seven of the 45 (15.5%) subjects in the BSCL2 group died prematurely between 14 and 35 years. Three of these deaths were the result of cardiac failure, two of renal failure, one of hepatic failure, and one of unknown cause. The non-informative subject died at the age of 19 months of cardiac failure. In contrast, all affected subjects in the BSCL1 pedigrees are still alive. Although this provides some evidence for a possible locus specific effect on premature mortality, the difference between BSCL2 and BSCL1/X/non-informative subgroups was not statistically significant.

Other clinical features

Skeletal muscular hypertrophy was reported in all affected subjects. The reported prevalence of hypertrophic cardiomyopathy was similar in BSCL1 (19%), BSCL2 (22%), and BSCLX (25%, including the non-informative patient), although the ability to make this diagnosis is highly dependent on access to

echocardiography. While the documentation of bone cysts is dependent on patients having undergone appropriate radiological examination, it is worth noting that the frequency of bone cysts was increased in BSCL1 index patients (23.5%) compared to BSCL2 (8.3%).

Intellectual impairment

Questionnaire based information regarding intellectual performance was available for BSCL1, BSCL2, and BSCLX subjects. The reported prevalence of mild or moderate intellectual impairment was remarkably high in the BSCL2 group with 78% of subjects being affected (table 2). In contrast, only 10% of the BSCL1 subgroup and the non-informative subject were reported to show evidence of intellectual impairment. The intellectual impairment found in BSCL2 was mild in 23 subjects, moderate in 11, and severe in one subject. Of the two affected BSCL1 subjects, one was mildly and the other moderately intellectually impaired. Table 2 indicates the spectrum of mutations in BSCL2 patients grouped according to the presence or absence of intellectual impairment. There was no obvious genotype-phenotype relationship. Missense, nonsense, and frameshift mutations were equally distributed between the intellectually impaired and the normal intelligence groups. Moreover, the same nonsense mutation (F108fsX113) was observed in two unrelated subjects, of whom one was mentally retarded and the other of normal intelligence.

When multivariate analysis was performed by logistic regression none of the aforementioned clinical features appeared significant, except intellectual impairment. Thus, intellectual impairment as a bimodal variable was associated with BSCL2 with $p < 0.0001$, OR 17.0, CI 3.6 to 79.0 with respect to all other types of lipodystrophies. If one compares the BSCL2 group with the BSCL1 group, again intellectual impairment appeared strongly associated with the BSCL2 genotype ($p < 0.0001$, OR 23.5, CI 3.9 to 128.3).

DISCUSSION

We have undertaken the most comprehensive study reported to date of genotype/phenotype relationships in inherited human syndromes of congenital generalised lipodystrophy (CGL) involving 70 affected subjects from 44 independent pedigrees (fig 1). Clinically relevant findings resulting from these studies include (1) the confirmation of the existence of at least one additional CGL locus other than those on chromosomes 9 and 11, (2) the confirmation of ethnic differences in

Figure 1 Family pedigrees of all participating families grouped by aetiology. For BSCL1 families, evidence for cosegregation with 9q34 is provided. For BSCL2, the nature of the mutation identified is indicated under the pedigree. For BSCLX, evidence against cosegregation with the 9q34 locus is also provided.

the relative frequency of *BSCL1* and *BSCL2* gene defects, and (3) the establishment of certain key phenotypic differences, in particular concerning intellectual impairment, between the three subtypes of CGL.

Two Europic families in our cohort had no detectable *seipin* mutations and also showed evidence against the involvement of a locus at 9q34. Taken together with the data of Heathcote

*et al*¹² in Omani pedigrees, this strongly suggests that there must be at least one additional locus that causes BSCL. For the moment, we suggest that this be referred to as *BSCLX*.

Among the 44 families studied, BSCL1 patients originated primarily from sub-Saharan Africa and Maghreb countries (Morocco, Algeria, Tunisia) with only a minority coming from Middle Eastern countries (for example, Turkey) and northern

Figure 1 continued

Europe. This is consistent with the results of Garg *et al*,¹⁰ who found cosegregation of the disease with chromosome 9q34 markers in the great majority of their families of African-American origin. In contrast, a large number of different *seipin* (BSCL2) mutations were found in subjects of varying

white ethnicities including Norway, United Kingdom, and Mediterranean countries, as well as Middle Eastern Arabs. Interestingly, patients from Portugal and its ancient colonies accounted for a significant proportion of subjects, suggesting a spread from the 15th century onwards.¹³ Notably, no

Figure 1 continued

subjects of African ancestry had *seipin* mutations. The two BSCLX families as well as the non-informative patient were of European origin.

The higher prevalence of recorded inbreeding in BSCL2 families than in BSCL1 families cannot easily be evaluated as long as the various local inbreeding patterns are not known. For example, chance inbreeding has been verified in one Norwegian BSCL2 genetic compound family. Taking into account also the different ethnic origins, the simplest explanation is that the high African origin of BSCL1 is the result of a single, very ancient mutation having reached an appreciable spread and allele frequency.

Phenotypic evaluation was undertaken by multiple independent physicians in several countries and patients were not examined by a single observer. Since a standardised formal assessment of intellectual capabilities was difficult to establish

because of highly variable attendance at school and the wide range of ethnic and racial background, it was decided to evaluate the degree of mental impairment using a grow scale measuring social adaptation and school performance. This classification allows some overlap between severe and moderate mental retardation.

Notwithstanding these caveats, the application of standardised questionnaires across these multiple centres has allowed us to attempt to establish, with reasonable confidence, whether there are any major and consistent phenotypic differences between the genetic subtypes of this disease. Lipoatrophy was present at birth and complete in all BSCL2 and BSCLX subjects. In contrast, a subset of BSCL1 subjects had evidence for lipoatrophy that was either delayed, incomplete, or both. For those subjects in whom the lipoatrophy was incomplete, the face tended to be spared. Thus, BSCL1

Figure 1 continued

may show phenotypic overlap with at least two other forms of lipodystrophy that have previously been considered distinct. Acquired total lipodystrophy (Lawrence syndrome) is generally considered to be a sporadic, non-inherited condition in which total lipodystrophy, of presumed autoimmune origin, occurs often after an infectious illness.¹⁴ Our findings suggest that some cases of BSCL1 in which the onset of the lipodystrophy is delayed and are of sporadic occurrence may be labelled incorrectly as having Lawrence syndrome. The distinction is made more complex by the fact that there are no diagnostic tests for Lawrence syndrome, although the

lipodystrophy of this syndrome does tend to appear more suddenly than in the BSCL1 cases. Familial partial lipodystrophy of the Dunnigan-Köberling type (FPLD, OMIM 151660) is an autosomal dominant, face sparing lipodystrophy caused by a particular cluster of mutations in the nuclear envelope proteins lamins A/C.¹⁵⁻¹⁹ However, these patients have a distinctive cushingoid appearance of the face that makes differential diagnosis easy.

The relative effects of the BSCL subtypes on premature mortality are difficult to quantify given factors such as incomplete ascertainment in families and possible referral bias. However, it is notable that while none of the BSCL1 patients included in our study died before the age of 40 (but family history death of affected subjects in early adulthood was positive in some cases), premature death was observed in seven patients from the BSCL2 subgroup. The reasons for this difference are not obvious as the prevalence of major metabolic derangements (for example, diabetes, hyperlipidaemia, and hepatic dysfunction) were similar between the groups. However, it is notable that three of the seven premature deaths in the BSCL2 group were attributed to cardiac failure, including one during pregnancy.

As these diseases are clearly autosomal recessive it was surprising that we found an excess of females in the BSCL1 group and an excess of males in the BSCL2 group. In the case of BSCL1, this is most likely to represent ascertainment bias, as the appearance of lipodystrophy is often more striking in females leading to earlier referral and diagnosis. However, this would not explain the opposite sex ratio found in the BSCL2 group. Within that group, affected female subjects tended to have a more severe phenotype than males (personal observations) and this subtype is generally more lethal than BSCL1. It is possible, therefore, that earlier mortality in affected females has resulted in a systematic bias towards the referral of surviving males to the consortium.

Interestingly, radiological evidence of bone cysts was observed in 10 patients (five with BSCL1, three with BSCL2, and two with BSCLX). A syndrome of cystic angiomas of bones and congenital lipodystrophy (OMIM 272500) has been reported by Brunzell *et al*²⁰ in 1968 and considered to be an entity separate from BSCL. Our findings suggest that this may not be a discrete entity.

The most remarkable difference between the groups was the prevalence of intellectual impairment. While it was not possible to undertake thorough and systematic quantitative assessment of IQ, it is notable that while 78% of BSCL2

Appendix 1 Criteria for the classification of intellectual impairment

Degree of mental retardation	Preschool years (0–5 y)	School years (6–21 y)	Adulthood
Simplified scale			
Mild (IQ 50–70)	Is rarely distinguishable from a normal child Mildly delayed development in the sensorimotor field Mild language delay Good social skills	Able to acquire a basic level of education, but doesn't reach the competence level of the last year of primary school	Able, with special education and special training to reach a satisfactory professional and social functioning Often needs guidance and supervision in stressful circumstances
Moderate (35 < IQ < 50)	Learns language and communication Motor development reaches a satisfactory level Able to acquire satisfactory autonomy Able to be self-sufficient with moderate controls	Is only able to learn very few pedagogic milestones (Some reading, some writing and calculation, usually during adolescence)	Able to do non-qualified work, usually in a protected environment Needs supervision and guidance Is unable to work autonomously
Severe (20 < IQ < 35)	Poor motor development Little or no language Low communication skills	Able to learn a few words Able to communicate Hygiene rules and minimal behavioural habits are acquired Unable to acquire school education	Is able partially to earn his or her income when working in a protected environment and under full supervision
Profound (IQ < 20)	Very low functioning capacity Needs nursing	Mild motor development Unable to acquire personal autonomy Needs permanent care	Some motor development is observed No development of language Needs permanent care

Appendix 2 Clinical features of all subjects listed individually

Family	No	Initials	Sex	Year of birth	Weight at birth (g)	Age at referral	Reason for referral		Age at onset of lipo	Age at onset of diabetes mellitus	Intellectual impairment	Other clinical features		Age at death	Cause of death	Ethnic background	Parental consang
BSCL1																	
1*	1	FV	F	1974	3100	6 mth	Failure to thrive	3 mth	26 y	Moderate		Bone cysts, microcephaly	–	–		Belgian	Y
2*	2	WX	M	1977	2760	5 mth	Lipoatrophy	Birth	–	–		Hypertrophic cardiomyopathy	–	–		African - Caribbean	N
3*	3	ZC	F	1987	–	7 y	Diabetes	–	7 y	–			–	–		Algerian	N
4*	4	DC	F	1968	–	5 y	Weight loss	5 y	13 y	–		Pancreatitis (at 7 y)	–	–		African - mixed	N
5*	5	DiH	F	1972	–	11 y	Diabetes mellitus	2 mth	7 y	Mild		Bone cysts	–	–		African - Senegal	Y
	6	Dil	M	1984	2570	6 wk	Lipoatrophy, family history	Birth	Transient hyperglycaemia at birth	–		–	–	–		African - Senegal	Y
	7	DiN	F	1987	–	9 mth	Lipoatrophy, family history	Birth	–	–		–	–	–	–	African - Senegal	Y
6*	8	AL	F	1983	2550	2 wk	Lipoatrophy	Birth	–	–		Normal cardiac ultrasound, clitoromegaly	–	–		African - Ghana	N
7*	9	J	F	1990	–	–	–	Birth	–	–		–	–	–	–	Moroccan	Y
8*	10	Rd	M	1991	–	–	–	Birth	–	–		–	–	–	–	Turkish	Y
9*	11	HL	F	1962	–	3 y	Lipoatrophy	3 wk	–	–		Bone cysts	–	–		African-American	N
	12	HHD	F	1965	–	9 mth	Family history	Birth	–	–		Bone cysts	–	–		African-American	N
	13	HRD	M	1969	–	6 y	Family history	–	–	–		–	–	–	–	African-American	N
10*	14	M	F	1990	3300	5 mth	Lipoatrophy	Birth	–	–		–	–	–	–	Tunisian	Y
11*	15	OU	F	1964	–	23 y	Diabetes mellitus	23 y	23 y	–		Hypertrophic cardiomyopathy	–	–		Algerian	Y
12*	16	SMG	F	1994	–	9 mth	Gigantism, lipoatrophy	Birth	–	–		Clitoromegaly	–	–		German (Brazil)	Y
13*	17	VV	F	1975	3000	4 y	Gigantism, precocious puberty	Birth	–	–		Bone cysts, goitre, macrocephaly	–	–		African-Indian (Caribbean)	N
14*	18	BJ	M	1993	2900	3 mth	Abdominal pain, hepatomegaly	Birth	–	–		Hypertrophic cardiomyopathy	–	–		African-Nigerian (France)	N
15*	19	EF	M	1996	3440	3 mth	Facial dysmorphism, hepatomegaly	3 mth	–	–		Hypertrophic cardiomyopathy	–	–		Turkish	Y
16*	20	PH	F	1970	–	22 y	Diabetes	17 y	22 y	–		–	–	–	–	Dutch (South Africa)	N
17*	21	ZA	F	–	–	6 mth	Lipoatrophy	Birth	–	–		–	–	–	–	Moroccan (France)	Y
Family	No	Seipin amino acid change	Type of seipin mutation	Initials	Sex	Year of birth	Weight at birth (g)	Age at referral	Reason for referral	Age at onset of lipo	Age at onset of diabetes mellitus	Intellectual impairment	Other clinical features	Age at death	Cause of death	Ethnic background	Parental consang
BSCL2																	
1	1	R138X/ R138X	Non	MALF	F	1965	Normal	24 y	Diabetes, acromegaloid features	6 mth	24 y /No	Mild	Hypertrophic cardiomyopathy	–	–	Portuguese	Y
	2	R138X/ R138X	Non	ML	F	1967	Normal	20 y	Diabetes	Infancy	15 y	Mild	Multinodular goitre	–	–	Portuguese	Y
2	3	R138X/ R138X	Non	HSDM	F	1980	–	15 y	Diabetes	Infancy	15 y	Mild	–	–	–	Portuguese	Y
3	4	F63fsX75/ F63fsX75	Ins-del	DD	M	1982	2900	Birth	Lipoatrophy	Birth	–	Moderate	Hypertrophic cardiomyopathy	–	–	Italian (Sicily)	Y
4	5	DV99-S146/ DV99-S146	Splice site	AS	M	1981	3120	3 mth	Rickets	Birth	–	Moderate	Persistent mullerian ducts, vitamin D resistant rickets	14 y	Heart failure	Turkish	Y
5	6	R138X/ R138X	Non	FJ	M	1992	3040	6 mth	Dysmorphism	First year	–	No (IQ 101)	Waddling gait, prognathism, hypertrophic cardiomyopathy	–	–	Portuguese (France)	Y
	7	R138X/ R138X	Non	FS	M	1996	3490	2 mth	Family history	–	–	No (IQ 101)	Waddling gait, surgical cure of macroglossia, hypertrophic cardiomyopathy	–	–	Portuguese (France)	Y

Appendix 2 continued

Family	No	Seipin amino acid change	Type of seipin mutation	Initials	Sex	Year of birth	Weight at birth	Age at referral	Reason for referral	Age at onset of lipo	Age at onset of diabetes mellitus	Intellectual impairment	Other clinical features	Age at death	Cause of death	Ethnic background	Parental consang
6	8	Ala212Pro/Ala212Pro	Miss	AE	F	1952	2000 at 33w	Birth	Lipoatrophy	Birth	12 y	Mild	Hypertrophic cardiomyopathy	32 y	Post-intoxication renal failure, pneumonia	Norwegian	N
	9	Ala212Pro/Ala212Pro	Miss	ME	M	1957	4000	1 y	Developmental delay	Birth	–	–	Angina pectoris, bone cysts	–	–	Norwegian	N
7	10	F63fsX75/F63fsX75	Ins-del	KS	M	1957	2500	–	Lipoatrophy	Birth	14 y	Moderate	Hypertrophic cardiomyopathy, bone cysts	35 y	Renal failure	Norwegian	N
8	11	Ala212Pro/Ala212Pro	Miss	LSN	M	1959	4100	–	–	Birth	–	Moderate	Hypertrophic cardiomyopathy	–	–	Norwegian	N
9	12	Ala212Pro/Ala212Pro	Miss	IT	F	1961	3000	5 mth	Enlarged tonsils	Birth	–	–	Hypertrophic cardiomyopathy	24 y	Heart failure during pregnancy	Norwegian	N
	13	Ala212Pro/Ala212Pro	Miss	TH	M	1966	4030	2.5 mth	Lipoatrophy	Birth	–	Moderate	Hypertrophic cardiomyopathy	–	–	Norwegian	N
10	14	Ala212Pro/F63fsX75	Miss/ins-del	OE	F	1973	3550	2 mth	Lipoatrophy	Birth	–	Mild	Bone cysts	–	–	Norwegian	Y
11	15	F105fsX111/F105fsX111	Del	Tba	M	1991	3000	3 y	Lipoatrophy	Birth	–	Moderate (IQ 40)	–	–	–	Lebanese	Y
	16	F105fsX111/F105fsX111	Del	TCh	M	1994	3000	9 mth	Lipoatrophy	Birth	–	Mild	–	–	–	Lebanese	Y
	17	F105fsX111/F105fsX111	Del	Tfa	M	1968	2800	31 y	Family history	Birth	31 y	Mild	–	–	–	Lebanese	Y
12	18	F105fsX111/F105fsX111	Del	EIA	M	1989	2200	7 mth	Lipoatrophy	Birth	–	Mild	–	–	–	Lebanese	Y
	19	F105fsX111/F105fsX111	Del	EIAN	M	1995	–	6 mth	Lipoatrophy	Birth	–	Mild	–	–	–	Lebanese	Y
13	20	F105fsX111/F105fsX111	Del	SJ	F	1993	2250	3 mth	Lipoatrophy	Birth	–	Moderate	–	–	–	Lebanese	Y
	21	F105fsX111/F105fsX111	Del	SR	F	1999	2400	2 mth	Lipoatrophy	Birth	–	–	–	4 mth	Sepsis	Lebanese	Y
14	22	F105fsX111/F105fsX111	Del	Jfa	F	1987	3000	2 y	–	Birth	8.5 y	Mild	–	–	–	Lebanese	Y
15	23	F105fsX111/F105fsX111	Del	SSi	F	1997	3120	Birth	Lipoatrophy & family history	Birth	–	Moderate	Precocious puberty at 4 y, hirsutism, liver cirrhosis, hypertrophic cardiomyopathy	–	–	Lebanese (USA)	Y
	24	F105fsX111/F105fsX111	Del	SSe	M	1996	3330	1 mth	Lipoatrophy	Birth	–	Moderate	Liver cirrhosis, hypertrophic cardiomyopathy	–	–	Lebanese (USA)	Y
16	25	F105fsX111/F105fsX111	Del	FE	M	1987	3500	9 mth	Hepato-megaly	Birth	12 y	Mild	–	–	–	Lebanese	Y
	26	F105fsX111/F105fsX111	Del	AS	F	1969	–	–	–	Birth	24 y	Mild	–	31 y	Liver cirrhosis	Lebanese	Y
	27	F105fsX111/F105fsX111	Del	AG	F	1972	–	–	–	Birth	26 y	Mild	–	–	–	Lebanese	Y
	28	F105fsX111/F105fsX111	Del	HS	M	1989	2700	2 mth	–	Birth	–	Mild	–	–	–	Lebanese	Y
	29	F105fsX111/F105fsX111	Del	HP	M	1992	3000	2 mth	–	1 mth	–	Mild	–	–	–	Lebanese	Y
	30	F105fsX111/F105fsX111	Del	GR	M	2000	3200	1 mth	–	–	–	Mild	–	–	–	Lebanese	Y
	31	F105fsX111/F105fsX111	Del	GG	M	1966	3100	33 y	Family history	Birth	–	Mild	–	31 y	Heart failure	Lebanese	Y
	32	F105fsX111/F105fsX111	Del	Gba	F	1968	3000	31 y	Family history	Birth	28 y	Mild	–	–	–	Lebanese	Y

Appendix 2 continued

Family	No	Seipin amino acid change	Type of seipin mutation	Initials	Sex	Year of birth	Weight at birth (g)	Age at referral	Reason for referral	Age at onset of lipo	Age at onset of diabetes mellitus	Intellectual impairment	Other clinical features	Age at death	Cause of death	Ethnic background	Parental consang
	33	F105fsX111/ F105fsX111	Del	Gbc	M	1971	3200	28 y	Family history	Birth	–	Mild	–	–	–	Lebanese	Y
	34	F105fsX111/ F105fsX111	Del	She	F	1966	2500	9 mth	Lipoatrophy	Birth	–	Mild	–	–	–	Lebanese	Y
	35	F105fsX111/ F105fsX111	Del	SEI	M	1998	2500	1 mth	Lipoatrophy & family history	1 mth	–	Mild	–	–	–	Lebanese	Y
17	36	F213fsX231/ F213fsX231	Del	SN	M	1985	–	4 mth	Organo-megaly	Birth	15 y	–	Normal cardiac ultrasound	–	–	Indian (UK)	Y
18	37	F101 fsX111/ F101 fsX111	Ins	MDSM	F	1960	Normal	29 y	Diabetes	3 mth	29 y	–	–	–	–	Portuguese (Brazil)	Y
19	38	F108fsX113/ F108fsX113	Ins	DSF	M	1975	3320	6 y	Diabetes	1 y	6 y	Moderate	–	–	–	Portuguese (South Africa)	Y
20	39	F108fsX113/ F108fsX113	Ins	GD	F	1981	–	2 mth	Lipoatrophy	Birth	14 y	–	–	–	–	Portuguese (South Africa)	N
21	40	Del exons 4–6	Del	HM twin 41	M	1990	2320 at 37 wk	5 mth	Failure to thrive	Birth	–	Mild	IUGR, polyhydramnios	–	–	Portuguese (France)	Y
	41	Del exons 4–6	Del	HD twin 40	M	1990	2050 at 37 wk	5 mth	Failure to thrive	Birth	–	Mild	IUGR, polyhydramnios	–	–	Portuguese (France)	Y
22	42	F105fsX112/ del E5-6	Del/del	RA	F	1970	–	Birth	Lipoatrophy	3–6 mth	13 y	Moderate	–	–	–	Portuguese (France)	N
23	43	F224ΔY225- Q271fsX288	Splice	EL	M	–	2000	Birth	Lipoatrophy	Birth	–	Severe psychomotor delay, pyramidal signs	–	–	–	Turkish	Y
24	44	F63fsX75/ F224fsX225	Ins-del/ splice	WM	M	1985	4700	6 mth	Lipoatrophy	Birth	–	–	–	–	–	British	N
	45	F63fsX75/ F224fsX225	Ins-del/ splice	WF	F	1989	3800	Birth	Family history	Birth	–	–	–	–	–	British	N
Family	No	Initials	Sex	Year of birth	Weight at birth (g)	Age at referral	Reason for referral	Age at onset of lipo	Age at onset of diabetes mellitus	Intellectual impairment	Other clinical features			Age at death	Cause of death	Ethnic background	Parental consang
BSCLX																	
1	1	HP	F	1980	2950	1 y	Failure to thrive	1 y	2 y	–	Liver cirrhosis, thick, curly hair			7.5 y	Liver failure	Czech Republic	N
2	2	RJ	F	1985	3250	Birth	Family history	Birth	–	–	Bone cysts			–	–	Belgian	N
	3	RK	F	1983	2350	13 mth	Failure to thrive	5 wk	–	–	Bone cysts			–	–	Belgian	N
Family	No	Initials	Sex	Year of birth	Weight at birth (g)	Age at referral	Reason for referral	Age at onset of lipo	Age at onset of diabetes mellitus	Intellectual impairment	Other clinical features			Age at death	Cause of death	Ethnic background	Parent consang
Non-informative																	
1	1	SV	M	1992	2100 at 37 wk	Birth	IUGR, hepatosplenomegaly	Birth	–	Moderate	Liver steatosis, hypertrophic cardiomyopathy			19 mth	Cardiac failure	Czech Republic	

subjects were reported as having some degree of intellectual impairment, this was true of only 10% of the BSCL1 subjects. Although this difference might be explained by the expression of seipin in the brain, such a tissue specific expression may also be independent from any type of mental retardation (Wolfram syndrome, Duchenne muscular dystrophy, merosin positive congenital muscular dystrophy) and does not therefore provide a definitive clue to its pathophysiology. The normal brain MRI reported in mentally retarded BSCL2 patients is also remarkable.

No clear correlation was apparent between the site and type of seipin mutation and intellectual impairment. Furthermore, affected subjects from different families, but harbouring the same mutation, were found to be discordant in terms of mental impairment in some instances. Clearly, given the complex and qualitative nature of cognitive functions, it is not surprising that its expression may be subject to multiple genetic and environmental modifiers.

In summary, BSCL1 is a form of congenital lipodystrophy, most prevalent among subjects of African origin, which appears to be a milder disease than BSCL2. This is manifested by its more frequent presentation as a delayed or partial lipodystrophy, a lower occurrence of premature death, and a markedly lower incidence of intellectual impairment. These observations suggest that the causative gene in BSCL1, yet to be described, may turn out to have more restricted functions and/or a more limited expression pattern than seipin. Our findings emphasize the importance of attempting to make a molecular diagnosis of the syndromes of congenital lipodystrophy as this may aid both prognostication and genetic counselling.

ACKNOWLEDGEMENTS

We thank the patients, their families, and the referring physicians for their participation; A Lemainque, P Hilbert, L Bodson, F Cavallin, A Fauré, V La Villa, and S Pavék for genotyping; and D Recan and her colleagues for immortalising the lymphoblasts. We thank the rare diseases programme of the European Commission (Directorate-General XII) for its support and L'Aide aux Jeunes diabétiques, l'Association Française contre les Myopathies (AFM), the Anders Jahres Fund, Assistance Publique-Hôpitaux de Paris, INSERM, Medinnova, and Ministère de la Recherche for their constant encouragement and financial support. We also want to thank Christian Melot and Maurice Jottrand for their help in statistical analysis of the data.

NOTE ADDED IN PROOF

After submission of this paper, identification of the BSCL1 gene was published by Agarwal A, et al (*Nat Genet* 2002;31:21–3). The AGPAT2 gene encodes a key enzyme in the biosynthesis of triacylglycerol. This is consistent with the limited expression pattern hypothesised in our conclusion.

Authors' affiliations

L Van Maldergem, C S Albott, Centre de Génétique Humaine, Institut de Pathologie et de Génétique, Lovreval, Belgium
J Magré, J Capeau, INSERM U 402, Faculté de Médecine Saint Antoine, Paris, France
T E Khallouf, Service de Pédiatrie, Hôtel-Dieu de France, Beirut, Lebanon
T Gedde-Dahl Jr, Department of Dermatology and Institute of Forensic Medicine, Rikshospitalet, Oslo, Norway
M Delépine, M Lathrop, Centre National de Génotypage, Evry, France
O Trygstad, Department of Paediatrics, Rikshospitalet, Oslo, Norway
E Seemanova, Department of Clinical Genetics, Charles University, Prague, Czech Republic
T Stephenson, Department of Child Health, University Hospital, Nottingham, UK
F Bonnici, Paediatric Endocrine Services, University of Cape Town, Cape, South Africa
V R Panz, Department of Medicine, University of the Witwatersrand, Johannesburg, South Africa
J-L Medina, Department of Endocrinology, Hospital S João, Porto, Portugal

P Bogalho, Unidade de Endocrinologia, Hospital Curry Cabral, Lisbon, Portugal
F Huet, Service de Pédiatrie, Hôpital d'Enfants du Bocage, Dijon, France
S Savasta, Clinica Pediatrica, Policlinico S Matteo, Università di Pavia, Pavia, Italy
A Verloes, Centre de Génétique Humaine, CHU Sart-Tilman, Liège, Belgium
J-J Robert, Service de Diabétologie Infantile, Hôpital Necker, Paris, France
H Loret (deceased), Service de Pédiatrie, CHU, Fort-de-France, Guadeloupe
M de Kerdanet, Unité d'Endocrinologie Pédiatrique, CHU Rennes, France
V R Panz, Department of Medicine, University of the Witwatersrand, Johannesburg, South Africa
J-L Medina, Department of Endocrinology, Hospital S João, Porto, Portugal
P Bogalho, Unidade de Endocrinologia, Hospital Curry Cabral, Lisbon, Portugal
F Huet, Service de Pédiatrie, Hôpital d'Enfants du Bocage, Dijon, France
S Savasta, Clinica Pediatrica, Policlinico S Matteo, Università di Pavia, Pavia, Italy
A Verloes, Centre de Génétique Humaine, CHU Sart-Tilman, Liège, Belgium
J-J Robert, Service de Diabétologie Infantile, Hôpital Necker, Paris, France
H Loret (deceased), Service de Pédiatrie, CHU, Fort-de-France, Guadeloupe
M de Kerdanet, Unité d'Endocrinologie Pédiatrique, CHU Rennes, France

REFERENCES

- Berardinelli W. An undiagnosed endocrinometabolic syndrome: report of two cases. *J Clin Endocrinol Metab* 1954;14:193-204.
- Seip M. Lipodystrophy and gigantism with associated endocrine manifestations: a new diencephalic syndrome? *Acta Paediatr Scand* 1959;48:555-74.
- Seip M, Trygstad O. Generalised lipodystrophy, congenital and acquired (lipodystrophy). *Acta Paediatr Suppl* 1996;413:2-28.
- Moller DE, O'Rahilly S. Syndromes of severe insulin resistance: clinical and pathophysiological features. In: Moller DE, ed. *Insulin resistance*. New York: Wiley and Sons, 1993:49-81.
- Taylor SI, Arioglu E. Genetically defined forms of diabetes in children. *J Clin Endocrinol Metab* 1999;84:4390-6.
- Garg A. Lipodystrophies. *Am J Med* 2000;108:143-52.
- Reitman ML, Arioglu E, Gavrilova O, Taylor SI. Lipodystrophy revisited. *Trends Endocrinol Metab* 2000;11:410-16.
- Gedde-Dahl T Jr, Trygstad O, Van Maldergem L, Magré J, van der Hagen C B, Olaisen B, Stenersen M, Mevag B. Genetics of the Berardinelli-Seip syndrome (congenital generalised lipodystrophy) in Norway: epidemiology and gene mapping. Berardinelli-Seip Study Group. *Acta Paediatr Suppl* 1996;413:52-8.
- Magré J, Hilbert P, Desbois-Mouton C, Vigouroux C, Fauré S, Weissenbach J, Capeau J, Bonnici A, Nivelon-Chevalier A, Gedde-Dahl T, O'Rahilly S, D'Abronzio FH, Bachy A, Van Maldergem L and the Berardinelli-Seip study group. Search for the gene(s) responsible for lipodystrophic diabetes. *Eur J Hum Genet* 1996;4:165A.
- Garg A, Wilson R, Barnes R, Arioglu E, Zaidi Z, Gurakan F, Kocak N, O'Rahilly S, Taylor SI, Patel SB, Bowcock AM. A gene for congenital lipodystrophy maps to human chromosome 9q34. *J Clin Endocrinol Metab* 1999;84:3390-4.
- Magré J, Delepine M, Khallouf E, Gedde-Dahl T Jr, Van Maldergem L, Sobel E, Papp J, Meier M, Megarbane A, Bachy A, Verloes A, D'Abronzio FH, Seemanova E, Assan R, Baudic N, Bourut C, Czernichow P, Huet F, Grigorescu F, de Kerdanet M, Lacombe D, Labrune P, Lanza M, Loret H, Matsuda F, Navarro J, Nivelon-Chevalier A, Polak M, Robert JJ, Tric P, Tubiana-Rufi N, Vigouroux C, Weissenbach J, Savasta S, Maassen JA, Trygstad O, Bogalho P, Freitas P, Medina HL, Bonnici F, Joffe BI, Loyson G, Panz VR, Raal FJ, O'Rahilly S, Stephenson T, Kahn CR, Lathrop M, Capeau J, BSCL Working Group. Identification of the gene altered in Berardinelli-Seip congenital lipodystrophy on chromosome 11q13. *Nat Genet* 2001;28:365-70.
- Heathcote K, Rajab A, Magré J, Syrris P, Besti M, Patton M, Delepine M, Lathrop M, Capeau J, Jeffery S. Molecular analysis of Berardinelli-Seip congenital lipodystrophy in Oman; evidence for multiple loci. *Diabetes* 2002;51:1291-3.
- Van Maldergem L, Da Silva H, Freitas P, D'Abronzio FH. Berardinelli-Seip syndrome: a new Portuguese disease? *Eur J Hum Genet* 1998;6:74A.
- Lawrence RD. Lipodystrophy and hepatomegaly with diabetes, lipemia, and other metabolic disturbances: a case throwing new light on the action of insulin. *Lancet* 1946;i:724-31, 773-5.
- Mounkes LC, Burke B, Stewart CL. The alpha-type lamins. Nuclear structural proteins as a focus for muscular dystrophy and cardiovascular diseases. *Trends Cardiovasc Med* 2001;11:280-5.
- Cao H, Hegele RA. Nuclear lamin A/C R482Q mutation in Canadian kindreds with Dunnigan-type familial partial lipodystrophy. *Hum Mol Genet* 2000;9:109-12.

- 17 **Shackleton S**, Lloyd DJ, Jackson SN, Evans R, Niermeijer MF, Singh BM, Schmidt H, Brabant G, Kumar S, Durrington PN, Gregory S, O'Rahilly S, Trembath RC. LMNA, encoding lamin A/C, is mutated in partial lipodystrophy. *Nat Genet* 2000;**24**:153-6.
- 18 **Speckman RA**, Garg A, Du F, Bennett L, Veile R, Arioglu E, Taylor SI, Lovett M, Bowcock AM. Mutational and haplotype analyses of families with familial partial lipodystrophy (Dunnigan variety) reveal recurrent missense mutations in the globular C-terminal domain of lamin A/C. *Am J Hum Genet* 2000;**66**:1192-8.
- 19 **Vigouroux C**, Magré J, Vantyghem MC, Bourut C, Lascols O, Shackleton S, Lloyd DJ, Guerci B, Padova G, Valensi P, Grimaldi A, Piquemal R, Touraine P, Trembath RC, Capeau J. Lamin A/C gene: sex-determined expression of mutations in Dunnigan-type familial partial lipodystrophy and absence of coding mutations in congenital and acquired generalised lipodystrophy. *Diabetes* 2000;**49**:1958-62.
- 20 **Brunzell JD**, Shankle SW, Bethune JE. Congenital generalized lipodystrophy accompanied by cystic angiomas. *Ann Intern Med* 1968;**69**:501-16.

Is your paper being cited?

CiteTrack service

CiteTrack will alert you by email whenever new content in the *Journal of Medical Genetics* or a participating journal is published that matches criteria you want to track

Topics: Tell CiteTrack which words or subjects to watch for in new content

Authors: Be alerted whenever key authors you are following publish a new paper

Articles: Know whenever a paper of interest to you is referenced by another paper

www.jmedgenet.com

PostScript

BOOK REVIEW

Molecular Cytogenetics Protocols and Applications

Methods in Molecular Biology Volume 204.
Editor Yao-Shan Fan. (Pp 424; \$135.00.)
Totowa, NJ: Humana Press. 2002. ISBN
1-58829-006-9.

This book is a very up to date manual covering the background, methodologies, and applications of molecular cytogenetics techniques. The emphasis is on the diagnostic applications of FISH in the many areas of medicine on which it impinges, including paediatrics, fetal and reproductive medicine, pathology, haematology, oncology, and, of course, medical genetics. With 27 chapters and over 60 authors, all of whom are experts in their field, this book clearly shows how far molecular cytogenetics techniques have developed over the last two decades.

The book is divided into three parts. Part 1 covers the basic concepts and techniques. The opening chapter by the editor Yao-Shan Fan provides a very helpful overview of the scope of the book and an extensive set of references for further reading. The following chapters cover probe labelling (DNA and RNA probes) and basic FISH techniques. The second part of the book is devoted to evolving techniques and applications and includes chapters on microdissection, PRINS, SKY FISH, M FISH, CGH, colour banding FISH, fibre FISH, multi-telomere FISH, fluorescence genotyping for telomeric regions, and microarray CGH. Special applications of molecular cytogenetic techniques in chromosomal disorders are covered in part 3 of the book. These include chapters on the application of FISH to the

delineation of marker chromosomes and the diagnosis of microdeletion syndromes. Other chapters cover FISH interphase nuclei screening for prenatal diagnosis including preimplantation diagnosis and fetal cells in the maternal circulation, in addition to the interphase FISH screening of routine amniotic fluid samples. This section concludes with a chapter on the application of FISH and CGH in reproductive pathology. The fourth and final section of the book covers the application of molecular cytogenetic techniques to cancer diagnosis. Chapters include the use of CGH in cancer investigations and the application of interphase FISH for the BCR/ABL rearrangements in CML and for HER2 amplification in breast cancer. Also included in this section are the interesting combined approaches, firstly of chromogenic in situ hybridisation with FISH in pathology and secondly simultaneous fluorescence immunophenotyping and FISH on tumour cells.

With any multiauthor book, there are bound to be differences in approach to the writing of individual contributions. This book has taken a surprisingly consistent approach, perhaps an illustration of good editorial control. All of the chapters have good introductory sections and are well referenced, as well as containing the authors' preferred methodologies. Each chapter also includes a comprehensive notes section (effectively tips and troubleshooting advice from the experts). However, a major problem with the book is the lack of comparison between different molecular techniques. In the preface, it is suggested that the book should help the technologist or cytogeneticist determine which procedure to perform for an informative result. I am not sure that the book provides this, as most if not all of the contributors are bound to select their own preferred methodology. For example, in the chapter on FISH screening for telomere abnormalities, an otherwise excellent chapter by Samantha Knight and Jonathan Flint, the practical application of only one of the two commercially available probe sets for this test is covered. More importantly, elsewhere in the book there is a first class chapter on interphase FISH for prenatal diagnosis of common aneuploidies by Baruch Feldman *et al.* This covers the topic extremely comprehensively and provides 89 literature citations. However,

there is no mention, as far as I could see, throughout the book of the other alternative molecular approach of quantitative fluorescence PCR for prenatal aneuploidy detection. While it may be reasonable to excuse the editor by saying that the book is not designed to cover purely molecular genetic techniques, in the same section as the prenatal FISH, there is a chapter on the molecular detection of uniparental disomy. This chapter sits rather incongruously among the others, but is in itself a useful and important topic. Another chapter which seems to have lost its place describes the BAC resource for molecular cytogenetics. This is the final chapter in the book, which appears to have been added as an afterthought. Surely this should have been in evolving techniques and applications rather than "special applications in oncology". The oncology applications would also have benefited from more chapters, for example, haematological disorders other than CML, solid tumour FISH (other than HER2), and perhaps a chapter on the screening of urine samples for bladder cancer.

Perhaps these topics are covered in the companion volume (Methods in molecular biology, volume 220. Cancer cytogenetics: methods and protocols). Another criticism is that the provision of colour plates is very variable. Some chapters are well illustrated, others less so. For example, the chapter on SKY FISH relies on black and white illustrations, whereas the MFISH chapter has glorious full colour images. Furthermore, the overall size of the book is relatively small (16 cm by 24 cm) and the size of typesetting and tight layout does not make for easy reading when compared with, for example, Rooney's "Human cytogenetics: a practical approach".

On the plus side, there is an incredibly large amount of information packed into this volume, none of it superfluous. Although other textbooks that cover FISH techniques are available, this book provides a more comprehensive, up to date, and thorough coverage of diagnostic molecular cytogenetics than any of the other books currently available.

In summary, I would recommend it as a reference source for everyone working in and interested in the exciting field of diagnostic and research molecular cytogenetics.

Lionel Willatt

NOTICE

European Human Genetics Conference 3 – 6 May 2003 ICC, Birmingham, England

Plenary sessions

- Low Penetrance Genes and Cancer Susceptibility
- Public Issues - Population DNA Banks
- Recent Developments in Neurogenetics

Symposia

- Bioinformatics
- Stem cells
- Sensory genetics
- Alternative splicing
- Cancer genetics
- Alzheimer disease
- SNPs and haplotypes
- Chromosomes in genetic disease
- Genetics and endocrine problems

Workshops

- Syndrome identification
- Cytogenetics
- Problems in counselling/ethics
- Genotyping and mutation detection arrays— practical problems
- Quality control
- Prenatal cytogenetics
- Community genetics

Abstract deadline

Will be via world wide web.
Closing date 13 January 2003.

Further information available from: The Vienna Medical Academy of Postgraduate Medical Education and Research, Alserstrasse 4, A-1090 Vienna, Austria. Tel +43 1 405 13 83 22 Email: eshg@medacad.org

CORRECTIONS

In the October 2002 issue of the journal, in the paper by Van Maldergem *et al* (*J Med Genet* 2002;39:722-33), we regret that some of the authors' names and affiliations were inadvertently omitted. They were:

N Tubiana-Ruffi

Service d'Endocrinologie Pédiatrique, Hôpital Robert-Debré, Paris, France

A Mégarbané

Unité de Génétique Médicale, Université Saint-Joseph, Beirut, Lebanon

J Maassen

Silvius Laboratory, University of Leiden Medical Centre, Leiden, The Netherlands

M Polak

INSERM U457, Université Paris VIII, Paris, France

D Lacombe

Service de Génétique Médicale, Hôpital Pellegrin-Enfants, Bordeaux, France

C R Kahn

Joslin Diabetes Center, Harvard University, Boston, USA

E L Silveira

Private Practice, Porto Alegre, Brazil

F H D'Abronzio

Department of Endocrinology, Faculdade de Medicina de Jundiaí, Brazil

F Grigorescu

Molecular Endocrinology, IURC, Montpellier, France

S O'Rahilly

Department of Medicine and Clinical Biochemistry, Addenbrooke's Hospital, Cambridge, UK

.....

In the July 2002 issue of the journal, in the Online mutation report by Olivieri *et al* (p e39), there was a missprint in table 2. For No 12 the mutation should have been 1031G>T instead of 1006G>T.

.....

.....

In the December 2002 issue of the journal, in the paper by Khoo *et al* (pp 906–912), all c.1732delTCinsAC mutations should read C.1732delTCinsA. This error occurs on page 906 (Abstract) and page 910 (Discussion).