ELECTRONIC LETTER

A novel pericentric inversion of chromosome 3 cosegregates with a developmental-behavioural phenotype

D Efron, M B Delatycki, M G de Silva, A Langbein, W Slaghuis, A Larson, H-H M Dahl, S M Forrest

J Med Genet 2003;40:e15(<http://www.jmedgenet.com/cgi/content/full/40/2/e15>)

Current advances in genetic knowledge and analysis have facilitated the shift in emphasis from single gene disorders to complex traits such as cancer and atopic disorders where changes in more than one gene usually contribute to the disease phenotype. A challenging area of complex trait research is the determination of the genetic basis of behavioural phenotypes. Early studies in this area focused on classical linkage analysis in pedigrees which segregated the trait of interest and led to the identification of genes for intellectual disability, such as *fraX(A)*.^{1,2} In this disorder, mutations in a single gene are sufficient to generate the disease phenotype.

In contrast, with other neurodevelopmental disorders, results to date have been much more complex. With ADHD, for example, a large number of genes affecting neurotransmitter function, particularly dopamine, have been implicated. However, results from linkage or association studies have not been reliably replicated. Even the most promising gene candidates are implicated in only a small percentage of ADHD cases.^{3,4} Identification of the genes involved is a challenging task. Firstly, accurate phenotype definition is crucial to successful outcomes in these pooled studies. Secondly, neurodevelopmental disorders such as ADHD are multifactorial, with a number of different genes, likely to be of varying effect and interacting with environmental influences, presumably contributing to the development of the observed phenotypes.

We have identified an extended family where the proband, a child with moderate intellectual disability as well as severe conduct disturbance, was found to have a pericentric inversion of chromosome 3. The inversion was found to cosegregate with developmental-behavioural problems in other members of the family. In order to describe the phenotype, a developmental-behavioural paediatrician undertook a standardised clinical evaluation of eight of the juvenile members of the extended family. The findings are presented in this paper.

METHODS

Participants

The parents/guardians of eight juvenile family members were approached by the geneticist and invited to participate in the research. Children were chosen for this evaluation because the measures for the study of behavioural symptoms in children are better established than those for adults. In addition, one other child (no inversion) and six adults in the family (four with the inversion and two without) also underwent IQ testing. (This child was unavailable for detailed assessment, and the adults were not assessed other than IQ testing, as discussed above). All agreed to be involved and gave informed consent. The project was approved by the Royal Children's Hospital Ethics in Human Research Committee (EHRC 98037C) and that of the University of Tasmania (approval No H4077).

Key points

- A child with a major neurodevelopmental impairment and severe impulse control problems was found to have a pericentric inversion of chromosome 3 [46N inv(3)(p14q21)]. As several relatives also had developmental-behavioural symptoms, chromosomal analysis was undertaken on a number of family members. This showed 10 other subjects who carried the inversion.
- In order to define which phenotypic characteristics were attributable to the inversion, as opposed to familial or individual traits, standardised comprehensive assessments of eight subjects (four with the inversion, four without) were undertaken by a developmental-behavioural paediatrician blinded as to the subjects' inversion status. In addition, intelligence testing was performed on 15 family members (eight with the inversion, seven without).
- Subjects with the inversion were found to have more significant developmental disabilities than those with normal karyotypes. The clinical phenotypic features associated with this inversion were intellectual impairment and an impulsive behaviour style. This is the first report of a behavioural phenotype associated with these loci and identifies a novel resource for defining the gene(s) involved.
- We hypothesise that one or more genes are disrupted by this inversion, resulting in the observed phenotype. Further molecular analysis of this family should provide an important contribution to the identification and characterisation of genes involved in this behavioural phenotype, which presents with some of the features commonly seen in attention deficit hyperactivity disorder (ADHD).

Cytogenetic analysis

Chromosome preparations were prepared using standard procedures⁵ and G banded chromosomes were assembled and described according to the recommendations of the International Standing Committee on Human Cytogenetic Nomenclature (ISCN).⁶

Clinical assessment

A standardised assessment procedure was undertaken, involving the completion of behaviour rating scales by the children's parents/guardians and teachers. The developmental-behavioural paediatrician performed a clinical

Figure 1 (A) Partial karyotype of the proband showing the normal chromosome 3 (a) and the chromosome with the pericentric inversion, $\text{inv}(3)(p14q21)$ (b). The extent of the inversion is indicated by arrows. (B) Ideogram of the chromosome 3 inversion showing the normal chromosome 3 (left) and the chromosome 3 with the inversion (right). The breakpoints of the inversion are indicated with arrows.

assessment. The clinical evaluation consisted of a semi-structured interview (pregnancy/perinatal period, early development, temperament/behaviour, learning, general health) and a neurodevelopmental assessment.

Psychological instruments

Wechsler Intelligence Scale for Children-III (WISC-III)⁷ and Wechsler Adult Intelligence Scale-Revised (WAIS-R)⁸ were undertaken by Amy Langbein. The Child Behaviour Checklist (CBCL)⁹ and complimentary Teacher's Report Form (TRF)¹⁰ are broad band standardised behaviour rating scales. These 113 item checklists provide a multidimensional profile of empirically derived problem behaviour syndromes. Conners' Parent Rating Scale - Revised and Conners' Teacher Rating Scale - Revised¹¹ have been the most frequently used behaviour rating scales in ADHD research.¹² The revised 48 item version of the Conners' Parent Rating Scale (CPRS-R) and the 28 item version of the Conners' Teacher Rating Scale (CTRS-R) were used in this study. DSM-IV ADHD Parent and Teacher Rating

Scales¹³ consists of the 18 items in The American Psychiatric Association's Diagnostic and Statistical Manual of Mental Disorders, 4th edition (DSM-IV) diagnostic criteria for ADHD, listed such that the Inattention items alternate with the Hyperactive/Impulsive items.

Blinding

The paediatrician knew that some of the children assessed had the inversion, but did not know which ones. An attempt was made to remain blinded to the inversion status of all subjects. This was successful in all cases except the proband (subject 1) whose carer inadvertently revealed that he had the inversion. As this subject clearly had the most severe developmental disability of all subjects assessed (and hence was very likely to be affected), this revelation did not constitute a major departure from the study protocol.

Figure 2 Pedigree of the family investigated in this study. Closed black symbols indicate the subjects who carry the chromosome inversion. N = normal karyotype. The children who underwent clinical assessment by the developmental/behavioural paediatrician in this study are indicated by the numbers 1 to 8. The proband is indicated with an arrow.

Table 1 Clinical assessment findings

Subject	Sex	Age (y.m)	Positive findings from assessment	IQ testing			Summary formulation
				Verbal	Perf	FS	
1*	M	13.9	Developmental concerns from 18/12; severe behavioural disturbance and learning difficulties; multiple neuro-developmental delays; conduct disturbance; features of anxiety and depression.	56	48	48	Moderate intellectual disability; marked impulsivity, conduct disturbance
2*	F	9.3	Behavioural problems from pre-school age; delayed short term auditory memory, poor fine motor skills	69	77	70	Consistent with treated ADHD (combined type)†
3	M	12.6	Learning difficulties, poor impulse control	67	66	64	Learning difficulties
4*	F	12.2	Learning difficulties; impulsive, uncooperative; delayed short term auditory memory.	74	78	74	Mild intellectual disability; impulsivity, behavioural disturbance
5	F	10.4		104	100	101	Normal child
6*	M	15.6	Behavioural disturbance, learning difficulties.			83	Borderline intellectual function with behavioural disturbance
7	M	12.8	Aggression, withdrawn, low self-esteem; onset 8 y; delayed short term auditory memory	90	104	96	Emotional disturbance
8	M	11.4	Learning difficulties	73	91	80	Learning difficulties

*Inversion present. †On methylphenidate for past two years. Perf = performance IQ. FS = full scale IQ.

Statistical methodology

Analysis of the difference in IQ between those with and those without the inversion was undertaken using a one tailed *t* test (Microsoft Excel for Windows).

RESULTS

The proband (subject 1), a 10 year old male, had been managed jointly by a child psychiatrist and paediatrician. He had moderate intellectual disability, as well as severe impulse control problems with features of ADHD and conduct disorder. His symptoms responded to high doses of stimulants. Cytogenetic analysis was performed as part of the investigative process. The child was found to have a pericentric inversion of chromosome 3 (46N inv(3)(p14q21) (fig 1A, B). A family member suggested that several other relatives had a similar behavioural phenotype and predicted that they too would carry the inversion. The family was then referred to the genetics clinic. Twenty other family members supplied blood

samples for karyotyping. Their inversion status was correctly predicted by the family member in 19 of the 20 cases. Of the 21 who had cytogenetic analysis, 11 had the inversion and 10 did not (fig 2). Most of the subjects with this inversion had clinically significant developmental-behavioural problems, and some had been diagnosed with attention deficit hyperactivity disorder (ADHD) and were being treated with psychostimulant medication (table 1).

Seven of the eight children assessed had clinical evidence of significant developmental difficulties (table 1). The types of clinical problems varied greatly within the sample studied.

Behavioural problems were present in six children. The results of the behavioural rating scales are shown in table 2. In subject 7 (no inversion), onset was at around the age of 8 years, suggesting environmental factors were a more important cause than constitutional factors. Two other subjects without the inversion (3 and 8) had primarily learning difficulties. These children were different from one another

Table 2 Results of the various rating scales used in evaluating the eight subjects. The results presented are those where the factors/scales with scores were >2 SD above the mean for CPRS-R, CTRS-R, CBCL, and TRF. For DSM-IV scales, results are given where six or more items were present

Subject	DSM-IV Parent	DSM-IV Teacher	CPRS-R	CTRS-R	CBCL	TRF
1	Inattent, hyp/imp	Inattent, hyp/imp	Conduct problem, learning problem, impulsive-hyperactive, anxiety	Conduct problem, hyperactivity, inattentive-passive	Anxious-depressed, social problems, thought problems, attention problems, delinquent behaviour, aggressive behaviour	Anxious-depressed, social problems, delinquent behaviour, aggressive behaviour
2	–	–	Conduct problem	–	Delinquent behaviour	–
3	–	–	–	–	–	–
4	–	–	Conduct problem, learning problem, psychosomatic, impulsive-hyperactive,	–	Withdrawn, thought problems, attention problems, delinquent behaviour, aggressive behaviour	–
5	–	–	–	–	–	–
6	Inattent	–	Learning problem, psychosomatic, impulsive-hyperactive	–	Withdrawn, somatic complaints, anxious-depressed, social problems, thought problems, attention problems, delinquent behaviour, aggressive behaviour	–
7	–	–	Conduct problem, impulsive-hyperactive	–	Withdrawn, anxious-depressed, social problems	Somatic complaints, anxious-depressed, thought problems
8	–	–	–	–	–	–

CPRS-R = Conners' Parent Rating Scale - Revised; CTRS-R = Conners' Teacher Rating Scale - Revised; CBCL = Child Behaviour Checklist; TRF = Teacher's Report Form.

temperamentally. Subject 3 had problems with impulsivity and aggression.

Of the eight children assessed, the clinical presentation in three (subjects 1, 4, and 6, all with inversion) strongly suggested a predominantly constitutional pathogenesis (early developmental difficulties, persistent, pervasive, and severe symptomatology with poor impulse control). Subject 2 had been taking stimulant medication for two years; however, historically she also had a constitutional problem, ADHD (table 1).

In addition, one other child (no inversion) and six adults in the family (four with the inversion and two without) also underwent IQ testing. Overall, the mean IQ of eight subjects with the inversion (four adults, four children) was 76.6, compared to 93.7 for the seven subjects (two adults, five children) without the inversion ($p=0.03$).

DISCUSSION

On clinical assessment at one point in time, those children who carried a pericentric inversion of chromosome 3 were found to have more significant developmental disabilities than those without the inversion. The clinical phenotypic features in common were intellectual difficulties and an impulsive behaviour style. The difference in IQ scores between subjects with the inversion and those without was statistically significant. Three of the four control family members had less severe developmental-behavioural problems. These tended to be qualitatively different from those with the inversion (one had learning difficulties, one reactive emotional disturbance), though the phenotype of subject 3 (learning difficulties with poor impulse control) did have similarities to that of subjects with the inversion.

These findings suggest that the inversion is causally related to the phenotype described. A number of disease causing genes have been localised by the cosegregation of the disease with an apparently balanced chromosomal rearrangement. An example of this is the localisation of the neurofibromatosis type 1 gene to chromosome 17q11.2 by mapping a translocation breakpoint cosegregating with the disease.¹⁴ More recently, chromosomal translocations have been identified in some patients with behavioural disorders such as autism. Breakpoints have been mapped to chromosome 7q31 and 22 and candidate genes are being examined.¹⁵

The chromosomal abnormality observed in this family may mediate the development of the observed phenotype by physically disrupting or altering a gene or genes involved in intelligence and/or behaviour. Alternatively, the disruption or alteration of a suppressor or an enhancer element at one or both of the breakpoints may change the level of expression of the target gene(s). Similarly, the physical rearrangement of the chromosome may alter the chromatin structure or reposition the gene of interest such that gene transcription is enhanced or suppressed. More remotely, it is possible that a novel gene is created at the inversion breakpoint region consisting of parts of genes derived from each arm of the chromosome. Similarly, a promoter of differing activity to the promoter of origin could be juxtaposed onto the 5' end of a gene involved in behaviour and/or intelligence altering the level of its expression.¹⁶

Many genes in the region of both breakpoints in the described family are expressed in the brain.¹⁷ This is an unsurprising finding since a high proportion of all genes are expressed in the central nervous system. Therefore the precise position of the inversion breakpoints will need to be defined at a molecular level to identify whether one or more genes have indeed been disrupted. Continuing studies are under way to identify gene(s) that are disrupted by this inversion.

It is of considerable interest to note that the first genome wide scan for loci involved in ADHD found a QTL peak (lod 1.37, $p=0.0060$) over the marker D3S1569 in band 3q21.¹⁸

Although the lod score did not exceed the QTL threshold for significance, it is located near the q arm breakpoint of the described inversion. Also intriguing, a genome screen for genes predisposing to bipolar affective disorder, a condition with phenomenological overlap with ADHD, found "suggestive" linkage to 3p14, the site of the other inversion breakpoint.¹⁹ Taken together the results of these two studies support our hypothesis that the chromosome 3 inversion is causally related to the phenotype described in the family in this paper.

The identification of a gene that is involved in the phenotype of impulsivity has a number of important clinical implications. In particular, the impulsive phenotype is part of the diagnostic profile of two of the three subtypes of patients with ADHD. Analysis of a cohort of such patients for mutations or polymorphisms at this locus will determine the impact this gene has on the aetiology of ADHD. If the impact is significant, better diagnostic testing based on genetic analysis may be possible for some children presenting with ADHD-like symptoms. This family affords the opportunity firmly to implicate a known biological pathway in the development of this behavioural trait or, alternatively, implicate a new gene family.

ACKNOWLEDGEMENTS

The first two authors contributed equally to this study. The authors are very grateful to the family described for allowing this study to go ahead. In particular the grandmother of the proband has been extremely helpful in coordinating interactions between family members and ourselves. The project was funded by the Collaborative Research Centre for the Discovery of Genes for Common Human Diseases. The CRC for Discovery of Genes for Common Human Diseases is established and supported by the Australian Government's Cooperative Research Centre's Programme. Professor Bob Williamson is thanked for his support and intellectual input into this study.

Authors' affiliations

D Efron, Centre for Community Child Health, Royal Children's Hospital, Parkville, Victoria, Australia

D Efron, M B Delatycki, M G de Silva, H-H M Dahl, S M Forrest, Murdoch Children's Research Institute, Royal Children's Hospital, Parkville, Victoria, Australia

D Efron, M B Delatycki, H-H M Dahl, S M Forrest, Department of Paediatrics, University of Melbourne, Parkville, Victoria, Australia

M B Delatycki, Genetic Health Services Victoria, Royal Children's Hospital, Parkville, Victoria, Australia

M B Delatycki, M G de Silva, H-H M Dahl, S M Forrest, Cooperative Research Centre for Discovery of Genes for Common Human Diseases, Parkville, Victoria, Australia

M B Delatycki, Department of Paediatrics, Monash University, Clayton, Victoria, Australia

A Langbein, W Slaghuis, Department of Psychology, University of Tasmania, Sandy Bay, Tasmania, Australia

A Larson, Calvary Rehabilitation Service, New Town, Tasmania, Australia

Correspondence to: Dr M B Delatycki, Genetic Health Services Victoria, Royal Children's Hospital, Flemington Road, Parkville, Victoria 3052, Australia; delatycm@cryptic.rch.unimelb.edu.au

REFERENCES

- Verkerk AJ**, Pieretti M, Sutcliffe JS, Fu YH, Kuhl DP, Pizzuti A, Reiner O, Richards S, Victoria MF, Zhang F, Eussen BE, van Ommen G-JB, Blonden LAJ, Riggings GJ, Chastain JL, Kunst CB, Galjaard H, Caskey CT, Nelson DL, Oostra BA, Warren ST. Identification of a gene (FMR-1) containing a CGG repeat coincident with a breakpoint cluster region exhibiting length variation in fragile X syndrome. *Cell* 1991;**65**:905-14.
- Yu S**, Pritchard M, Kremer E, Lynch M, Nancarrow J, Baker E, Holman K, Mulley JC, Warren ST, Schlessinger D, Sutherland GR, Richards RI. Fragile X genotype characterized by an unstable region of DNA. *Science* 1991;**252**:1179-81.
- Madras BK**, Miller GM, Fischman AJ. The dopamine transporter: relevance to attention deficit hyperactivity disorder (ADHD). *Behav Brain Res* 2002;**130**:57-63.
- Comings DE**. Clinical and molecular genetics of ADHD and Tourette syndrome. Two related polygenic disorders. *Ann NY Acad Sci* 2001;**931**:50-83.

- 5 **Webber LM**, Garson OM. Fluorodeoxyuridine synchronization of bone marrow cultures. *Cancer Genet Cytogenet* 1983;**26**:123-32.
- 6 **ISCN 1995**. In: Mitelman F, ed. *An international system for human cytogenetic nomenclature*. Basel: S Karger, 1995.
- 7 **Wechsler D**. *Wechsler intelligence scale for children*. 3rd ed. San Antonio: The Psychological Corporation, 1991.
- 8 **Wechsler D**. *Wechsler adult intelligence scale*. Revised manual. New York: Psychological Corporation: 1981.
- 9 **Achenbach TM**. *Manual for the child behavior checklist/4-18 and 1991 profile*. Burlington, VT: University of Vermont, 1991.
- 10 **Achenbach TM**. *Manual for the teacher's report form and 1991 profile*. Burlington, VT: University of Vermont, 1991.
- 11 **Goyette CH**, Conners CK, Ulrich RF. Normative data on revised Conners parent and teacher rating scales. *J Abnorm Child Psychol* 1978;**6**:221-36.
- 12 **Barkley RA**. A review of stimulant drug research with hyperactive children. *J Child Psychol Psychiatry* 1977;**18**:137-65.
- 13 **DuPaul GJ**, Barkley RA. *ADHD rating scale*, 1995 (unpublished).
- 14 **Ledbetter DH**, Rich DC, O'Connell P, Leppert M, Carey JC. Precise localization of NF1 to 17q11.2 by balanced translocation. *Am J Hum Genet* 1989;**44**:20-4.
- 15 **Bonora E**, Bacchelli E, Levy ER, Blasi F, Marlow A, Monaco AP, Maestrini E. Mutation screening and imprinting analysis of four candidate genes for autism in the 7q32 region. *Mol Psychiatry* 2002;**7**:289-301.
- 16 **Mickley LA**, Spengler BA, Knutsen TA, Biedler JL Fojo T. Gene rearrangement: a novel mechanism for *MDR-1* gene activation. *J Clin Invest* 1997;**99**:1947-57.
- 17 **Online Mendelian Inheritance in Man**. <http://www3.ncbi.nlm.nih.gov/Omim/>
- 18 **Fisher SE**, Francks C, McCracken JT, McGough JJ, Marlow AJ, MacPhie IL, Newbury DF, Crawford LR, Palmer CG, Woodward JA, Del'Homme M, Cantwell DP, Nelson SF, Monaco AP, Smalley SL. A genomewide scan for loci involved in attention-deficit/hyperactivity disorder. *Am J Hum Genet*. 2002;**70**:1183-196.
- 19 **Cichon S**, Schumacher J, Muller DJ, Hurter M, Windemuth C, Strauch K, Hemmer S, Schulze TG, Schmidt-Wolf G, Albus M, Borrmann-Hassenbach M, Franzek E, Lanczik M, Fritze J, Kreiner R, Reuner U, Weigelt B, Minges J, Lichtermann D, Lerer B, Kanyas K, Baur MP, Wienker TF, Maier W, Rietschel M, Propping P, Nothen MM. A genome screen for genes predisposing to bipolar affective disorder detects a new susceptibility locus on 8q. *Hum Mol Genet* 2001;**10**:2933-44.