ELECTRONIC LETTER

Germline *TP53* mutations in breast cancer families with multiple primary cancers: is *TP53* a modifier of *BRCA1*?

A-M Martin, P A Kanetsky, B Amirimani, T A Colligon, G Athanasiadis, H A Shih, M R Gerrero, K Calzone, T R Rebbeck, B L Weber

J Med Genet 2003;40:e34(<http://www.jmedgenet.com/cgi/content/full/40/4/e34>)

Somatic mutations in *TP53* are the most frequent events in human cancer and lead to inactivation of the gene, loss of tumour suppressor function, and in some cases generation of a dominant negative form of p53.¹⁻³ Eleven exons make up the primary transcript of *TP53*, of which exons 2-11 encode the protein. Five conserved domains exist in exons 1, 4, 5, 7, and 8,⁴ which are considered essential for normal p53 function. Approximately 90% of disease associated mutations occur in these domains, with mutations in five codons (175, 245, 248, 249, and 273) accounting for approximately 20% of all mutations reported to date.

Germline mutations in *TP53* cause Li-Fraumeni syndrome (LFS), a familial association of childhood leukaemia, brain cancer, soft tissue sarcoma, and adrenal cortical carcinoma,^{5,6} as well as other cancers such as breast cancer, melanoma, germ cell tumours, and carcinomas of the lung, pancreas, and prostate.^{7,8} Cancers characteristically develop at unusually early ages and multiple primary tumours are frequent. Susceptibility to cancer in these families follows an autosomal dominant pattern of inheritance⁷ and among families with a known germline *TP53* mutation the probability of developing any invasive cancer (excluding carcinomas of the skin) approaches 50% by the age of 30, compared to an age adjusted population incidence of cancer of 1%. It is estimated that more than 90% of *TP53* mutation carriers will develop cancer by the age of 70.⁹

In addition to the numerous mutations, *TP53* also contains several polymorphisms that may alter its activity. In particular, at nucleotide 215 (codon 72) there is a single base pair variant (g.215G>C) in the coding region, which results in a substitution of proline for arginine in the protein sequence.¹⁰ The frequency of this polymorphism varies from 26-35%¹¹⁻¹³ and it appears to affect protein function. The R72 variant of *TP53* is believed to be more sensitive to human papillomavirus (HPV) induced degradation by the E6 oncoprotein than the 72P variant, and is thought to be of functional significance in HPV associated tumours¹⁴ such as cervical tumours.¹⁵⁻¹⁷ Furthermore, some, but not all studies document an overrepresentation of R72 variant in cervical cancer patients compared to a control population.^{18,19} However, other reports suggest the association of the 72P variant with incidence of squamous cell carcinoma of the head and neck²⁰ and lung adenocarcinomas in smokers.²¹

In families with multiple cases of breast cancer that do not fit the criteria for LFS, the frequency of *TP53* germline mutations has been investigated in multiple studies,²²⁻²⁸ documenting that *TP53* mutations account for <1% of site specific breast cancer families.²³ However, among LFS families, there is a very high incidence of early onset breast cancer. Taken together, these data suggest that germline *TP53* mutations are strongly associated with hereditary breast cancer susceptibility but almost exclusively in the context of LFS.

Because of the high penetrance of early onset breast cancer and the known increased incidence of multiple primary cancers in LFS families (50% by 30 years of age),⁹ we investi-

Key points

- Eighty eight women with breast cancer and a personal or family history of multiple primary cancers (MPC) (including ovarian cancer) and 84 women with a personal and family history of breast cancer only (BC) were studied. All women had been previously screened for germline *BRCA1* and *BRCA2* mutations; 38 (43%) of MPC women and 10 (12%) of BC women had a mutation in one of these two genes.
- We determined the frequency of deleterious germline *TP53* mutations, as well as the common R72P polymorphism in *TP53* and investigated the association of this polymorphism with the development of cancers in the entire study set. We also evaluated the association between R72P and breast cancer penetrance in the subset of women with known *BRCA1* or *BRCA2* mutations.
- One woman, from a family with breast cancer only, was found to have a deleterious *TP53* mutation (exon 7, G245S); no deleterious *TP53* mutations were detected in the families with cases of multiple primary cancers. The common R72P polymorphism was seen at a frequency of 41% in the entire sample. MPC women were more likely to be homozygous for R72 compared to BC women ($p=0.05$, OR 2.83, 95% CI 1.2 to 6.9), an association that was more striking in women with a *BRCA1* or *BRCA2* mutation (OR 6.1, 95% CI 1.4 to 26.4).
- We also found that the presence of a 72P allele was associated with an earlier age of breast cancer diagnosis among *BRCA1* mutation carriers ($p=0.05$), suggesting that the R72P polymorphism may be a modifier of *BRCA1* penetrance.

gated whether deleterious germline mutations in *TP53* and/or the R72P polymorphism were associated with multiple primary cancers (in which one was breast cancer) in families with ≥ 2 breast cancers but no evidence of LFS. One previous study investigated the frequency of germline *TP53* mutations with bilateral breast cancer²⁹ and found no *TP53* mutations; however, only 19 samples were tested. In the current study we determined the frequency of deleterious *TP53* germline mutations in 172 breast cancer families, with and without multiple primary cancers. Germline *BRCA1* or *BRCA2* mutation status was known in all subjects³⁰; 43% of women with multiple primary cancers and 12% of women with breast cancer only had

Abbreviations: MPC, multiple primary cancers; BC, breast cancer only; LFS, Li-Fraumeni syndrome; HPV, human papillomavirus

Table 1 Non-breast primary cancers in families with multiple primary cancers

Cancer	No of families with cancer*
Ovarian	29
Colorectal	13
Non-melanoma skin	11
Thyroid	6
Endometrial	6
Cervix	5
Leukaemia	4
Lymphoma	3
Others†	13

*Five patients had two or more non-breast cancers, so the number of cancers does not equal the number of patients.

†Other primary cancers include melanoma, brain, head/neck, sarcoma, lung, kidney, and pharynx.

either a *BRCA1* or *BRCA2* mutation. In addition, we established the frequency of the common exon 4 polymorphism (R72P) in this sample and evaluated whether this polymorphism may be a modifier of breast cancer penetrance in the presence of *BRCA1* or *BRCA2* mutations.

MATERIALS AND METHODS

Patient population

All families were recruited from clinics at the University of Michigan (1993-1995) and the University of Pennsylvania (1995-1998). Patients were either self- or physician referred because of a perceived risk of inherited susceptibility to breast cancer. All women consented to genetic testing for clinical and/or research purposes. Personal and family histories of all cancers were recorded, including age of diagnosis of all cancers and the number of related women in each family at risk for breast cancer (age ≥ 20 years). Pathology reports were obtained on all probands and on other family members when possible. The testing protocol was approved by duly constituted institutional review boards at both the University of Michigan and the University of Pennsylvania.

Eighty-eight women were from families with at least two cases of breast cancer and at least one woman affected with both a primary breast cancer and a primary non-breast cancer (denoted MPC). Eighty-four (95%) MPC women had two primary cancers, and four MPC women (5%) had three or more primary cancers. All non-breast malignancies were considered, including non-melanoma skin cancers. An additional 84 women were from families with at least two cases of breast

Table 2 Description of relatives providing sample for testing

Sample number tested (relative)	Age of breast cancer in relative	Average age of diagnosis of additional cancer(s) in multiply affected subject	Second primary cancer in multiply affected subject	Relationship of subject providing DNA for testing
229	30	58	Ovarian	Daughter
522	56	63	Ovarian	Niece
743	69	74	BCC, cervical NHL	Niece
673	40	43	BCC	Daughter
513	38	30	Cervical	Daughter
641	44	30	Colon	Cousin
813	36	44	Ovarian	Niece
852	56	53	BCC, endometrial	Niece
834	45	40	Cervical	Sister
1019	50	63	Ovarian	Daughter
842	27	47	Ovarian	Sister
975	57	50	BCC	Sister
1965	57	25	Thyroid	Daughter
910	74	55	Thyroid	Daughter
1946	44	84	Leukaemia	Daughter
1901	30	65	Lung	Niece
1907	46	86	Colon, rectal	Daughter
1208	41	51	Ovarian	Sister
1708	72	69	Skin, cervical, endometrial	Sister
1785	51	49	Colon	Granddaughter
1844	52	67	Colon	Daughter
1773	50	51	Cervical	Niece
1762	45	74	Colon	Daughter
1797	36	50	Ovarian	Daughter
1748	42	72	Hodgkin's lymphoma	Granddaughter
1722	50	83	Endometrial	Granddaughter
1719	40	23	Ovarian	Sister
1718	35	36	Ovarian	Granddaughter
1763	41	56	Colon	Daughter
1794	40	40	Ovarian	Cousin
1783	53	53	Throat	Double cousin
1853	57	57	Brain	Grandniece
320	40	40	Pituitary	Niece
1987	55	55	Ovarian	Daughter
1909	70	70	Leukaemia	Granddaughter
1954	61	61	Thyroid, colon	Sister
1851	43	74	Leukaemia	Cousin
1806	49	49	Melanoma	Daughter
1863	39	39	Ovarian	Daughter
1993	67	67	Colon	Cousin
2074	95	95	Colon	Niece
2038	50	52	Ovarian	Daughter
2067	50	50	BCC	Daughter
2233	41	42	ALL	Niece
2241	61	52	Ovarian	Cousin

Table 3 Primer sequences

Exon	Primer sequences	Annealing temp (°C)	Reference
2/3	Forward: 5'-ggatcccttcttctt-3' Reverse: 5'-agcatcaaatcatcattgc-3'	57	
4	Forward: 5'-gacctggctctgactgt-3' Reverse: 5'-atacggccaggcattgaagt-3'	54	
5/6	Forward: 5'-tgccctgacttcaacttgt-3' Reverse: 5'-taacccctctccagaga-3'	54	
7	Forward: 5'-tgccacagggtctcccaagg-3' Reverse: 5'-aggggtcagcggaagcaga-3'	55	Evans <i>et al</i> ⁵³
8/9	Forward: 5'-caagggtgggtggagtaga-3' Reverse: 5'-actgataagagggtcccaag-3'	54	
10	Forward: 5'-atgttgctttgatcgtca-3' Reverse: 5'-ctttcaacctaggaaggca-3'	54	
11	Forward: 5'-agccacctgaactcaaaa-3' Reverse: 5'-aatggcagggaggagaga-3'	55	Evans <i>et al</i> ⁵³

cancer but no cases of multiple primary cancers (denoted BC). DNA was available from at least one woman with multiple primary cancers in 43 families. In the remaining 45 multiple primary cancer families, the multiply affected woman was dead (n=33) or unavailable (n=12). In these families, *TP53* screening was undertaken using DNA from the closest female relative diagnosed with breast cancer. Table 1 provides a description of the cancers reported in subjects with multiple primary cancers and table 2 is a detailed description of the female relatives of a multiply affected woman, who provided samples for testing. All samples were previously screened for germline mutations in *BRCA1* and *BRCA2*³⁰; 38 MPC women and 10 BC women had a *BRCA1* or *BRCA2* germline mutation.³⁰

Mutation analysis

DNA was extracted from peripheral blood mononuclear cells and stored in TE at 4°C. The entire 10 exon coding domain and flanking splice site regions of *TP53* were amplified using seven PCR primer sets (table 3). PCR amplification was performed in a final volume of 20 µl containing 80 ng of DNA, 1.5 mmol/l MgCl₂, 10 mmol/l Tris-HCl (pH 8.3), 50 mmol/l KCl, 0.2 mmol/l each of dCTP, dATP, dTTP, dGTP (Amersham Pharmacia Biotech), each primer at 1.0 µmol/l, and 1.0 unit of *Taq* polymerase (Boehringer Mannheim). Annealing temperatures were optimised for each primer set and ranged from 55–60°C. Variants were identified by conformation sensitive gel electrophoresis (CSGE) as previously described³⁰ and characterised by direct sequencing using the ABI Prism 377 after reamplification from source DNA. All mutation nomenclature is reported using the recommendations of den Dunnen and Antonarakis.³¹

Statistical analysis

Differences in *TP53* mutation frequency between the MPC and BC groups were assessed using χ^2 analysis. Odds ratios (OR)

and 95% confidence intervals (95% CI) were reported. In those instances where expected cell counts fell below five, we used exact methods to determine the 95% CI.³² Furthermore, we used the Mann-Whitney U-Wilcoxon rank sum test to determine whether *TP53* genotypes altered the median age of first breast cancer diagnosis within categories defined by *BRCA1* or *BRCA2* mutation status.

RESULTS

DNA from one woman from a BC family and no *BRCA1* or *BRCA2* mutation showed an abnormal CSGE profile in *TP53* exon 7. Sequence analysis of this variant showed a G to A transition at the first nucleotide in codon 245 resulting in a glycine-serine change at this position (G245S). No presumed deleterious *TP53* mutations were seen in the MPC group (MPC=0%, BC=1.2%, p=0.31).

The proline allele of the R72P polymorphism was seen at a frequency of 41% in the entire sample. The distribution of R72P genotypes within groups is presented in table 4. Owing to the small number of homozygous R72P genotypes, all R72P alleles were combined into one group (P/*). When we performed subgroup analyses in women with *BRCA1* or *BRCA2* mutations, MPC women were six times more likely to have the homozygous R72P genotype than BC women (OR=6.1, 95% CI 1.4 to 26.4) (table 5). However, because of the small sample size, a statistically significant association between the homozygous R72P genotype and MPC could not be confirmed separately in an analysis of only *BRCA1* mutation carriers or only *BRCA2* mutation carriers.

Table 5 *TP53* R72P genotypes in MPC and BC families by *BRCA1/2* mutation status

	MPC (n=43)		BC (n=84)		OR 95% CI
<i>TP53</i> genotype	No	(%)	No	(%)	
<i>BRCA1</i> or <i>BRCA2</i> mutation					
R/R	19	(83)	7	(44)	6.1 (1.4, 26.4)
P/*	4	(17)	9	(56)	1.0
<i>BRCA1</i> mutation					
R/R	16	(80)	3	(43)	5.3 (0.83, 34.1)
P/*	4	(20)	4	(57)	1.0
<i>BRCA2</i> mutation					
R/R	5	(83)	4	(44)	6.3 (0.50, 77.5)
P/*	1	(17)	5	(56)	1.0
No detectable mutation					
R/R	16	(80)	44	(65)	2.2 (0.66, 7.3)
P/*	4	(20)	24	(35)	1.0

P/* = R72P or P72P.

Three women in the sample set had both a *BRCA1* and a *BRCA2* mutation.

Table 4 *TP53* exon 4 R72P genotypes in subjects with multiple primary cancers (MPC) compared to subjects with breast cancer only (BC)

<i>TP53</i> genotype	MPC (n=43)		BC (n=84)		
	No	(%)	No	(%)	
R/R	35	(81)	51	(61)	
R/P	6	(14)	28	(33)	
P/P	2	(5)	5	(6)	
°P/*	8	(19)	33	(39)	OR=2.83 (1.2, 6.9)

°P/* is the combined genotypes of R/P and P/P used for statistical analysis owing to the rarity of the P allele.

primary cancers in breast cancer families and support previous studies suggesting that *TP53* mutations account for less than 1% of hereditary susceptibility to breast cancer. However, we found presence of the homozygous R72 allele was associated with a six-fold increased risk for the development of multiple primary cancers among subjects with a germline *BRCA1* or *BRCA2* mutation. Finally, we provide preliminary evidence that the arginine allele of R72P in exon 4 of *TP53* may modify *BRCA1* associated breast cancer risk, using age of diagnosis as a surrogate for penetrance.

ACKNOWLEDGEMENTS

This work was supported by grants from the Susan G Komen Breast Cancer Foundation (PDF99-003024 to A-MM) and a grant from the NCI (CA 76417) to BLW.

Authors' affiliations

A-M Martin, Laboratory of Molecular Pathology, Department of Pathology, Pennsylvania Hospital, Philadelphia, USA
A-M Martin, B Amirani, T A Colligon, G Athanasiadis, H A Shih, M R Gerrero, B L Weber, Department of Medicine, University of Pennsylvania, Philadelphia, USA
A-M Martin, T A Colligon, G Athanasiadis, M R Gerrero, K Calzone, B L Weber, Abramson Family Cancer Research Institute, University of Pennsylvania Cancer Center, Philadelphia, USA
P A Kanetsky, T R Rebbeck, Department of Biostatistics and Epidemiology, University of Pennsylvania, Philadelphia, USA
P A Kanetsky, T R Rebbeck Center for Clinical Epidemiology and Biostatistics, University of Pennsylvania, Philadelphia, USA

Correspondence to: Dr B L Weber, Room 514, BRB II/III, 421 Curie Boulevard, Philadelphia, PA 19104, USA; weberb@mail.med.upenn.edu

REFERENCES

- Kern SE, Pietenpol JA, Thiagalingam S, Seymour A, Kinzler KW, Vogelstein B. Oncogenic forms of p53 inhibit p53-regulated gene expression. *Science* 1992;**256**:827-30.
- Vogelstein B, Kinzler KW. p53 function and dysfunction. *Cell* 1992;**70**:523-6.
- Marutani M, Tonoki H, Tada M, Takahashi M, Kashiwazaki H, Hida Y, Hamada J, Asaka M, Moriuchi T. Dominant-negative mutations of the tumor suppressor p53 relating to early onset of glioblastoma multiforme. *Cancer Res* 1999;**59**:4765-9.
- Ko LJ, Prives C. p53: puzzle and paradigm. *Genes Dev* 1996;**10**:1054-72.
- Li FP, Fraumeni JF. Soft-tissue sarcomas, breast cancer, and other neoplasms: familial syndrome? *Ann Intern Med* 1969;**71**:747.
- Li FP, Fraumeni JF. Prospective study of a family cancer syndrome. *JAMA* 1982;**247**:2692-4.
- Strong LC, Stine M, Norsted TL. Cancer in survivors of childhood soft tissue sarcoma and their relatives. *J Natl Cancer Inst* 1987;**79**:1213-20.
- Li FP, Fraumeni JF, Mulvihill JJ. A cancer family syndrome in 24 kindreds. *Cancer Res* 1988;**48**:5358.
- Malkin D, Li FP, Strong LC, Fraumeni JF Jr, Nelson CE, Kim DH, Kassel J, Gryka MA, Bischoff FZ, Tainsky MA, et al. Germ line p53 mutations in a familial syndrome of breast cancer, sarcomas, and other neoplasms. *Science* 1990;**250**:1233-8.
- Harris N, Brill E, Shohat O, Prokocimer M, Wolf D, Arai N, Rotter V. Molecular basis for heterogeneity of the human p53 protein. *Mol Cell Biol* 1986;**6**:4650-6.
- Kawajiri K, Nakachi K, Imai K, Watanabe J, Hayashi S. Germ line polymorphisms of p53 and CYP1A1 genes involved in human lung cancer. *Carcinogenesis* 1993;**14**:1085-9.
- Sjalander A, Birgander R, Hallmans G, Cajander S, Lenner P, Athlin L, Beckman G, Beckman L. p53 polymorphisms and haplotypes in breast cancer. *Carcinogenesis* 1996;**17**:1313-16.
- Wang-Gohrke S, Rebbeck TR, Besenfelder W, Kreienberg R, Runnebaum IB. p53 germline polymorphisms are associated with an increased risk for breast cancer in German women. *Anticancer Res* 1998;**18**:2095-9.
- Soussi T, Beroud C. Assessing TP53 status in human tumours to evaluate clinical outcome. *Nat Rev Cancer* 2001;**1**:233-40.
- Kammer C, Tommasino M, Syrjanen S, Delius H, Hebling U, Warthorst U, Pfister H, Zehbe I. Variants of the long control region and the E6 oncogene in European human papillomavirus type 16 isolates: implications for cervical disease. *Br J Cancer* 2002;**86**:269-73.
- Zehbe I, Tachezy R, Mytilineos J, Voglino G, Mikyskova I, Delius H, Marongiu A, Gissmann L, Wilander E, Tommasino M. Human papillomavirus 16 E6 polymorphisms in cervical lesions from different European populations and their correlation with human leukocyte antigen class II haplotypes. *Int J Cancer* 2001;**94**:711-16.
- Zehbe I, Wilander E, Delius H, Tommasino M. Human papillomavirus 16 E6 variants are more prevalent in invasive cervical carcinoma than the prototype. *Cancer Res* 1998;**58**:829-33.
- Zehbe I, Voglino G, Wilander E, Genta F, Tommasino M. Codon 72 polymorphism of p53 and its association with cervical cancer. *Lancet* 1999;**354**:218-19.
- Zehbe I, Voglino G, Wilander E, Delius H, Marongiu A, Edler L, Klimek F, Andersson S, Tommasino M. p53 codon 72 polymorphism and various human papillomavirus 16 E6 genotypes are risk factors for cervical cancer development. *Cancer Res* 2001;**61**:608-11.
- Shen H, Zheng Y, Sturgis EM, Spitz MR, Wei Q. P53 codon 72 polymorphism and risk of squamous cell carcinoma of the head and neck: a case-control study. *Cancer Lett* 2002;**183**:123-30.
- Fan R, Wu MT, Miller D, Wain JC, Kelsey KT, Wiencke JK, Christiani DC. The p53 codon 72 polymorphism and lung cancer risk. *Cancer Epidemiol Biomarkers Prev* 2000;**9**:1037-42.
- Prosser J, Elder PA, Condie A, MacFadyen I, Steel CM, Evans HJ. Mutations in p53 do not account for heritable breast cancer: a study in five affected families. *Br J Cancer* 1991;**63**:181-4.
- Borresen AL, Andersen TI, Garber J, Barbier-Pirau N, Thorlacius S, Eyfjord J, Ottestad L, Smith-Sorensen B, Hovig E, Malkin D, et al. Screening for germ line TP53 mutations in breast cancer patients. *Cancer Res* 1992;**52**:3234-6.
- Jolly KW, Malkin D, Douglass EC, Brown TF, Sinclair AE, Look AT. Splice-site mutation of the p53 gene in a family with hereditary breast-ovarian cancer. *Oncogene* 1994;**9**:97-102.
- Sun XF, Johannsson O, Hakansson S, Sellberg G, Nordenskjold B, Olsson H, Borg A. A novel p53 germline alteration identified in a late onset breast cancer kindred. *Oncogene* 1996;**13**:407-11.
- Warren W, Eeles RA, Ponder BA, Easton DF, Averill D, Ponder MA, Anderson K, Evans AM, DeMars R, Love R, et al. No evidence for germline mutations in exons 5-9 of the p53 gene in 25 breast cancer families. *Oncogene* 1992;**7**:1043-6.
- Lehman TA, Haffty BG, Carbone CJ, Bishop LR, Gumbs AA, Krishnan S, Shields PG, Modali R, Turner BC. Elevated frequency and functional activity of a specific germ-line p53 intron mutation in familial breast cancer. *Cancer Res* 2000;**60**:1062-9.
- Rapakko K, Allinen M, Syrjakoski K, Vahteristo P, Huusko P, Vahakangas K, Eerola H, Kainu T, Kallioniemi OP, Nevanlinna H, Winqvist R. Germline TP53 alterations in Finnish breast cancer families are rare and occur at conserved mutation-prone sites. *Br J Cancer* 2001;**84**:116-19.
- Lidereau R, Soussi T. Absence of p53 germ-line mutations in bilateral breast cancer patients. *Hum Genet* 1992;**89**:250-2.
- Shih HA, Nathanson KL, Seal S, Collins N, Stratton MR, Rebbeck TR, Weber BL. BRCA1 and BRCA2 mutations in breast cancer families with multiple primary cancers. *Clin Cancer Res* 2000;**6**:4259-64.
- den Dunnen JT, Antonarakis SE. Mutation nomenclature extensions and suggestions to describe complex mutations: a discussion. *Hum Mutat* 2000;**15**:7-12.
- Mehta CR, Patel NR, Gray R. Computing an exact confidence interval for the common odds ratio in several 2x2 contingency tables. *J Am Stat Assoc* 1985;**80**:969-73.
- Toguchida J, Yamaguchi T, Dayton SH, Beauchamp RL, Herrera GE, Ishizaki K, Yamamoto T, Meyers PA, Little JB, Sasaki MS, et al. Prevalence and spectrum of germline mutations of the p53 gene among patients with sarcoma. *N Engl J Med* 1992;**326**:1301-8.
- Brose MS, Rebbeck TR, Calzone KA, Stophel JE, Nathanson KL, Weber BL. Cancer risk estimates for BRCA1 mutation carriers identified in a risk evaluation program. *J Natl Cancer Inst* 2002;**94**:1365-72.
- Storey A, Thomas M, Kalita A, Harwood C, Gardiol D, Mantovani F, Breuer J, Leigh IM, Matlashewski G, Banks L. Role of a p53 polymorphism in the development of human papillomavirus-associated cancer. *Nature* 1998;**393**:229-34.
- Makni H, Franco EL, Kaiano J, Villa LL, Labrecque S, Dudley R, Storey A, Matlashewski G. P53 polymorphism in codon 72 and risk of human papillomavirus-induced cervical cancer: effect of inter-laboratory variation. *Int J Cancer* 2000;**87**:528-33.
- Agorastos T, Lambropoulos AF, Constantinidis TC, Kotsis A, Bontis JN. p53 codon 72 polymorphism and risk of intra-epithelial and invasive cervical neoplasia in Greek women. *Eur J Cancer Prev* 2000;**9**:113-18.
- Dokianakis DN, Spandidos DA. P53 codon 72 polymorphism as a risk factor in the development of HPV-associated cervical cancer. *Mol Cell Biol Res Commun* 2000;**3**:111-14.
- Tenti P, Vesentini N, Rondo Spauldo M, Zappatore R, Migliora P, Carnevali L, Ranzani GN. p53 codon 72 polymorphism does not affect the risk of cervical cancer in patients from northern Italy. *Cancer Epidemiol Biomarkers Prev* 2000;**9**:435-8.
- Tachezy R, Mikyskova I, Salakova M, Van Ranst M. Correlation between human papillomavirus-associated cervical cancer and p53 codon 72 arginine/proline polymorphism. *Hum Genet* 1999;**105**:564-6.
- Yamashita T, Yaginuma Y, Saitoh Y, Kawai K, Kurakane T, Hayashi H, Ishikawa M. Codon 72 polymorphism of p53 as a risk factor for patients with human papillomavirus-associated squamous intraepithelial lesions and invasive cancer of the uterine cervix. *Carcinogenesis* 1999;**20**:1733-6.
- Klaes R, Ridder R, Schaefer U, Benner A, von Knebel Doeberitz M. No evidence of p53 allele-specific predisposition in human papillomavirus-associated cervical cancer. *J Mol Med* 1999;**77**:299-302.
- Rosenthal AN, Ryan A, Al-Jehani RM, Storey A, Harwood CA, Jacobs U. p53 codon 72 polymorphism and risk of cervical cancer in UK. *Lancet* 1998;**352**:871-2.
- Ouchi T, Monteiro AN, August A, Aaronson SA, Hanafusa H. BRCA1 regulates p53-dependent gene expression. *Proc Natl Acad Sci USA* 1998;**95**:2302-6.

- 45 **Zhang H**, Somasundaram K, Peng Y, Tian H, Bi D, Weber BL, El-Deiry WS. BRCA1 physically associates with p53 and stimulates its transcriptional activity. *Oncogene* 1998;**16**:1713-21.
- 46 **Marmorstein LY**, Kinev AV, Chan GK, Bochar DA, Beniya H, Epstein JA, Yen TJ, Shiekhattar R. A human BRCA2 complex containing a structural DNA binding component influences cell cycle progression. *Cell* 2001;**104**:247-57.
- 47 **Chai YL**, Cui J, Shao N, Shyam E, Reddy P, Rao VN. The second BRCT domain of BRCA1 proteins interacts with p53 and stimulates transcription from the p21WAF1/CIP1 promoter. *Oncogene* 1999;**18**:263-8.
- 48 **Harkin DP**, Bean JM, Miklos D, Song YH, Truong VB, Englert C, Christians FC, Ellisen LW, Maheswaran S, Oliner JD, Haber DA. Induction of GADD45 and JNK/SAPK-dependent apoptosis following inducible expression of BRCA1. *Cell* 1999;**97**:575-86.
- 49 **Crook T**, Crossland S, Crompton MR, Osin P, Gusterson BA. p53 mutations in BRCA1-associated familial breast cancer. *Lancet* 1997;**350**:638-9.
- 50 **Schuyler M**, Berns EM. Is TP53 dysfunction required for BRCA1-associated carcinogenesis? *Mol Cell Endocrinol* 1999;**155**:143-52.
- 51 **Bertwistle D**, Ashworth A. Functions of the BRCA1 and BRCA2 genes. *Curr Opin Genet Dev* 1998;**8**:14-20.
- 52 **Xu X**, Qiao W, Linke S, Cao L, Li W, Furth P, Harris C, Deng C. Genetic interactions between tumor suppressors Brca1 and p53 in apoptosis, cell cycle and tumorigenesis. *Nat Genet* 2001;**28**:266-71.
- 53 **Evans SC**, Mims B, McMasters KM, Foster CJ, de Andrade M, Amos CI, Strong LC, Lozano G. Exclusion of a p53 germline mutation in a classic Li-Fraumeni syndrome family. *Hum Genet* 1998;**102**:681-6.