ELECTRONIC LETTER

Prevalence of *SMN1* deletion and duplication in carrier and normal populations: implication for genetic counselling

V Cusin, O Clermont, B Gérard, D Chantereau, J Elion

J Med Genet 2003;40:e39(<http://www.jmedgenet.com/cgi/content/full/40/4/e39>)

Spinal muscular atrophy (SMA) is the second most frequent autosomal recessive disease, with a prevalence of 1 in 6000 live born infants.¹ It is characterised by degeneration of motor neurones of the anterior horn of the spinal cord, leading to symmetrical muscular weakness and atrophy. The International SMA Consortium classification² defines several degrees of severity in the SMA phenotype, depending on the age of onset and motor development milestones. Type I SMA, Werdnig-Hoffmann I disease, is the most severe form with onset within 6 months of birth. Patients are unable to sit up and have serious respiratory dysfunction. Type II SMA is the intermediate form with onset within the first 2 years; children can sit up but are unable to walk. The clinical course is variable. Type III (also called Kugelberg-Welander disease) begins after 2 years of age and usually has a chronic evolution. Children can stand and walk unaided at least in infancy. Adult form (type IV) is the mildest, with onset after 30 years of age; few cases have been reported and its prevalence is not accurately known.

Spinal muscular atrophy is linked to locus 5q13 in more than 95% of patients.³⁻⁶ The critical region, containing several genes including the survival motor neurone (*SMN*) gene, is inverted and duplicated. Homozygous deletion of *SMN1*, located in the telomeric position, accounts for the disease in 98% of these cases and has been reported in infantile, intermediate, and adult onset disease.⁷⁻¹⁰ Linkage analysis in families with SMA shows large de novo deletions in 2% of patients.¹¹⁻¹³ *SMN2* is a highly homologous gene located in the centromeric duplicated region.^{7,14} Most of the *SMN1* transcripts are full length, whereas most of the *SMN2* transcripts lack exon 7. In fact, a nucleotide substitution (C→T) in exon 7 of the centromeric gene causes the skipping of exon 7 without altering the coded amino acid. Although homozygous deletion of the entire *SMN1* gene is responsible for the disease in almost all patients, hybrid genes involving *SMN1* and *SMN2* have been described.^{7,15-17} Therefore the absence of *SMN1* exon 7 is used for molecular diagnosis of the disease. Homozygous *SMN2* deletion is found in 5-9% of normal controls, and is not considered to be pathological.^{7,20}

Even before *SMN* dosage, the presence of more than two fragments corresponding to *SMN2* on the same chromosome had already been found in type II and III SMA.^{7,21} Three mechanisms have been suggested to explain the duplication of the *SMN2* gene on the same chromosome: (1) gene conversion, (2) illegitimate recombination, and (3) a two step mechanism, that is, deletion of an *SMN* gene followed by duplication of the remaining *SMN2* gene. Nevertheless, if gene conversion is the most probable event to explain the origin of duplication *in cis* of *SMN* genes, it has never been directly observed, except in cases of hybrid formation.^{7,15-17}

Since the introduction of SMA carrier detection methods for genetic counselling in families with SMA and relatives, risk assessment of recurrence has improved.¹⁸⁻²⁴ Carrier screening of parents is indicated before recommending prena-

Key points

- To establish the risk of recurrence in families with SMA correctly, we calculated the frequency of different *SMN1* genotypes in a large population.
- Characterisation of atypical genotypes in parents of patients with SMA shows a prevalence of *SMN1* duplication *in cis* associated with *SMN1* deletion of 3%.
- Studying the general population, we established a real carrier prevalence of 1/34, and deduced the frequency of the genotype carrying *SMN1* duplication associated with deletion to be 1/1000.
- Assessment of risk of recurrence in relatives depends on the molecular mechanism of the *SMN1* deletion and on genotypes of the proband and his or her parents.

tal diagnosis as de novo deletion has been found in 1% of them. However, the prevalence of parents carrying an *SMN1* duplication on the same chromosome associated with a deletion on the other one is currently unknown. Chen *et al*²² reported two duplications in 60 parents and relatives, whereas Mailman *et al*²³ reported one duplication in 100 parents. However, they could not establish the molecular mechanism for all the parents carrying two *SMN1* genes. It is extremely important to distinguish *SMN1* duplication to ensure correct interpretation of the genotypes of the families with SMA and their relatives.

Here, we report the results of carrier screening in parents of patients with SMA. To characterise the molecular mechanism of *SMN1* deletion in these families, we used two approaches: the *SMN* genotype was established first, followed by analysis of *SMN* flanking markers to interpret atypical genotypes. Thus, the prevalence of the genotype carrying two *SMN1* genes *in cis* was established. Also, we studied these genotypes in the general population and thus could establish true carrier prevalence. As we will show, the results permit more reliable assessment of risk of recurrence in families with SMA.

SUBJECTS AND METHODS

Gene dosage was performed in 202 parents of patients with SMA with homozygous deletion of *SMN1* exon 7, and in 375 people from the general population as described by Gérard *et al*.²⁰ Briefly, a two step method was used: firstly, measurement of all *SMN* genes, secondly, calculation of the *SMN1*/*SMN2* ratio by primer extension. Combining the results of the two techniques, we found the exact number of *SMN1* and *SMN2*

Abbreviations: SMA, spinal muscular atrophy; *SMN*, survival motor neurone gene

copies. This method is routinely used for carrier detection in the Genetic Centre of our hospital. More than 900 tests have been performed. The *SMN* genotype was established for all family members when one parent carrying two *SMN1* copies was detected. Analysis of markers flanking *SMN* in the duplicated region (C212 and Ag1-CA) distinguished *SMN1* duplication from de novo deletion in all cases. Amplification of these markers was performed with primers already described,^{11,26} modified using fluorescent dye in 5'. Products were visualised on an ABI PRISM 3100 DNA sequencer and analysed with GeneScan software (Applied Biosystems). When results were not fully informative, markers flanking the duplicated region were also studied (D5S637 and D5S629).²⁷ Informed consent was obtained from all tested patients.

RESULTS

Genotypes in SMA carriers

In the parent population, carrier status (only one *SMN1* gene detected) was confirmed in 193 (95.5%). Nine families had one parent carrying two *SMN1* copies (4.5%). We found two de novo deletions (1.0%), one on a maternally and one on a paternally derived chromosome. Six parents with *SMN1* duplication (3.0%) were detected. Finally, one de novo gene conversion on a paternally derived chromosome, converting an *SMN1* gene into *SMN2*, was found (0.5%) in which a normal chromosome carrying one *SMN1* and one *SMN2* was changed into a pathological chromosome carrying two *SMN2* copies. Three examples of the tested families are shown in fig 1A–C.

Genotypes in the general population

Most of the people from the general population were spouses of relatives with SMA who reported no neuromuscular diseases in their families. Among them, 11 heterozygous genotypes (one *SMN1* copy) were detected. These data set the carrier prevalence at 1/34 in the general population (table 1). Moreover, we estimated the prevalence of the genotype with *SMN1* duplication and *SMN1* deletion by multiplying the two chromosome frequencies as independent probabilities (table 2). About 1/1000 people with two *SMN1* genes is expected to carry a duplication plus deletion genotype. This result overlaps with the estimation deduced from our data in the carrier population. Indeed, if 3% of parents show *SMN1* duplication and carrier frequency in the general population is 1/34, a duplication plus deletion genotype should be found in 1/1133 people with a normal genotype, including two or more *SMN1* genes.

DISCUSSION

Carrier testing in parents and relatives of patients with SMA raises several problems about genotype interpretation and consequently assessment of recurrence risk. Carrier screening in parents of children with SMA with *SMN1* homozygous deletion is the method for detecting atypical genotypes. We found that 4.5% of parents carried two *SMN1* genes. We therefore characterised the molecular mechanism to establish the recurrence of risk in these families accurately. In fact, detecting a de novo deletion reduces the risk of recurrence in a family from 25% to less than 1% as germinal mosaicism has never been detected. Conversely, *SMN1* duplication is rarely considered when analysing families with SMA, although it is even more frequent than de novo deletion and leads to errors in genotype interpretation in relatives. Finally, conversion of *SMN1* into *SMN2* confirms that this mechanism induces duplication of *SMN* genes and de novo loss of the *SMN1* gene that is not detected by marker analysis.

When calculating the risk of recurrence in a couple where one is a carrier (one *SMN1* detected) and the other has a two *SMN1* genotype, three possibilities should be considered: (1) the presence of a subtle mutation in the person with the two *SMN1* genotypes (as in 1% of SMA parents^{7,10} estimated at 1%

Figure 1 Three examples of the tested families. The *SMN* genotype (T for telomeric gene *SMN1*, C for centromeric gene *SMN2*) and results of the marker analysis are given above each individual symbol. Patients asking for genetic counselling are indicated with arrows. Full rectangle shows *SMN1*, open rectangle shows *SMN2*. (A) Family 1 is an example of de novo deletion. This molecular mechanism could be suspected from the *SMN* gene dosage alone, mainly because dosage of a single *SMN2* in the proband was not coherent with the genotypes of her sister and parents. This result was confirmed by marker analysis, which showed the loss of the maternal contribution. To establish which chromosome was deleted in the affected child, we analysed D5S637 and D5S629, two markers flanking the duplicated region. We found that the deleted chromosome was the one carrying one *SMN1* gene. (B) In family 2, an *SMN1* duplication was detected in family member I.2. Gene dosage in family member II.2, being tested for carrier detection, showed two *SMN1* genes: only by combining results of both analyses (quantitative and marker) in all family members could we show that she was an *SMN1* deletion carrier. (C) Family 3 shows gene conversion. In family member II.2, the affected daughter, the five known divergences between *SMN1* and *SMN2* genes were tested by sequencing and restriction enzyme digestion and all showed *SMN2* origin.⁷ Moreover, she carried three *SMN2* genes, whereas her parents each carried only one *SMN2* gene. Paternity was confirmed. Analysis of markers C212 and Ag1-CA showed that both daughters had inherited the same chromosome from their father. Taking all these findings together, a de novo conversion of *SMN1* into *SMN2* was the only mechanism that could account for them. Therefore, we deduced a normal genotype for family member I.1.

Table 1 SMN1 genotypes in the general population

Genotypes	No SMN1	1 SMN1	2 SMN1	3 SMN1	4 SMN1
n=375	0 0 %	11 2.9 %	313 83.5 %	48 12.8 %	3 0.8 %

Table 2 Chromosomes in the general population

Chromosomes with	No in general population	Frequency
No SMN1	11	1.5%
1 SMN1	685	91.3%
2 SMN1	54	7.2%
Total	750	100%

of 1/34 in the general population); (2) the transmission of a de novo deletion (as in 1% of SMA parents, estimated at 1% of 1/34 in the general population); (3) the presence of a SMN1 duplication associated with a deletion (estimated at 1/1000 of the two SMN1 genotypes in the general population (our study)). Adding the risks derived from the three different molecular mechanisms, we calculate that the risk of recurrence will be 1/2520 for each pregnancy.

Intriguingly, carrier prevalence is higher than previously deduced from the disease prevalence: direct estimation sets carrier prevalence in the general population at 1/34 rather than 1/40. This finding increases the prevalence of SMN1 deleted genotypes to 1/4624. Underestimation of the frequencies of the extreme phenotypes could probably explain this discrepancy. The genotype with complete deletion of SMN1 and SMN2 should exist even if it has never been detected, as it would almost certainly be lethal, maybe at the embryonic stage, leading to early miscarriage. Conversely, non-symptomatic SMN1 homozygously deleted people have already been described.^{28,29} The question about the predictive value of this finding should be considered along with data concerning the age of evaluation and the clinical course of such patients.^{29,30} Only a few patients with SMA who have type IV and type III SMA with chronic evolution have been reported^{8,9} as they often remain undiagnosed because of the delayed onset of the disease and reasonable level of fitness. Further investigations are necessary to determine the exact prevalence of all SMA phenotypes. Studies about the level of expression of SMN2 in these patients will probably elucidate the phenotypic variability in SMA.

Authors' affiliations

V Cusin, B Gérard, D Chanterreau, J Elion, Laboratoire de Biochimie Génétique, Hôpital R Debré, 48 Boulevard Sérurier, 75019 Paris, France
O Clermont, Laboratoire de Génétique Bactérienne EA3105, Hôpital R Debré, 48 Boulevard Sérurier, 75019 Paris, France

Correspondence to: Dr V Cusin, Laboratoire de Biochimie Génétique, Hôpital R Debré, 48 Boulevard Sérurier, 75019 Paris, France; veronica.cusin@wanadoo.fr

REFERENCES

- Pearn J. Incidence, prevalence, and gene frequency studies of chronic childhood spinal muscular atrophy. *J Med Genet* 1978;**15**:409–13.
- Munsat TL. Workshop report. International SMA collaboration. *Neuromusc Disord* 1991;**1**:81.
- Melki J, Abdelhak S, Sheth P, Bachelot MF, Burlet P, Marcadet A, Aicardi J, Barois A, Carriere JP, Fardeau M, Fontan D, Ponsot G, Billette T, Angelini C, Barbosa C, Ferriere G, Lanzi G, Ottolini A, Babron MC, Cohen D, Hanauer A, Clerget-Darpoux F, Lathrop M, Munnich A, Frézal J. Gene for proximal spinal muscular atrophies maps to chromosome 5q. *Nature* 1990;**344**:767–8.
- Melki J, Sheth P, Abdelhak S, Burlet P, Bachelot MF, Lathrop M, Frézal J, Munnich A, the French spinal muscular atrophy investigators. Mapping of acute (type I) spinal muscular atrophy to chromosome 5q12-q14. *Lancet* 1990;**336**:271–3.
- Brzustowicz LM, Lehner T, Castilla LH, Penchaszadeh GK, Wilhelmson KC, Daniels RJ, Davies KE, Leppert M, Ziter F, Wood D, Dubowitz V, Zerres K, Hausmanova-Petrusewicz I, Ott J, Munsat TL, Gilliam TC. Genetic mapping of chronic childhood-onset spinal muscular atrophy to chromosome 5q11.2-q13.3. *Nature* 1990;**344**:540–1.
- Gilliam TC, Brzustowicz LM, Castilla LH, Lehner T, Penchaszadeh GK, Daniels RJ, Byth BC, Knowles J, Hislop JE, Shapira Y, Dubowitz V, Munsat TL, Ott J, Davies KE. Genetic homogeneity between acute and chronic forms of spinal muscular atrophy. *Nature* 1990;**345**:823–5.
- Lefebvre S, Burglen L, Reboullet S, Clermont O, Burlet P, Viollet L, Benichou B, Cruaud C, Millasseau P, Zeviani M, Le Paslier D, Frezal J, Cohen D, Weissenbach J, Munnich A, Melki J. Identification and characterization of a spinal muscular atrophy-determining gene. *Cell* 1995;**80**:155–65.
- Brahe C, Servidei S, Zappata S, Ricci E, Tonali P, Neri G. Genetic homogeneity between childhood-onset and adult-onset autosomal recessive spinal muscular atrophy. *Lancet* 1995;**346**:741–2.
- Clermont O, Burlet P, Lefebvre S, Burglen L, Munnich A, Melki J. SMN gene deletions in adult-onset spinal muscular atrophy. *Lancet* 1995;**346**:1712–3.
- Rodrigues NR, Owen N, Talbot K, Ignatius J, Dubowitz V, Davies KE. Deletions in the survival motor neuron gene on 5q13 in autosomal recessive spinal muscular atrophy. *Hum Mol Genet* 1995;**4**:631–4.
- Melki J, Lefebvre S, Burglen L, Burlet P, Clermont O, Millasseau P, Reboullet S, Benichou B, Zeviani M, Le Paslier D, Cohen D, Weissenbach J, Munnich A. De novo and inherited deletions of the 5q13 region in spinal muscular atrophies. *Science* 1994;**264**:1474–7.
- Raclin V, Veber PS, Burglen L, Munnich A, Melki J. De novo deletions in spinal muscular atrophy: implications for genetic counselling. *J Med Genet* 1997;**34**:86–7.
- Wirth B, Schmidt T, Hahnen E, Rudnik-Schoneborn S, Krawczak M, Muller-Myhsok B, Schonling J, Zerres K. De novo rearrangements found in 2% of index patients with spinal muscular atrophy: mutational mechanisms, parental origin, mutation rate, and implications for genetic counseling. *Am J Hum Genet* 1997;**61**:1102–11.
- Burglen L, Lefebvre S, Clermont O, Burlet P, Viollet L, Cruaud C, Munnich A, Melki J. Structure and organization of the human survival motor neurone (SMN) gene. *Genomics* 1996;**32**:479–82.
- Bussaglia E, Clermont O, Tizzano E, Lefebvre S, Burglen L, Cruaud C, Urtizberea JA, Colomer J, Munnich A, Baiget M, Melki J. A frame-shift deletion in the survival motor neuron gene in Spanish spinal muscular atrophy patients. *Nat Genet* 1995;**11**:335–7.
- Hahnen E, Schonling J, Rudnik-Schoneborn S, Zerres K, Wirth B. Hybrid survival motor neuron genes in patients with autosomal recessive spinal muscular atrophy: new insights into molecular mechanisms responsible for the disease. *Am J Hum Genet* 1996;**59**:1057–65.
- Cusco I, Barcelo MJ, del Rio E, Martin Y, Hernandez-Chico C, Bussaglia E, Baiget M, Tizzano EF. Characterisation of SMN hybrid genes in Spanish SMA patients: de novo, homozygous and compound heterozygous cases. *Hum Genet* 2001;**108**:222–9.
- McAndrew PE, Parsons DW, Simard LR, Rochette C, Ray PN, Mendell JR, Prior TW, Burghes AH. Identification of proximal spinal muscular atrophy carriers and patients by analysis of SMNT and SMNC gene copy number. *Am J Hum Genet* 1997;**60**:1411–22.
- Wirth B, Herz M, Wetter A, Moskau S, Hahnen E, Rudnik-Schoneborn S, Wienter T, Zerres K. Quantitative analysis of survival motor neuron copies: identification of subtle SMN1 mutations in patients with spinal muscular atrophy, genotype-phenotype correlation, and implications for genetic counseling. *Am J Hum Genet* 1999;**64**:1340–56.
- Gerard B, Ginet N, Matthijs G, Evrard P, Baumann C, Da Silva F, Gerard-Blanluet M, Mayer M, Grandchamp B, Elion J. Genotype determination at the survival motor neuron locus in a normal population and SMA carriers using competitive PCR and primer extension. *Hum Mutat* 2000;**16**:253–63.
- Campbell L, Potter A, Ignatius J, Dubowitz V, Davies K. Genomic variation and gene conversion in spinal muscular atrophy: implications for disease process and clinical phenotype. *Am J Hum Genet* 1997;**61**:40–50.
- Chen KL, Wang YL, Rennett H, Joshi I, Mills JK, Leonard DG, Wilson RB. Duplications and de novo deletions of the SMN1 gene demonstrated by fluorescence-based carrier testing for spinal muscular atrophy. *Am J Med Genet* 1999;**85**:463–9.
- Scheffer H, Cobben JM, Mensink RG, Stulp RP, van der Steege G, Buys CH. SMA carrier testing: validation of hemizygous SMN exon 7 deletion

- test for the identification of proximal spinal muscular atrophy carriers and patients with a single allele deletion. *Eur J Hum Genet* 2000;**8**:79–86.
- 24 **Saugier-Veber P**, Drouot N, Lefebvre S, Charbonnier F, Vial E, Munnich A, Frebourg T. Detection of heterozygous SMN1 deletions in SMA families using a simple fluorescent multiplex PCR method. *J Med Genet* 2001;**38**:240–3.
 - 25 **Mailman MD**, Hemingway T, Darsey RL, Glasure CE, Huang Y, Chadwick RB, Heinz JW, Papp AC, Snyder PJ, Sedra MS, Schafer RW, Abuelo DN, Reich EW, Theil KS, Burghes AH, de la Chapelle A, Prior TW. Hybrids monosomal for human chromosome 5 reveal the presence of a spinal muscular atrophy (SMA) carrier with two SMN1 copies on one chromosome. *Hum Genet* 2001;**108**:109–15.
 - 26 **DiDonato CJ**, Morgan K, Carpten JD, Fuerst P, Ingraham SE, Prescott G, McPherson JD, Wirth B, Zerres K, Hurko O, Wasmuth JJ, Mendell JR, Burghes AH, Simard L. Association between Ag1-CA alleles and severity of autosomal recessive proximal spinal muscular atrophy. *Am J Hum Genet* 1994;**55**:1218–29.
 - 27 **Clermont O**, Burlet P, Burglen L, Lefebvre S, Pascal F, McPherson J, Wasmuth JJ, Cohen D, Le Paslier D, Weissenbach J, Lathrop M, Munnich A, Melki J. Use of genetic and physical mapping to locate the spinal muscular atrophy locus between two new highly polymorphic DNA markers. *Am J Hum Genet* 1994;**54**:687–94.
 - 28 **Hahnen E**, Forkert R, Merke C, Rudnik-Schöneborn S, Schönling J, Zerres K, Wirth B. Molecular analysis of SMA candidate genes on chromosome 5q13 in autosomal recessive spinal muscular atrophy: evidence of homozygous deletions of the SMN gene in unaffected individuals. *Hum Mol Genet* 1995;**4**:1927–33.
 - 29 **Clermont O**, Bürglen L, Burlet P, Lefebvre S, Viollet L, Hausmanowa-Petrusewicz I, Munnich A, Melki J. Genetics analysis of unusual SMA pedigrees using SMN, NAIP genes and markers C212-C272. *Am J Hum Genet* 1995;**57**:A237.
 - 30 **Zerres K**, Rudnik-Schöneborn S, Forkert R, Wirth B. Genetic basis of adult-onset spinal muscular atrophy. *Lancet* 1995;**346**:1162.