ORIGINAL ARTICLE

HRPT2 mutations are associated with malignancy in sporadic parathyroid tumours

V M Howell, C J Haven, K Kahnoski, S K Khoo, D Petillo, J Chen, G J Fleuren, B G Robinson, L W Delbridge, J Philips, A E Nelson, U Krause, K Hammje, H Dralle, C Hoang-Vu, O Gimm, D J Marsh, H Morreau, B T Teh

J Med Genet 2003;40:657–663

Background: Hyperparathyroidism is a common endocrinopathy characterised by the formation of parathyroid tumours. In this study, we determine the role of the recently identified gene, *HRPT2*, in parathyroid tumorigenesis.

Methods: Mutation analysis of *HRPT2* was undertaken in 60 parathyroid tumours: five HPT-JT, three FIHP, three MEN 1, one MEN 2A, 25 sporadic adenomas, 17 hyperplastic glands, two lithium associated tumours, and four sporadic carcinomas. Loss of heterozygosity at 1q24-32 was performed on a subset of these tumours.

Results: *HRPT2* somatic mutations were detected in four of four sporadic parathyroid carcinoma samples, and germline mutations were found in five of five HPT-JT parathyroid tumours (two families) and two parathyroid tumours from one FIHP family. One HPT-JT tumour with germline mutation also harboured a somatic mutation. In total, seven novel and one previously reported mutation were identified. "Two-hits" (double mutations or one mutation and loss of heterozygosity at 1q24-32) affecting *HRPT2* were found in two sporadic carcinomas, two HPT-JT-related and two FIHP related tumours.

Conclusions: The results in this study support the role of *HRPT2* as a tumour suppressor gene in sporadic parathyroid carcinoma, and provide further evidence for *HRPT2* as the causative gene in HPT-JT, and a subset of FIHP. In light of the strong association between mutations of *HRPT2* and sporadic parathyroid carcinoma demonstrated in this study, it is hypothesised that *HRPT2* mutation is an early event that may lead to parathyroid malignancy and suggest intragenic mutation of *HRPT2* as a marker of malignant potential in both familial and sporadic parathyroid tumours.

See end of article for authors' affiliations

Correspondence to: Dr B T Teh; bin.teh@vai.rog, Dr H Morreau; j-morreau@lumc.nl, or Dr D J Marsh; Debbie_Marsh@med.usyd.edu.au

Revised version received 26 June 2003
Accepted for publication 26 June 2003

Hyperparathyroidism (HPT) is one of the most common endocrinopathies, believed to affect approximately three individuals per 1000 adults.¹ HPT is characterised by the formation of parathyroid tumours and, if left untreated, patients develop bone disease, renal stones, and neuromuscular dysfunction. The majority of tumours are sporadic, but approximately 5% are associated with the autosomal dominant hereditary cancer syndromes multiple endocrine neoplasia type 1 and 2A (MEN 1 and 2A), familial isolated hyperparathyroidism (FIHP), and hyperparathyroidism-jaw tumour syndrome (HPT-JT).²

Sporadic HPT may occur as primary, secondary, or tertiary disease. Primary HPT can be attributed to a single adenoma in 80–85% of cases, multiglandular hyperplasia in 15–20% of cases, and carcinoma in less than 1% of cases.³ Secondary HPT arises in response to 1,25-dihydroxyvitamin D₃ deficiency, hyperphosphatemia, or hypocalcemia, due to renal failure, and presents as multiglandular hyperplasia. Tertiary HPT is defined as autonomous parathyroid hyperfunction in a patient with a previously well documented history of secondary HPT. Its presentation has been described as nodular hyperplasia or multiple adenoma. HPT may also arise in response to lithium treatment as a therapy for bipolar disorder.⁴

Our understanding of the molecular basis of parathyroid tumourigenesis has increased over the past 10 years. It has been clearly established that increased expression of the oncogene *cyclin D1* (formerly named *PRAD1*) is associated with the formation of parathyroid hyperplastic glands, adenomas and carcinomas.^{5–7} The MEN 1 gene, *MEN1*, located at 11q13,^{8–9} has also been shown to play a major role

in both familial^{10–12} and sporadic^{13–16} parathyroid tumourigenesis, functioning as a tumour suppressor. However the genetic basis of many sporadic and familial parathyroid tumours is still unknown. Parathyroid carcinoma, for example, is not a feature of MEN 1, and neither loss of heterozygosity (LOH) at 11q13, nor mutations in *MEN1* have been reported.^{17–18} The tumour suppressor gene, *Retinoblastoma (Rb1)*, located at 13q14, was suggested to have a primary role in parathyroid carcinogenesis following the finding of allelic loss of *Rb1* in the majority of parathyroid carcinomas but rarely in benign parathyroid tumours.^{19–20} However, subsequent immunohistochemical studies found decreased Rb1 expression to be unreliable as a marker of malignancy.^{21–23}

Recently, germline mutations in a newly identified gene, *HRPT2* (AF312865, Hs.5722), previously identified as *Chromosome 1 open reading frame 28*²⁴ have been identified in 14 of 24 HPT-JT kindreds and one of two FIHP kindreds.²⁵ *HRPT2* maps to 1q25, consists of 17 exons, containing 1596 nucleotides and encodes a 531 amino acid protein termed parafibromin with unknown function.²⁵ Patients with HPT-JT present in childhood or early adulthood with HPT, and are also at risk of developing fibro-osseous jaw tumours and renal lesions.^{26–27} While sporadic parathyroid carcinoma is rare, the incidence of parathyroid carcinoma is reported to be

Abbreviations: FIHP, familial isolated hyperparathyroidism; HPT, hyperparathyroidism; HPT-JT, hyperparathyroidism-jaw tumour syndrome; LOH, loss of heterozygosity; MEN, multiple endocrine neoplasia

15% in HPT-JT.² The majority of parathyroid tumours in these patients are aggressive, occasionally recurrent adenomas, notable also for their cystic histology.²⁸ LOH studies at 1q24-32 have identified allelic loss in some, but not all, HPT-JT associated tumours, suggestive of a tumour suppressor role for *HRPT2*.^{29–30} Similar loss of the *HRPT2* locus in tumours from members of a FIHP family has been reported.³¹ Studies in sporadic parathyroid tumours report that 9–13% of adenomas have LOH at 1q24-32.^{14–32–33} In addition to germline *HRPT2* mutations in HPT-JT and FIHP kindreds, Carpten *et al* also detected somatic *HRPT2* mutations in two of 47 sporadic cystic parathyroid adenomas.²⁵ One of these was also found to have LOH at 1q24-32.²⁵ LOH at this locus has not been identified in lithium-associated parathyroid tumours³⁴ or secondary HPT.³³

To determine the role of *HRPT2* in parathyroid tumourigenesis, mutation analysis was undertaken in both sporadic and familial parathyroid neoplasms. A subset of tumours was also assessed for LOH at the *HRPT2* locus, 1q24-32.

MATERIALS AND METHODS

Subjects and samples

Sixty parathyroid samples were obtained from subjects (31 female and 29 male) who underwent parathyroidectomy for hyperparathyroidism at Royal North Shore Hospital, Sydney, Australia; Leiden University Medical Centre, Leiden, Netherlands; or Martin Luther University, Halle-Wittenberg, Germany. Peripheral blood samples were also available from 42 of these patients. Patients gave informed consent according to protocols approved by each centre's human ethics committee.

The tumours were classified as familial or sporadic, including lithium-associated. Further classification of adenoma, hyperplasia (secondary or tertiary), or carcinoma was established according to detailed WHO guidelines.³⁵ Twelve familial tumour specimens were collected. Three were from two FIHP families, five from two HPT-JT families, three from three MEN 1 families and one from an MEN 2A patient. Orthopantomography of the jaw and renal ultrasound were performed on all affected members of the FIHP families. No case of jaw or renal tumour was found. Twenty-five sporadic adenomas, two lithium-associated tumours, 11 secondary and six tertiary hyperplastic glands, as well as four parathyroid carcinomas were also collected. The percentage of neoplastic tissue in each sample was assessed histologically from a paraffin embedded representative piece of tissue. All samples were composed of at least 70% neoplastic cells.

A panel of 65 anonymised peripheral blood samples collected from healthy volunteers constituted the normal germline DNA panel.

DNA/RNA preparation

Parathyroid tissue was frozen in liquid nitrogen immediately after surgical removal and stored at -70°C or below. Peripheral blood was collected into EDTA anti-coagulant tubes and stored at -70°C . DNA was extracted from the frozen tissue and peripheral blood leucocytes according to standard procedures. RNA was extracted from frozen tissue using TRI Reagent (Sigma-Aldrich Corporation, St Louis, MO) according to the manufacturer's protocol.

LOH studies

Where DNA was available from matched tumour and blood samples, allelic deletion of the chromosome 1q24-32 region flanking *HRPT2* was assessed using a selection of the following microsatellite markers: centromeric-*DIS218-DIS238-DIS422-DIS2625-DIS081-HRPT2DIS533-DIS2757-DIS2794-DIS477*-telomeric.^{25–36} PCR was performed in a 7.5 μL reaction volume containing 0.17 μM each of

HEX-labelled forward and unlabelled reverse primer (Invitrogen, Life Technologies, Carlsbad, CA), 4 mM MgCl_2 , 0.3 units AmpliTaq Gold polymerase and 1 $\times$ Buffer II (Applied Biosystems, Foster City, CA), 250 μM dNTPs (Invitrogen, Life Technologies), and 15 ng of genomic DNA. Amplification was performed in a DNA Engine Tetrad (MJ Research, Incline Village, NV) with an initial denaturation of 95°C for 10 min, followed by 10 cycles of 94°C for 15 seconds, 55°C for 15 seconds, 72°C for 30 seconds, and 20 cycles of 89°C for 15 seconds, 55°C for 15 seconds, 72°C for 30 seconds, and a final extension at 72°C for 10 min. The resulting microsatellite PCR products for each specimen were then pooled, and 1 μL was added to 10 μL of Hi-Di formamide (Applied Biosystems) and 0.2 μL of ROX 400HD size standard (Applied Biosystems), denatured at 95°C for 5 min and loaded into an ABI Prism 3700 Genetic Analyser (Applied Biosystems). Assessment of LOH was performed using Genescan v. 3.7 and Genotyper v. 3.7 software (Applied Biosystems). LOH was defined according to the following formula: LOH index = $(T_2/T_1)/(N_2/N_1)$, where T was the tumour sample, N was the matched normal sample, 1 and 2 were the intensities of smaller and larger alleles, respectively. If the ratio was <0.5 or >1.5 , the result was determined to be LOH.³³

HRPT2 mutation analysis

Initially, normal thyroid tissue RNA was sequenced in 4 overlapping segments between 5'UTR and 3'UTR to clarify the intron/exon boundaries and confirm the published sequence of the recently identified *HRPT2* gene.²⁵ cDNA was synthesized using the SUPERScript First-Strand Synthesis System for RT-PCR (Invitrogen, Life Technologies) and amplified in a 20 μL reaction containing 200 ng of cDNA, 200 μM dNTPs (Invitrogen, Life Technologies), 1.5–4 mM MgCl_2 , 1 μM of each primer, 1 unit of Platinum *Taq* DNA Polymerase and 1 $\times$ PCR buffer (Invitrogen, Life Technologies). Following an initial denaturation at 94°C for 5 minutes, 35 cycles of 15 second steps at 94°C , 55 – 60°C (depending on the primer set) and 72°C were performed in a DNA Engine Dyad (MJ Research), finishing with an extension at 72°C for 10 minutes. The PCR products were purified using the QIAquick PCR Purification kit (QIAGEN Pty Ltd, Clifton Hill, Vic, Australia), cycle sequenced using the ABI PRISM BigDye Terminator Cycle Sequencing Ready Reaction kit (Applied Biosystems) and sequenced on an ABI PRISM 3700 Genetic Analyser, according to the manufacturer's recommended protocol. The organisation of the gene is detailed in table 1 and primer sequences are available on request.

Sequencing of *HRPT2* was then performed for all tumour samples. All 17 exons and the intron-exon boundaries were amplified in 15 separate reactions for each sample. PCR was performed in 40 μL reactions containing 75 ng of template DNA, 200 μM dNTPs, 1.5 mM MgCl_2 , 0.5 μM each primer, 1.5 units of *Taq* DNA Polymerase, and 1 $\times$ reaction buffer (Invitrogen, Life Technologies). Following an initial denaturation at 94°C for 5 minutes, 35 cycles of 30 second steps at 94°C , 55°C , and 45 seconds at 72°C were performed, finishing with an extension at 72°C for 10 minutes in a DNA Engine Tetrad. Primer sequences are available on request. The PCR products were purified through a Multiscreen PCR Filter plate (Millipore Corp., Billerica, MA) and sequenced as above.

Where matched blood was available, the status of mutations identified in the tumours were determined as either somatic or germline by sequencing of the germline DNA.

RNA was available from three tumours to test whether there were aberrant splicing products of the gene. cDNA was synthesised, amplified and sequenced as above.

Table 1 *HRPT2* (Unigene cluster HS.5722) exon/intron organisation

Exon							Intron	
No	Bp	Nucleotides	Amino acids	Genomic sequence no*	3'splice acceptor region	5'splice donor region	No	Kb
1	131	1–131	1–44	2583–2713	agggggggaag/ATG	TGGGG/gtaag	1	2.8
2	106	132–237	45–79	5494–5599	atttcag/GACT	AGCT/gtaag	2	5.0
3	70	238–307	80–102	10,556–10,625	gttttag/ACTG	GCGT/gtgag	3	5.1
4	63	308–370	103–123	15,773–15,835	cttcag/CAAC	CAAG/gtatg	4	0.1
5	53	371–423	124–141	15,919–15,971	ctttatag/TCAA	ATTGAG/gtaaa	5	2.5
6	89	424–512	142–170	18,467–18,555	atttcag/GATG	ATTAG/gtaag	6	3.7
7	217	513–729	171–243	22,232–22,448	cttttag/GTCT	GGAAAG/gtaatt	7	5.8
8	99	730–828	244–276	28,249–28,347	atttttag/AATTTT	TGTG/gtaag	8	2.4
9	79	829–907	277–302	30,686–30,764	attttttacag/GATC	GAAG/gcaagt	9	2.0
10	65	908–972	303–324	32,762–32,826	cttttttcgtag/AAAC	AACG/gtaag	10	51.3
11	58	973–1030	325–343	84,177–84,234	ttttaag/GAGGTG†C	CCAG/gtag	11	8.2
12	36	1031–1066	344–355	92,447–92,482	tcacag/TTTCT	GAAAG/gtga	12	0.3
13	88	1067–1154	356–385	92,772–92,859	ctttatag/GATC	TGAA/gtaag	13	20.5
14	162	1155–1316	385–438	113,375–113,536	gtttttcaag/ATT	ACTG/gtaag	14	3.1
15	101	1317–1417	439–472	116,638–116,738	cttgtag/GGAC	TAAAA/gtaag	15	13.4†
16	142	1418–1559	473–519	9248–9389	tatag/TTAA	ACAG/gtaat	16	0.8
17	37	1560–1596	520–531	10,194–10,230	attttttcag/GTAC	CTGA/attatt	3'UTR	0.7

*The genomic sequence no. is taken from clones RP11-239J11 (AL390863/AF312865) and RP11-185C19 (AL139133/AF312865).

†Position of overlap of these clones.

‡Base found to be discrepant with the published sequence is in bold.

Polymorphism scanning of the normal DNA panel

All amplifications were performed with 1 µM of each primer (Invitrogen, Life Technologies), 200 µM dNTPs (Invitrogen, Life Technologies), 2 units of AmpliTaq Gold polymerase (Applied Biosystems), 1 × Buffer II (Applied Biosystems), 1.5–4 mM MgCl₂, and cycling conditions as stated for PCR of cDNA above, with an annealing temperature of 55°C, unless otherwise stated. Primer sequences are available on request.

RsaI digest for detection of c.33C→T (Y11Y)

DNA was amplified in a 20 µL volume with exon 1 primers and 1 unit of AmpliTaq Gold polymerase. Four microlitres of the resulting 228 bp amplicon was digested with 1 unit of restriction endonuclease *RsaI* (Fermentas AB, Vilnius, Lithuania) in 1 × Buffer Y+/Tango (Fermentas AB) with a final volume of 8 µL, and incubated at 37°C for 8 hours. One microlitre of the resulting digest was mixed with 1 µL of loading dye (5 g/L Bromophenol Blue, 12.5 mM EDTA, 50% formamide) and electrophoresed in a Corbett Research Gel-Scan 2000 Real-time Electrophoresis System (Corbett Research, Mortlake, NSW, Australia) through a 5% polyacrylamide, 0.6 × TBE gel under suggested non-denaturing conditions. Wild type amplicons were distinguished by the presence of bands at 109 bp and 119 bp. Heterozygotes contained an additional band at 228 bp.

dHPLC detection of polymorphisms

DNA was amplified in a 50 µL volume using exon 2, 7, or 13 primers and following amplification the resulting products underwent enhancement of heteroduplex formation. This involved a denaturation at 94°C for 5 min immediately followed by a slow ramp over 35 cycles, starting at 94°C and decreasing 2°C/1 min cycle, finishing with 1 min incubation at 25°C. Five microlitres of amplicon was then injected into the flowpath of a DNasep-MD cartridge in the WAVE-MD Mutation Detection System Model 2000 denaturing HPLC (Transgenomic, Omaha, NE) under WAVE-MD Standard Method gradient conditions and WAVEMAKER (Transgenomic) predicted temperatures for each amplicon. Eluted fragments were detected by the system's UV detector and analysed as chromatograms. Wild type amplicons eluted as a single homoduplex peak whereas heterozygous samples presented with up to four peaks representing homo- and heteroduplexes. For the

IVS2+28C→T variant, homozygotes for either the C or T allele were mixed with an equal aliquot of an exon 2 amplicon homozygous for the C allele, denatured, slowly reannealed and re-injected into the dHPLC to determine which allele was present.

Statistical analysis

The χ^2 contingency test was performed to compare the occurrence of mutations between four cases of sporadic carcinomas and 44 cases of sporadic non-carcinomas. A *p* value less than 0.05 is considered as significant.

RESULTS

HRPT2 mutation analysis

HRPT2 mutations were detected in the DNA from four of four sporadic parathyroid carcinoma samples, five of five HPT-JT parathyroid tumours (two families) and parathyroid tumours from one of two FIHP families (fig 1). No mutations were detected in any of the other 49 tumours sequenced. Eight different *HRPT2* mutations were detected. With the exception of L64P, that has been previously published,²⁵ all the mutations identified were novel (fig 2). One of the mutations (c.76delA) was detected both in a sporadic carcinoma and in a HPT-JT family. Where available, matched constitutive DNA was analysed to determine the germline versus somatic nature of these mutations (fig 1). HPT-JT tumour #1613 harboured both a germline and a somatic *HRPT2* mutation, and two somatic *HRPT2* mutations were detected in carcinoma #2077.

LOH studies

LOH analysis of 1q24-32 was performed on 42 of the 60 tumours. All samples tested were informative for at least one of the markers (data not shown). LOH was detected in six samples: one sporadic carcinoma (#10977), one HPT-JT tumour (#1765), two FIHP tumours (#4 and #54) and two sporadic adenomas (#9 and #101) (fig 1). LOH at 1q24-32 has been previously reported in two of these samples (#9³⁴ and #4³¹). LOH was confirmed in sporadic carcinoma sample #10977, HPT-JT sample #1765 and FIHP samples #4 and #54 by the presence of only the mutant allele in tumour DNA sequence. Retention of heterozygosity was demonstrated in the remaining 18 tumours not assessed for LOH by the

Figure 1 Mutations detected within *HRPT2*, and LOH analysis of flanking markers. "Age" is the age in years at the time of parathyroidectomy. *HRPT2* mutations detected are listed as nucleotide changes and amino acid changes where relevant. Where two mutations were detected in the same patient, both are listed. Germline mutations are denoted "[G]", and somatic mutations are denoted "[S]". For the microsatellite markers, see key in figure. Blank spaces indicate LOH testing not performed (germline DNA not available). *The approximate location of *HRPT2* within the 1q24-32 region. The results in this shaded column represent retention or loss of heterozygosity at the *HRPT2* locus by combined microsatellite and intragenic mutation results. †*D1S218* and *D1S477* have been reported previously.³¹ ‡Previously reported as having LOH at 1q and 1p and CGH loss at 1q31-qter.³⁴

presence of a heterozygous mutation or polymorphism in *HRPT2*.

Polymorphisms

Five intronic and one exonic sequence variants, five of which were also identified in a screen of 65 normal subjects, were identified in parathyroid lesions, and where available,

matched constitutive DNA (table 2). An additional variant, not detected in the tumours, was found in the normal panel (fig 3). Twenty six of the 54 parathyroid tumour samples not displaying LOH were heterozygous for the common polymorphism, IVS2+28C→T, confirming retention of heterozygosity for *HRPT2*. A dinucleotide repeat variant in intron 7, IVS7+33(GA)₈ (wild type IVS7+33 (GA)₉) was found in five

Figure 2 Distribution of *HRPT2* mutations and polymorphisms. Exons 8–12 inclusive, 15, and 16, have not been shown for clarity. Underlined variants are from this study, others are from Carpten *et al.*²⁵ Variants predicted to be polymorphisms are italicised.

Table 2 HRPT2 predicted polymorphisms found in this study			
Polymorphism	Heterozygote frequency in tumour DNA* (allele frequency)	Classification of tumours harbouring a polymorphism	Heterozygote frequency in germline DNA from a normal panel (allele frequency)
Y11Y (c.33C→T)	1 in 56 (0.01)	1 sporadic adenoma [G]	0 in 65
IVS2+28delCCTA	2 in 56 (0.02)	1 secondary hyperplasia [G] 1 tertiary hyperplasia [?]	5 in 65 (0.04)
IVS2+28C→T	26 in 56 (Heterozygous) (0.30)	all tumour types [G]	33 in 65 (Heterozygous) (0.32)
IVS7+33(GA) ₈	4 in 56 (Homozygous for T) 4 in 56 (0.04)	1 lithium [G] 1 secondary hyperplasia [G] 1 MEN 1 [?] 1 FIHP† (2 affected family members [G])	4 in 65 (Homozygous for T) 1 in 65 (0.01)
IVS12-86C→T	3 in 56 (0.03)	1 sporadic adenoma [?] 2 secondary hyperplasia [?]	3 in 65 (0.02)
IVS12-109T→G	5 in 56 (0.05)	1 secondary hyperplasia [G] 1 tertiary hyperplasia [?] 3 sporadic adenomas [?]	6 in 65 (0.05)
IVS13+20A→C	0 in 56		1 in 65 (0.01)

*Only 1 sample from each family was included in the heterozygote frequency estimation.
[G]Present also in germline.
[?]Germline status not ascertained.
†The variant IVS7+33(GA)₈ was found in tumour and in germline in two affected members of the same family. Both tumours demonstrated LOH at 1q24-32 and the polymorphism was found on the mutated allele in tumour DNA.

tumours. We were unable to detect aberrant splicing in transcripts generated from three of these tumours (#4, #54, and #76).

Statistical analysis

A χ^2 contingency test showed that the occurrence of mutation in sporadic parathyroid carcinomas is highly significant compared with sporadic non-carcinomas ($\chi^2 = 48$; $p < 0.001$).

Figure 3 Denaturing HPLC chromatograms of HRPT2 polymorphisms. The chromatograms plot the absorbance at 260 nm in mVolts v elution time in minutes for the various amplicons and show the different profiles that distinguish between the wild type (WT) and various polymorphisms for each amplicon. Wild type amplicons elute as a single homoduplex peak, whereas samples heterozygous for a polymorphism present with up to four peaks, representing homo- and heteroduplexes. Samples harbouring both polymorphisms in intron 2 (A) are also distinguishable from those heterozygous for a single intron 2 polymorphism.

DISCUSSION

Here we report the first finding of HRPT2 mutations in 100% (four of four) of sporadic parathyroid carcinomas studied (fig 1). All are predicted to prematurely truncate the parafibromin protein. No HRPT2 mutations were found in any of the 44 sporadic, non-carcinoma samples in this study. These results demonstrate a strong association between intragenic mutation of HRPT2 and malignancy in parathyroid tumours ($p < 0.001$). Furthermore, our findings of two somatic mutations as well as one mutation with LOH at 1q24-32 in two of four carcinomas are consistent with Knudson's "two-hit" hypothesis, and suggest the role for HRPT2 as a tumour suppressor gene in sporadic parathyroid tumorigenesis.

Our finding of germline mutations in two of two HPT-JT families and confirmation of a germline mutation in one of two FIHP families support the conclusions of Carpten *et al*,²⁵ that HRPT2 is the causative gene in HPT-JT and in a subset of FIHP (fig 1). Of additional significance was our finding of the "second-hit" in four of the seven samples with germline mutations. Three familial tumour samples demonstrated loss of the wild-type allele, and one tumour contained a somatic frameshift mutation (fig 1). These results provide evidence of a role for HRPT2 as a tumour suppressor gene in familial as well as sporadic parathyroid tumorigenesis.

The two germline mutations in the HPT-JT families are, like those found in the sporadic carcinomas, predicted to prematurely truncate the protein, whereas the FIHP mutation is a missense mutation of unknown consequence. Whether there is a phenotype/genotype correlation between mutation type and the presence of parathyroid disease alone, or in conjunction with jaw tumours or renal lesions (that is, FIHP v HPT-JT) will require a larger cohort of FIHP families with HRPT2 mutations to establish. Similarly, it is uncertain whether the severity or number of "hits" affecting HRPT2 determines the presentation of familial tumours as benign, cystic, or malignant. An alternative scenario may be that intragenic mutation of HRPT2 is an early event in a subset of patients with parathyroid lesions, and that additional, yet unknown events are required for the progression to malignancy. It is noteworthy that HRPT2 mutations have been previously detected in two of 47 (4%) sporadic cystic adenomas, and one (2%) had additional 1q24-32 LOH.²⁵ Given the strong association demonstrated between

intragenic mutation of *HRPT2* and malignancy in the current study, such a finding in a benign tumour might be considered a marker of malignant potential and long term monitoring is suggested for these patients.

Twenty four different *HRPT2* mutations have now been reported,²⁵ including seven in this paper (fig 2). Over 80% (20 of 24) of *HRPT2* mutations are located in exons 1, 2, or 7 or flanking intronic sequences. Exon 1 contains 42% (10 of 24) of all mutations detected to date, and exon 2 appears to contain a mutation hot spot at c.165C. Exon 7 is by far the largest exon, hinting at an important role for this exon, and to date, has been shown to harbour 21% (five of 24) of all *HRPT2* mutations detected. This exon contains a repeat element (AG)₅, and three different frameshift mutations altering this motif have been detected. This study also detected seven apparent polymorphisms. We found these, in particular, IVS2+28C→T, to be useful markers for exclusion of LOH at the *HRPT2* locus.

The consistent detection of LOH at 1q24-32 in 9–13% of sporadic adenomas^{14 32 33} is interesting. Our finding of LOH in two of 25 (12.5%) sporadic adenomas is in agreement with these previous reports, as is our finding of no LOH in 17 hyperplasia or two lithium-associated tumours.^{33 34} We have previously shown that extensive loss of both arms of chromosome 1 occurs in approximately 10% of sporadic adenomas,^{14 32} including adenoma #9³⁴ from this study. We did not find a *HRPT2* mutation on the remaining allele of either of the sporadic adenomas with LOH at 1q24-32 in this study and suggest that the LOH at 1q24-32 in the majority of sporadic adenomas may in fact be targeted at a tumour suppressor gene elsewhere on chromosome 1. To date, there is no evidence that LOH at 1q24-32 alone confers a higher likelihood of malignancy.

In conclusion, the results in this study support a role for *HRPT2* as a tumour suppressor gene in agreement with Knudson's "two-hit" hypothesis in sporadic parathyroid carcinoma. These results also provide further evidence for *HRPT2* as the causative gene in HPT-JT, and a subset of FIHP. Efficient detection of *HRPT2* mutations should begin with exons 1, 2, and 7, which together harbour 80% of mutations found to date. In light of the strong association between mutations of *HRPT2* and sporadic parathyroid carcinoma demonstrated in this study, we hypothesise that *HRPT2* mutation is an early event that may lead to parathyroid malignancy and therefore suggest intragenic mutation of *HRPT2* as a marker of parathyroid malignant potential in both familial and sporadic tumours. Further studies will clarify the exact nature of the relationship between intragenic mutation of *HRPT2* and parathyroid malignancy.

ACKNOWLEDGEMENTS

We are indebted to A-L Richardson, L Cheung and G Theodosopoulos, from the Kolling Institute of Medical Research, for their assistance with the DNA extractions; and to surgeon, J Kievit, (Department of Surgery) and endocrinologist JWA Smit, (Department of Endocrinology), both from Leiden University Medical Centre for assistance with the collection and clinical classification of tumours. Supported by a Dora Lush Postgraduate Research Scholarship, National Health and Medical Research Council (NHMRC), Australia; a Cancer Memorial Research Scholarship, RNSH, Australia; and a Northern Sydney Health Ramsay HealthCare Study Fellowship (VMH); by an R.D. Wright Fellowship, NHMRC, Australia (DJM); in part by the Pinguin-Stiftung (CH-V); and by the Van Andel Foundation, USA.

Authors' affiliations

V M Howell*, **C J Haven***, **K Kahnoski**, **S K Khoo**, **D Petillo**, **J Chen**, **B T Teh**, Laboratory of Cancer Genetics, Van Andel Research Institute, 333 Bostwick NE, Grand Rapids, Michigan, 49503, USA
V M Howell, **B G Robinson**, **D J Marsh**, Kolling Institute of Medical

Research, Royal North Shore Hospital (RNSH), St Leonards, NSW, Australia, 2065, and Department of Molecular Medicine, University of Sydney, Sydney, NSW, Australia

C J Haven, **G J Fleuren**, **H Morreau**, Department of Pathology, Leiden University Medical Centre, PO Box 9600, L1Q, NL-2300, Leiden, Netherlands

L W Delbridge, Department of Surgery, RNSH, Sydney, NSW, Australia

J Philips, Department of Anatomical Pathology, Pacific Laboratory Medicine Services, RNSH, Sydney, NSW, Australia, and Department of Pathology, University of Sydney, Sydney, NSW, Australia

A E Nelson, Garvan Institute of Medical Research, Darlinghurst, Sydney, NSW, Australia

K Hammje, **H Dralle**, **C Hoang-Vu**, **O Gimm**, Department of General, Visceral and Vascular Surgery, Martin Luther University, Halle-Wittenberg, Germany

U Krause, Department of Pathology, Martin Luther University, Halle-Wittenberg, Germany

*VMH and CJH contributed equally to this work

REFERENCES

- Adami S, Marcocci C, Gatti D. Epidemiology of primary hyperparathyroidism in Europe. *J Bone Miner Res* 2002;17(Suppl 2):N18–23.
- Marx SJ, Simonds WF, Agarwal SK, Burns AL, Weinstein LS, Cochran C, Skarulis MC, Spiegel AM, Libutti SK, Alexander HR Jr, Chen CC, Chang R, Chandrasekharappa SC, Collins FS. Hyperparathyroidism in hereditary syndromes: special expressions and special managements. *J Bone Miner Res* 2002;17(Suppl 2):N37–43.
- Marx SJ. Hyperparathyroid and hypoparathyroid disorders. *N Engl J Med* 2000;343:1863–75.
- Garfinkel PE, Ezrin C, Stancer HC. Hypothyroidism and hyperparathyroidism associated with lithium. *Lancet* 1973;2:331–2.
- Arnold A, Kim HG, Gaz RD, Eddy RL, Fukushima Y, Byers MG, Shows TB, Kronenberg HM. Molecular cloning and chromosomal mapping of DNA rearranged with the parathyroid hormone gene in a parathyroid adenoma. *J Clin Invest* 1989;83:2034–40.
- Hsi ED, Zukerberg LR, Yang WL, Arnold A. Cyclin D1/PRAD1 expression in parathyroid adenomas: an immunohistochemical study. *J Clin Endocrinol Metab* 1996;81:1736–9.
- Vasef MA, Brynes RK, Sturm M, Bromley C, Robinson RA. Expression of cyclin D1 in parathyroid carcinomas, adenomas, and hyperplasias: a paraffin immunohistochemical study. *Mod Pathol* 1999;12:412–6.
- Lemmens I, Van de Ven WJ, Kas K, Zhang CX, Giraud S, Wautot V, Buisson N, De Witte K, Salandre J, Lenoir G, Pugeat M, Calender A, Parente F, Quincey D, Gaudray P, De Wit MJ, Lips CJ, Hoppener JW, Khodaei S, Grant AL, Weber G, Kytola S, Teh BT, Farnebo F, Thakker RV, et al. Identification of the multiple endocrine neoplasia type 1 (MEN1) gene. The European Consortium on MEN1. *Hum Mol Genet* 1997;6:1177–83.
- Chandrasekharappa SC, Guru SC, Manickam P, Olufemi SE, Collins FS, Emmert-Buck MR, Debelenko LV, Zhuang Z, Lubensky IA, Liotta LA, Crabtree JS, Wang Y, Roe BA, Weisemann J, Boguski MS, Agarwal SK, Kester MB, Kim YS, Heppner C, Dong Q, Spiegel AM, Burns AL, Marx SJ. Positional cloning of the gene for multiple endocrine neoplasia-type 1. *Science* 1997;276:404–7.
- Giraud S, Zhang CX, Serova-Sinilnikova O, Wautot V, Salandre J, Buisson N, Waterlot C, Bauters C, Porchet N, Aubert JP, Emy P, Cadiot G, Delemer B, Chabre O, Niccoli P, Leprat F, Duron F, Emperauger B, Cougard P, Goudet P, Sarfati E, Riou JP, Guichard S, Rodier M, Calender A, et al. Germ-line mutation analysis in patients with multiple endocrine neoplasia type 1 and related disorders. *Am J Hum Genet* 1998;63:455–67.
- Teh BT, Kytola S, Farnebo F, Bergman L, Wong FK, Weber G, Hayward N, Larsson C, Skogseid B, Beckers A, Phelan C, Edwards M, Epstein M, Alford F, Hurley D, Grimmond S, Silins G, Walters M, Stewart C, Cardinal J, Khodaei S, Parente F, Tranebjaerg L, Jorde R, Salmela P, et al. Mutation analysis of the MEN1 gene in multiple endocrine neoplasia type 1, familial acromegaly and familial isolated hyperparathyroidism. *J Clin Endocrinol Metab* 1998;83:2621–6.
- Villablanca A, Wassif WS, Smith T, Hoog A, Vierimaa O, Kassem M, Dwight T, Forsberg L, Du Q, Learoyd D, Jones K, Stranks S, Juhlin C, Teh BT, Carling T, Robinson B, Larsson C. Involvement of the MEN1 gene locus in familial isolated hyperparathyroidism. *Eur J Endocrinol* 2002;147:313–22.
- Carling T, Correa P, Hessman O, Hedberg J, Skogseid B, Lindberg D, Rastad J, Westin G, Akerstrom G. Parathyroid MEN1 gene mutations in relation to clinical characteristics of nonfamilial primary hyperparathyroidism. *J Clin Endocrinol Metab* 1998;83:2960–3.
- Dwight T, Twigg S, Delbridge L, Wong FK, Farnebo F, Richardson AL, Nelson A, Zedenius J, Philips J, Larsson C, Teh BT, Robinson B. Loss of heterozygosity in sporadic parathyroid tumours: involvement of chromosome 1 and the MEN1 gene locus in 1q13. *Clin Endocrinol (Oxf)* 2000;53:85–92.
- Farnebo F, Teh BT, Kytola S, Svensson A, Phelan C, Sandelin K, Thompson NW, Hoog A, Weber G, Farnebo LO, Larsson C. Alterations of the MEN1 gene in sporadic parathyroid tumours. *J Clin Endocrinol Metab* 1998;83:2627–30.
- Heppner C, Kester MB, Agarwal SK, Debelenko LV, Emmert-Buck MR, Guru SC, Manickam P, Olufemi SE, Skarulis MC, Doppman JL, Alexander RH, Kim YS, Saggart SK, Lubensky IA, Zhuang Z, Liotta LA,

- Chandrasekharappa SC, Collins FS, Spiegel AM, Burns AL, Marx SJ. Somatic mutation of the MEN1 gene in parathyroid tumours. *Nat Genet* 1997;**16**:375–8.
- 17 **Agarwal SK**, Schrock E, Kester MB, Burns AL, Heffess CS, Ried T, Marx SJ. Comparative genomic hybridization analysis of human parathyroid tumours. *Cancer Genet Cytogenet* 1998;**106**:30–6.
 - 18 **Kytola S**, Farnebo F, Obara T, Isola J, Grimelius L, Farnebo LO, Sandelin K, Larsson C. Patterns of chromosomal imbalances in parathyroid carcinomas. *Am J Pathol* 2000;**157**:579–86.
 - 19 **Dotzenrath C**, Teh BT, Farnebo F, Cupisti K, Svensson A, Toell A, Goretzki P, Larsson C. Allelic loss of the retinoblastoma tumour suppressor gene: a marker for aggressive parathyroid tumours? *J Clin Endocrinol Metab* 1996;**81**:3194–6.
 - 20 **Cryns VL**, Thor A, Xu HJ, Hu SX, Wierman ME, Vickery AL, Jr, Benedict WF, Arnold A. Loss of the retinoblastoma tumour-suppressor gene in parathyroid carcinoma. *N Engl J Med* 1994;**330**:757–61.
 - 21 **Farnebo F**, Auer G, Farnebo LO, Teh BT, Twigg S, Aspenblad U, Thompson NW, Grimelius L, Larsson C, Sandelin K. Evaluation of retinoblastoma and Ki-67 immunostaining as diagnostic markers of benign and malignant parathyroid disease. *World J Surg* 1999;**23**:68–74.
 - 22 **Lloyd RV**, Carney JA, Ferreira JA, Jin L, Thompson GB, Van Heerden JA, Grant CS, Wollan PC. Immunohistochemical Analysis of the Cell Cycle-Associated Antigens Ki-67 and Retinoblastoma Protein in Parathyroid Carcinomas and Adenomas. *Endocr Pathol* 1995;**6**:279–87.
 - 23 **Subramaniam P**, Wilkinson S, Shepherd JJ. Inactivation of retinoblastoma gene in malignant parathyroid growths: a candidate genetic trigger? *Aust N Z J Surg* 1995;**65**:714–6.
 - 24 **Sood R**, Bonner TI, Makalowska I, Stephan DA, Robbins CM, Connors TD, Morgenbesser SD, Su K, Faruque MU, Pinkett H, Graham C, Baxevas AD, Klinger KW, Landes GM, Trent JM, Carpten JD. Cloning and characterization of 13 novel transcripts and the human RGS8 gene from the 1q25 region encompassing the hereditary prostate cancer (HPC1) locus. *Genomics* 2001;**73**:211–22.
 - 25 **Carpten JD**, Robbins CM, Villablanca A, Forsberg L, Presciutti S, Bailey-Wilson J, Simonds WF, Gillanders EM, Kennedy AM, Chen JD, Agarwal SK, Sood R, Jones MP, Moses TY, Haven C, Petillo D, Leotlela PD, Harding B, Cameron D, Pannett AA, Hoog A, Heath H 3rd, James-Newton LA, Robinson B, Zarbo RJ, Cavaco BM, Wassif W, Perrier ND, Rosen IB, Kristoffersson U, Turnpenny PD, Farnebo LO, Besser GM, Jackson CE, Morreau H, Trent JM, Thakker RV, Marx SJ, Teh BT, Larsson C, Hobbs MR. HRPT2, encoding parafibromin, is mutated in hyperparathyroidism-jaw tumour syndrome. *Nat Genet* 2002;**32**(4):676–80.
 - 26 **Szabo J**, Heath B, Hill VM, Jackson CE, Zarbo RJ, Mallette LE, Chew SL, Besser GM, Thakker RV, Huff V, et al. Hereditary hyperparathyroidism-jaw tumour syndrome: the endocrine tumour gene HRPT2 maps to chromosome 1q21-q31. *Am J Hum Genet* 1995;**56**:944–50.
 - 27 **Teh BT**, Farnebo F, Kristoffersson U, Sundelin B, Cardinal J, Axelsson R, Yap A, Epstein M, Heath H, Cameron D 3rd, Larsson C. Autosomal dominant primary hyperparathyroidism and jaw tumour syndrome associated with renal hamartomas and cystic kidney disease: linkage to 1q21-q32 and loss of the wild type allele in renal hamartomas. *J Clin Endocrinol Metab* 1996;**81**:4204–11.
 - 28 **Mallette LE**, Malini S, Rappaport MP, Kirkland JL. Familial cystic parathyroid adenomatosis. *Ann Intern Med* 1987;**107**:54–60.
 - 29 **Cavaco BM**, Barros L, Pannett AA, Ruas L, Carvalho M, Ruas MM, Krausz T, Santos MA, Sobrinho LG, Leite V, Thakker RV. The hyperparathyroidism-jaw tumour syndrome in a Portuguese kindred. *QJM* 2001;**94**:213–22.
 - 30 **Haven CJ**, Wong FK, van Dam EW, van der Juijt R, van Asperen C, Jansen J, Rosenberg C, de Wit M, Roijers J, Hoppener J, Lips CJ, Larsson C, Teh BT, Morreau H. A genotypic and histopathological study of a large Dutch kindred with hyperparathyroidism-jaw tumour syndrome. *J Clin Endocrinol Metab* 2000;**85**:1449–54.
 - 31 **Teh BT**, Farnebo F, Twigg S, Hoog A, Kytola S, Korpi-Hyövalti E, Wong FK, Nordenstrom J, Grimelius L, Sandelin K, Robinson B, Farnebo LO, Larsson C. Familial isolated hyperparathyroidism maps to the hyperparathyroidism-jaw tumour locus in 1q21-q32 in a subset of families. *J Clin Endocrinol Metab* 1998;**83**:2114–20.
 - 32 **Villablanca A**, Farnebo F, Teh BT, Farnebo LO, Hoog A, Larsson C. Genetic and clinical characterization of sporadic cystic parathyroid tumours. *Clin Endocrinol (Oxf)* 2002;**56**:261–9.
 - 33 **Farnebo F**, Teh BT, Dotzenrath C, Wassif WS, Svensson A, White I, Betz R, Goretzki P, Sandelin K, Farnebo LO, Larsson C. Differential loss of heterozygosity in familial, sporadic, and uremic hyperparathyroidism. *Hum Genet* 1997;**99**:342–9.
 - 34 **Dwight T**, Kytola S, Teh BT, Theodosopoulos G, Richardson AL, Philips J, Twigg S, Delbridge L, Marsh DJ, Nelson AE, Larsson C, Robinson BG. Genetic analysis of lithium-associated parathyroid tumours. *Eur J Endocrinol* 2002;**146**:619–27.
 - 35 **Solcia E**, Kloppel G, Sobin LH. *World Health Organization. International Histological Classification of Tumours. Histological Typing of Endocrine Tumours*, 2nd ed. Berlin: Springer-Verlag, 2000:48–55.
 - 36 **Kong A**, Gudbjartsson DF, Sainz J, Jonsdottir GM, Gudjonsson SA, Richardsson B, Sigurdardottir S, Barnard J, Hallbeck B, Masson G, Shlien A, Palsson ST, Frigge ML, Thorgeirsson TE, Gulcher JR, Stefansson K. A high-resolution recombination map of the human genome. *Nat Genet* 2002;**31**:241–7.

Screening for genomic rearrangements of the MMR genes must be included in the routine diagnosis of HNPCC

F Di Fiore, F Charbonnier, C Martin, S Frerot, S Olschwang, Q Wang, C Boisson, M-P Buisine, M Nilbert, A Lindblom, T Frebourg

J Med Genet 2004;41:18–20

In hereditary non-polyposis colorectal cancer (HNPCC), the most common form of inherited colorectal cancer, detection of the causal alteration of the mismatch repair (MMR) gene involved is essential for proper management of the families. This will allow the identification of relatives with high risk for colorectal or endometrial cancer, who require the appropriate screening and, conversely, will avert useless surveillance in non-carrier relatives. Mutational studies,¹ based on conventional screening methods, have indicated that point mutations of *MSH2*, *MLH1*, or *MSH6* can be detected in approximately 55% of the families, fulfilling the Amsterdam (AMS) criteria. These stipulate:

- at least three relatives with colorectal cancer, or cancer of the endometrium, small bowel, ureter, or renal pelvis
- one of whom is a first degree relative of the other two
- at least two successive generations affected
- and at least one cancer diagnosed before the age of 50 years.²

In a recent study, we showed that genomic rearrangements of *MSH2* are involved in approximately 20% of the AMS+ HNPCC families without detectable point mutations within *MSH2* or *MLH1*.³ This study was performed using quantitative multiplex PCR of short fluorescent fragments (QMPSF), which can easily detect heterozygous genomic deletions and duplications.^{3–7} This method is based on the simultaneous amplification of short genomic sequences under quantitative

Key points

- In hereditary non-polyposis colorectal cancer (HNPCC), point mutations of *MSH2*, *MLH1*, or *MSH6* are detected in approximately half of the families involved, which therefore fulfil the Amsterdam criteria (AMS).
- We analysed *MSH2* in 120 AMS+ and 212 AMS– HNPCC families without *MSH2* or *MLH1* point mutations, using quantitative multiplex PCR of short fluorescent fragments (QMPSF). We identified in 22% of the AMS+ and in 8% of the AMS– families 19 distinct exonic deletions and two cases of duplication of *MSH2*. We detected seven distinct 5' breakpoints in the deletions removing exon 1. Specific QMPSF analysis of the *MSH2* promoter in 65 AMS+ families, without *MSH2/MLH1* point mutations or *MSH2* exonic deletion, revealed only one case of promoter deletion. Among 86 AMS+ and 106 AMS– families, we detected seven distinct *MLH1* exonic deletions in 7% of the AMS+ families and in 3% of the AMS– families. We found that the selective extinction of MMR protein in the tumours was highly predictive of an MMR rearrangement.
- We conclude that *MSH2* rearrangements are involved in at least 10% of the AMS+ families, which justifies screening for these in the routine diagnosis of HNPCC. The presence of *MLH1* rearrangements should be considered in AMS+ HNPCC patients, when there is a selective loss of *MLH1* expression in the tumours.

Table 1 Frequency of *MSH2* and *MLH1* exonic rearrangements detected by QMPSF in HNPCC families according to their status

Status	<i>MSH2</i>	<i>MLH1</i>
AMS+ families	120*	86†
Without IHC‡ information	101	75
-with a rearrangement	16 (16%)	2 (3%)
With selective extinction of the MMR protein	19	11
-with a rearrangement	11 (58%)	4 (36%)
-AMS+ families with a rearrangement	27 (22%)	6 (7%)
AMS– families	212*	106†
Without IHC‡ information	200	85
-with a rearrangement	9 (4%)	2 (2%)
With selective extinction of the MMR proteins§	12	21
-with a rearrangement	7 (58%)	1 (5%)
-AMS families with a rearrangement	16 (8%)	3 (3%)

*Without *MSH2* or *MLH1* point mutation.

†Without *MSH2* or *MLH1* point mutation, or *MSH2* exonic rearrangement.

‡IHC staining of the tumour.

§In the tumour.

conditions, using dye labelled primers, and the superimposition of the electropherograms of patients and controls.

We have now integrated QMPSF into the routine diagnosis of HNPCC. We first analysed, as previously described,³ the 16 exons of *MSH2* in 332 families, without point mutations within *MSH2* and *MLH1* (table 1). These families corresponded to 120 families fulfilling AMS criteria and 212 AMS– families. Immunohistochemical (IHC) staining of the tumours was performed in 19 AMS+ and 12 AMS– patients, and revealed a selective extinction of the *MSH2* protein. Among the AMS+ patients without IHC information, an *MSH2* genomic rearrangement was detected in 16% of the cases; the detection rate reached 58% in AMS+ patients, showing a selective loss of expression of *MSH2* in their tumours. Among the AMS– patients negative for *MSH2* and *MLH1* mutations, we found an *MSH2* genomic rearrangement in 4% of the cases, when IHC analysis had not been performed, and in 58% of the cases with IHC *MSH2* extinction. QMPSF analysis of the 19 exons of *MLH1* was then carried out in 192 families, corresponding to 86 AMS+

Table 2 Summary of *MSH2* and *MLH1* exonic rearrangements detected by QMPFSF in HNPCC families

<i>MSH2</i>	Families	<i>MLH1</i>	Families
del* exon 1	R1†, R2†, P14	del exons 1–19	U4, U5
del exons 1–2	P15, R12, R13, R20	del exon 4–6	Lu4
del exons 1–4	L7†	del exon 6	S10
del exons 1–6	Li8†, R14, P16, U1, U2, S1, R18	del exons 7–9	R19
del exons 1–7	R9†, Lu1, Lu2, S2, R21	del exons 9–10	P20
del exons 1–8	R10†, R11†, S3, S4	del exon 11	S11, U6
del exons 1–11	U3	del exon 14	S12
del exons 1–15	P12†		
del exon 2	S5		
del exon 3	R3†, P17		
del exons 4–6	Lu3		
del exon 5	P5†		
del exon 5–6	P13†		
del exon 7	L6†		
del exon 7–10	S6		
del exon 8	P18, R15, S7, R16		
del exons 9–10	P19		
del exons 12–13	R17		
del exons 13–15	S8		
Dup† exons 7–8	S9		
Dup exons 9–10	L14†		

*del, deletion.

†Previously published in Charbonnier *et al.*³

‡Dup, duplication.

We have not included in this table two *MLH1* rearrangements removing exon 2 and exons 12–13 that we had initially detected by RT-PCR and documented in Charbonnier, *et al.*⁴

families (including 11 families in which IHC staining of the tumours had been undertaken and revealed a selective extinction of *MLH1*) and 106 AMS– families. Among the AMS+ patients, we found an *MLH1* genomic deletion in 3% of the cases when IHC had not been performed, and in 36% of the cases when IHC had revealed a selective *MLH1* extinction. Although the patients for whom IHC staining of the tumours

revealed a selective loss of expression were low in number, our results indicate that the selective extinction of an *MMR* protein within a tumour is predictive, in HNPCC families, of the genetic alteration.

We detected a total of 21 distinct *MSH2* exonic rearrangements, including 19 deletions and two duplications, in 43 families; and seven exonic deletions of *MLH1* in nine families (table 2). In families with an *MSH2* rearrangement removing exon 1, QMPFSF scanning of 50 Kb of genomic sequences upstream of the *MSH2* transcription initiation site (table 3) revealed at least seven distinct 5' breakpoints. Furthermore, this analysis showed that the recurrent exonic deletions that we detected (deletions of exon 1, exons 1–2, 1–6, 1–7) had been independently generated (tables 2 and 3), thus excluding a founder effect. The numerous breakpoints within the 5' *MSH2* region led us to screen for rearrangements affecting the promoter selectively, which would have escaped the initial QMPFSF analysis of *MSH2* restricted to the 16 exons. We therefore performed a specific QMPFSF assay for the 4.4 kb promoter region, using the promoter amplicons indicated in table 3, and reanalysed 65 AMS+ families without *MSH2*/*MLH1* point mutations or *MSH2* exonic deletion. We identified in a single family a 1.7 kb partial deletion of the promoter removing the -1770–60 region.

This study confirms the following.

- The frequency of *MSH2* exonic rearrangements in AMS+ HNPCC families without detectable point mutations of *MSH2* or *MLH1* can be estimated to approximately 20% (table 1).
- We identified, on the basis of exonic and promoter rearrangements, 30 distinct genomic alterations that demonstrate the remarkable heterogeneity of *MSH2* rearrangements (tables 2 and 3).
- The rearrangements that affect the *MSH2* promoter selectively occur in less than 2% of AMS+ families.
- *MLH1* rearrangements are involved in 7% of AMS+ families without point mutations.

Table 3 QMPFSF scanning of the 5' *MSH2* region in HNPCC families with exonic deletions removing exon 1

Family	Exonic rearrangement	QMPFSF amplicons*								
		–39258 –39107	–34123 –33918	–23323 –23186	–9381 –9243	–5112 –4942	–4244† –4084	–3591† –3387	–1714† –1494	+200‡ 327
R2	del exon 1							+	del*	del
L7	del exons 1–4							+	del	del
R18	del exons 1–6							+	del	del
U1	del exons 1–6							+	del	del
R14	del exons 1–6						+	del	del	del
U2	del exons 1–6						del	del	/	del
R1	del exon 1			+	del	del	del	del	/	del
P15	del exons 1–2			+	del	del	/	del	del	del
R9	del exons 1–7			+	del	del	del	del	del	del
R11	del exons 1–8			+	del	del	del	del	del	del
U3	del exons 1–11			+	del	del	/	del	/	del
R12	del exons 1–2		+	del	del	del	/	del	del	del
R13	del exons 1–2		+	del	/	del	/	del	/	del
S2	del exons 1–7	+	del	del	/	del	/	del	/	del
R10	del exons 1–8	+	del	del	del	del	del	del	del	del
Li8	del exons 1–6	del	del	del	del	del	del	del	del	del
S1	del exons 1–6	del	/	del	/	del	/	del	/	del
Lu1	del exons 1–7	del	/	del	/	del	/	del	/	del
Lu2	del exons 1–7	del	/	del	/	del	/	del	/	del
S3	del exons 1–8	del	/	del	/	del	/	del	/	del
S4	del exons 1–8	del	/	del	/	del	/	del	/	del
P12	del exons 1–15	del	del	del	del	del	del	del	del	del

*Numbered from the *MSH2* transcription initiation site (–68 bp from the ATG) according to the chromosome 2.†These amplicons correspond to the *MSH2* promoter defined by Iwahashi, *et al* (1998).

‡Amplicon corresponding to exon 1.

§Non-deleted.

*Deleted.

Working draft sequence (contig NT_034483). Primers and QMPFSF conditions are available upon request.

Two recent papers,⁸⁻⁹ have reported higher detection rates of *MSH2* and *MLH1* rearrangements, respectively, but these remarkable percentages are probably due to the differences between the populations analysed. The first study,⁸ performed on 24 AMS+ families without point mutations, using Southern Blot analysis, documented the detection of an *MSH2* genomic deletion in 50% of the cases. This remarkable detection rate is probably explained by the fact that the exons 1–6 deletion, detected in seven families, were shown to be associated to a founder effect.⁸ The second study,⁹ based on the QMPFS analysis of 52 AMS+ families without point mutations, reported the detection of genomic deletions of *MSH2* and *MLH1* in 12% of the families, for each gene. In this work, the existence of a common haplotype, in four families harbouring a deletion of *MLH1* exons 1–10, suggested a founder effect and may also have led to an overestimation of the relative contribution of *MLH1* deletions in HNPCC.

In conclusion, we recommend that investigation for *MSH2* rearrangements be included systematically in the routine diagnosis of HNPCC. The contribution of these alterations to HNPCC is higher than that of *MSH6* mutations.¹⁰⁻¹² Considering the lower frequency of *MLH1* rearrangements, except in certain populations where they are associated with a founder effect,⁸⁻⁹ it is probably more efficient to search in HNPCC families only when IHC staining of the tumours has revealed a selective loss of *MLH1* expression.

ACKNOWLEDGEMENTS

We are grateful to Mario Tosi for critical review of the manuscript, and to the clinicians who referred HNPCC families.

Authors' affiliations

F Di Fiore, F Charbonnier, C Martin, S Frerot, T Frebourg, Department of Genetics, University Hospital, and INSERM EMI 9906 -IFRMP, Faculty of Medicine, Rouen, France

S Olschwang, C Boisson, INSERM U434 and Saint Antoine Hospital, Paris, France

Q Wang, Department of Molecular Oncology, Centre Léon Bérard, Lyon, France

M-P Buisine, Laboratory of Biochemistry and Molecular Biology, University Hospital, Lille, France

M Nilbert, Department of Oncology, University Hospital, Lund, Sweden
A Lindblom, Department of Clinical Genetics, Karolinska Hospital, Stockholm, Sweden

This work was supported by L' Association pour la Recherche sur le Cancer, and La Ligue Nationale de Lutte contre le Cancer.

Correspondence to: Dr T Frebourg, Department of Genetics, University Hospital, and INSERM EMI 9906-IFRMP, Faculty of Medicine, 22 Bd. Gambetta, 76183 Rouen, France; Frebourg@chu-rouen.fr

Received 14 July 2003
 Accepted 16 August 2003

REFERENCES

- 1 **Peltomäki P**, Vasen H. The International Collaborative Group on Hereditary Nonpolyposis Colorectal Cancer. Mutations predisposing to hereditary nonpolyposis colorectal cancer: database and results of a collaborative study, *Gastroenterology* 1997;**113**:1146–58.
- 2 **Vasen HF**, Watson P, Mecklin JP, Lynch HT. New clinical criteria for hereditary nonpolyposis colorectal cancer (HNPCC, Lynch syndrome) proposed by the International Collaborative Group on HNPCC. *Gastroenterology* 1999;**116**:1453–6.
- 3 **Charbonnier F**, Olschwang S, Wang Q, Boisson C, Martin C, Buisine MP, Puisieux A, Frebourg T. *MSH2* in contrast to *MLH1* and *MSH6* is frequently inactivated by exonic and promoter rearrangements in hereditary nonpolyposis colorectal cancer. *Cancer Res* 2002;**62**:848–53.
- 4 **Charbonnier F**, Raux G, Wang Q, Drouot N, Cordier F, Limacher JM, Saurin JC, Puisieux A, Olschwang S, Frebourg T. Detection of exon deletions and duplications of the mismatch repair genes in hereditary nonpolyposis colorectal cancer families using multiplex polymerase chain reaction of short fluorescent fragments. *Cancer Res* 2000;**60**:2760–3.
- 5 **Casilli F**, Di Rocco ZC, Gad S, Tournier I, Stoppa-Lyonnet D, Frebourg T, Tosi M. Rapid detection of novel *BRCA1* rearrangements in high-risk breast-ovarian cancer families using multiplex PCR of short fluorescent fragments. *Hum Mut* 2002;**20**:218–26.
- 6 **Bougeard G**, Charbonnier F, Moerman A, Martin C, Ruchoux MM, Drouot N, Frebourg T. Early onset brain tumor and lymphoma in *MSH2* deficient children. *Am J Hum Genet* 2003;**72**:213–16.
- 7 **Bougeard G**, Brugieres L, Chompret A, Gesta P, Charbonnier F, Valent A, Martin C, Raux G, Feunteun J, Bressac-de Paillerets B, Frebourg T. Screening for *TP53* rearrangements in Li-Fraumeni syndrome reveals a complete deletion of the *TP53* gene. *Oncogene* 2003;**22**:840–6.
- 8 **Wagner A**, Barrows A, Wijnen JT, Van Der Klift H, Franken PF, Verkuijlen P, Nakagawa H, Geugien M, Jaghmohan-Changur S, Breukel C, Meijers-Heijboer H, Morreau H, Van Puijenbroek M, Burn J, Coranel S, Kinarski Y, Okimoto R, Watson P, Lynch JF, De La Chapelle A, Lynch HAT, Fodde R. Molecular analysis of hereditary nonpolyposis colorectal cancer in the United States: high mutation detection rate among clinically selected families and characterization of an American founder genomic deletion of the *MSH2* gene. *Am J Hum Genet* 2003;**72**:1088–100.
- 9 **Wang Y**, Friedl W, Lamberti C, Jungck M, Mathiak M, Pagenstecher C, Propping P, Mangold E. Hereditary nonpolyposis colorectal cancer: frequent occurrence of large genomic deletions in *MSH2* and *MLH1* genes. *Int J Cancer* 2003;**103**:636–41.
- 10 **Wijnen J**, de Leeuw W, Vasen H, Van Der Klift H, Moller P, Stormorken A, Meijers-Heijboer H, Lindhout D, Menko F, Vossen S, Moslein G, Tops C, Brocker-Vriends A, Wu Y, Hofstra R, Sijmons R, Cornelisse C, Morreau H, Fodde R. Familial endometrial cancer in female carriers of *MSH6* germline mutations. *Nat Genet* 1999;**23**:142–4.
- 11 **Huang J**, Kuismanen SA, Liu T, Chadwick RB, Johnson CK, Stevens MW, Richards SK, Meek JE, Gao X, Wright FA, Mecklin JP, Jarvinen HJ, Gronberg H, Bisgaard ML, Lindblom A, Peltomäki P. *MSH6* and *MSH3* are rarely involved in genetic predisposition to nonpolyposis colon cancer. *Cancer Res* 2001;**61**:1619–23.
- 12 **Berends MJW**, Wu Y, Sijmons RH, Mensik RGJ, van der Sluis T, Hordijk-Hos JM, de Vries EGE, Hollema H, Karrenbeld A, Buys CHCM, van der Zee AGJ, Hofstra RMW, Kleibeuker JH. Molecular and clinical characteristics of *MSH6* variants: an analysis of 25 index carriers of a germline variant. *Am J Hum Genet* 2002;**70**:26–37.
- 13 **Iwahashi Y**, Ito E, Yanagisawa Y, Akiyama Y, Yuasa Y, Onodera T, Maruyama K. Promoter analysis of the human mismatch repair gene hMSH2. *Gene* 1998;**213**:141–7.

CORRECTION

The authors of the paper by Howell *et al* in the September issue (HRPT2 mutations are associated with malignancy in sporadic parathyroid tumours. *J Med Genet* 2003;**40**:657–63) have notified us of an error. In figure 1, third row from the bottom, for Family ID (Family F1)4†, the histology of the tumour should be Adenoma and not Carcinoma. The authors apologise for the error.