SHORT REPORT

Mutations of *ESPN* cause autosomal recessive deafness and vestibular dysfunction

S Naz, A J Griffith, S Riazuddin, L L Hampton, J F Battey Jr, S N Khan, S Riazuddin, E R Wilcox, T B Friedman

J Med Genet 2004;41:591–595. doi: 10.1136/jmg.2004.018523

We mapped a human deafness locus *DFNB36* to chromosome 1p36.3 in two consanguineous families segregating recessively inherited deafness and vestibular areflexia. This phenotype co-segregates with either of two frameshift mutations, 1988delAGAG and 2469delGTCA, in *ESPN*, which encodes a calcium-insensitive actin-bundling protein called espin. A recessive mutation of *ESPN* is known to cause hearing loss and vestibular dysfunction in the *jerker* mouse. Our results establish espin as an essential protein for hearing and vestibular function in humans. The abnormal vestibular phenotype associated with *ESPN* mutations will be a useful clinical marker for refining the differential diagnosis of non-syndromic deafness.

Most congenital non-syndromic hearing loss is inherited as a recessive Mendelian disorder.¹ Forty recessive deafness loci (*DFNB*) have been mapped and the genes responsible at 20 of these loci have been reported.² With Institutional Review Board (IRB) approval and written informed consent we ascertained families PKSN32 and PKSR5A from Punjab, Pakistan. Detailed clinical histories were obtained for affected individuals (ages ranging from 6 to 40 years), who exhibited prelingual, profound, sensorineural hearing loss and independent ambulation delayed beyond 1.5 years of age. Affected members (ages 14, 16, and 39) of family PKSN32 denied vestibular symptoms although caloric testing with electronystagmography (ENG) revealed vestibular areflexia. The four deaf individuals in family PKSR5A reported balance problems, and caloric-ENG testing confirmed vestibular areflexia. Although congenital, profound hearing loss with vestibular areflexia is accompanied by retinitis pigmentosa (RP) in Usher syndrome type 1,³ affected individuals from PKSN32 or PKSR5A denied nyctalopia or other symptoms of RP. Electroretinography (ERG) on two affected individuals from families PKSN32 (age 14) and PKSR5A (age 19) ruled out the presence of RP.

The deafness segregating in family PKSN32 was not linked to any of the known recessive or dominant deafness loci (Hereditary Hearing Loss Homepage, <http://dnalab-www.uia.ac.be/dnalab/hhh/>). Therefore, a genome-wide search was performed with the Marshfield Weber 9 marker set. We found a 6 cM region of homozygosity on chromosome 1p36.3 among four affected individuals of this family (fig 1). A maximum two-point LOD score of 4.1 ($\theta = 0$) was obtained with marker *DIS214*, defining a new recessive deafness locus, *DFNB36*. In the second family, PKSR5A, the hearing loss was consistent with linkage to *DFNB36* with a maximum two-point LOD score of 3.4 ($\theta = 0$) for marker *DIS3774* (fig 1).

ESPN, a gene in the *DFNB36* critical interval at 1p36.3, was a good positional candidate because a mutation of *Espin* is

known to cause deafness and vestibular dysfunction in the *jerker* mouse.⁴ We screened the coding sequence of *ESPN* (GenBank accession number AL136880) by sequencing PCR amplified *ESPN* exons from genomic DNA of affected individuals in the two families. However, the sequencing analysis of *ESPN* was complicated due to the presence of a second *ESPN*-like sequence on chromosome 1p36.13, annotated as *LOC284729* (GenBank accession number AL035288). As we identified frequent frameshift mutations in *LOC284729* disrupting its open reading frame (data not shown) and due to the absence of sequence encoding homologous residues that are important for actin-bundling activity of espin,⁵ *LOC284729* appears to be an unprocessed pseudogene. Based upon our analyses of *LOC284729*, this putative pseudogene has been assigned the symbol *ESPNP*. Except for the absence of sequences comparable to exon 1 and exon 13, *ESPNP* has all of the other exons and introns of *ESPN* to which it shows 95% nucleotide sequence identity. *ESPNP* is 10 Mb away from *ESPN* in reverse orientation with respect to *ESPN* and is outside the *DFNB36* linkage interval defined by families PKSN32 (fig 1) and PKSR5A (data not shown).

Intronic primers for amplification of exons 1 and 13 were unique for *ESPN* since *ESPNP* lacks homologous sequences, and intronic primers for amplification of exons 2, 6, 7, 8, 9, and 10 were specific for *ESPN* due to multiple mismatches with *ESPNP* sequence (table 1). However, intronic primers for amplification of other exons were either identical to *ESPN* and *ESPNP* (exons 11 and 12) or failed to discriminate between the mismatches of the two sequences (exons 3, 4, and 5). Therefore, we subcloned the PCR products from exons 3, 4, 5, 11, and 12 and sequenced individual clones. The nucleotide mismatches allowed us to unambiguously assign individual clones to either *ESPN* or *ESPNP*.

We detected two different mutations of *ESPN* segregating with the deafness phenotype in families PKSR5A and PKSN32. In family PKSR5A, affected individuals were homozygous for a 4 bp deletion, 1988delAGAG, in exon 9 (fig 2A), while in family PKSN32 all affected individuals were homozygous for a 4 bp deletion, 2469delGTCA, in exon 13 (fig 2B). The obligate carriers in both families were heterozygous and neither mutation was detected among 150 normal-hearing individuals from Pakistan. The 1988delAGAG mutation in exon 9 of *ESPN* was not present in *ESPNP* in deaf individuals of family PKSR5A. *ESPNP* lacks the site of 2469delGTCA mutation in family PKSN32 (fig 2C).

ESPN is predicted to encode an 854 amino acid protein (fig 3) referred to as espin, a name derived from ectoplasmic specialisations (+ in) on the basis of its discovery and localisation in parallel F-actin bundles of ectoplasmic

Abbreviations: ABM, actin-bundling module; ENG, electronystagmography; ERG, electroretinography; IRB, Institutional Review Board; RP, retinitis pigmentosa

Figure 1 Pedigrees and mapping of *DFNB36*. Families PKSN32 and PKSR5A are shown with the haplotypes of chromosome 1p36.3 STR markers. The grey shaded haplotypes represent the ancestral chromosome harbouring the *ESPN* mutation.

specialisation in testes.⁶ Domain prediction algorithms of human espin sequence revealed eight ankyrin-like repeats at the N-terminus, two proline rich regions, a consensus site for ATP or GTP binding (P-loop), which is contained within an actin-binding WH2 motif (amino acids 651–668), and a coiled coil (amino acids 756–831) (figs 2D and 3). There are 66 residues (amino acids 739–804) that show 33% sequence identity to a domain in forked proteins, which are essential for formation of actin bundles in bristles of *Drosophila melanogaster*.⁷ Deletion mutagenesis experiments have

suggested that espin contains three actin-binding sites (figs 2D and 3).^{5, 8} Two of the actin-binding sites at the C-terminus constitute the ABM (figs 2D and 3) and are important for espin activity. *Espin* expression constructs lacking either one or both of these actin-binding sites are unable to cross-link actin filaments when transfected into BHK fibroblasts.⁵

The two mutations of *ESPN*, segregating with hearing loss linked to *DFNB36*, cause frameshifts in the *ESPN* translation reading frame. If translated, 1988delAGAG is predicted to

Table 1 Primers for amplification of *ESPN* exons* and flanking intronic sequences

Oligo	Sequence	Size (bp)	Exon(s) amplified
1F	ATTCGAACCCAGTTTGCTG	1010	1
1R	CCACCCACTTCCAGGACTAC		
2F	AGGAAGGGTGGAGAGATC	859	2
2R	ATGTTGAGTGGGAGCCATT		
3F	GAGGTCAGACACAGCAGGTG	1954	3–5
5R	AGCGTGGGTTCCAGTTATG		
6F	GGAACTGGGTCCTGCTG	552	6
6R	CCTCCCATGTTTAAAGCA		
7F	TACGACTCTGCTCTCCAGCCACT	6909	7 and 8
8R	CCCTCTGCAGCCCTTCTTAAGC		
9F	CCATCAATCTTGGCTTAGG	377	9
9R	ACTGGTGACAGTGCAGGTGA		
10F	CACAGTGTCTCAGGCATCG	444	10
10R	GATGGGCTGTGCATCCAG		
11F	AGCTCTGAGGGGTGTGAC	462	11 and 12
12R	TACAAGGGCCAAGAGACAGG		
13F	CTAGCCCTCTGTCTCAGC	439	13
13R	GGTCAGGGAGGGTTCCAG		

*Sequence corresponds to AL136880.
Nucleotides in each primer mismatched to *ESPN* sequence are underlined. Amplification conditions and sequences of additional primers for mutational analysis of exons 2, 3, 4, 5, 7, and 8 are available as supplemental material (<http://img.bmjournals.com/supplemental/>).

Figure 2 Electropherograms demonstrating *ESPN* mutations and diagrammatic representation of *ESPN* and *ESPNP*. (A) Sequence traces from exon 9 of *ESPN* for control individual VI:3 and affected individual VI:6 in family PKSR5A, showing the wild type sequence and 1988delAGAG, respectively (underlined in the control trace). (B) Sequence traces from exon 13 of *ESPN* for a control and affected individual V:4 in family PKSN32, showing 2469delGTCA (underlined). The traces show the reverse complement strand. (C) Exon/intron structure of *ESPN* (GenBank# AL136880). Exons are represented as numbered boxes, and introns are indicated by lines joining the boxes (not drawn to scale). The locations of mutations found in the sequencing analysis of *ESPN* from the affected individuals in the two families are shown (numbered vertical lines). *ESPNP*, a putative pseudogene, is depicted below *ESPN*. Exons 1 and 13 of *ESPN* are absent and exon 8 is interrupted by an intron. There is only one expressed sequence tag (EST) corresponding to *ESPNP* (GenBank# CB987978), drawn below *ESPNP*. (D) Diagrammatic representation of espin depicting ankyrin repeats encoded by exons 1–4 and the proline rich peptides (PR1 and PR2) encoded by exons 7 and 8. xAB is an actin-binding domain⁵ and is encoded by exon 7 of *ESPN*. The WH2 domain (Wiskott Aldrich syndrome homology region 2) containing the P-loop is encoded by exon 9. The coiled coil and other residues important for actin bundling are formed by amino acids encoded by exons 10–13 constituting the actin-bundling module (ABM).

introduce one substituted amino acid after residue 662, followed by a stop codon at nucleotide 1990 and 2469delGTCA is predicted to introduce 27 substituted amino acids after residue 821, followed by a stop codon at nucleotide 2533 resulting in a truncated 844 amino acid protein with 821 correct and 23 substituted residues. These mutations are presumed to result in loss of function of espin. The resulting mutant proteins are predicted to have either no actin-bundling module, (1988delAGAG), or lack one of the C-terminal actin-binding sites (2469delGTCA), which is necessary for espin activity.⁵ There is evidence for multiple isoforms of Espin in the mouse^{4 5 6 9} and the same may be true in humans as well. However, the mutations associated with *DFNB36* occur in *ESPN* exons, which are known to be present in all reported isoforms of *Espin* in mice.^{4 5 7 9}

PCR analysis of human fetal inner ear cDNA revealed expression of *ESPN* in the inner ear (data not shown). In both the cochlea and the vestibule of the mouse inner ear, espin is

localised mostly to the stereocilia of hair cells.⁴ Stereocilia are specialised microvilli projecting from the apical surfaces of inner ear hair cells and are vital for transduction of sound and for detection of linear and angular acceleration. They contain a densely packed core of parallel bundles of actin filaments.¹⁰ The formation and ordered arrangement of these filaments into bundles requires different actin-binding proteins, which cross-link actin filaments.¹¹ Multiple actin-binding proteins such as fimbrin, plastin, and espin, are expressed in the inner ear.^{11 12} Espin plays a crucial role in cross-linking parallel actin bundles and varying concentrations of espin may determine the extent of elongation and, consequently, the length of parallel actin bundles.¹³

Espin is absent from the stereocilia of *jerker* mice,⁴ and consequently by postnatal day 10 the stereocilia are shortened and have reduced stiffness.¹⁴ Within 3 months of birth, a degenerative process leads to the complete loss of all sensory hair cells in the *jerker* mouse.^{4 14} In addition to espin,

Rat	ESPN	MALEQAMQAARRGDLVLRSLHAAGLLGPSLRDPLDALPVHHAARSGKLHCLRYLVEEVA	60
Mouse	ESPN	MALEQALQAARRGDLVLRSLHAAGLLGPSLRDSDALPVHHAARSGKLHCLRYLVEEVA	60
Human	ESPN	MALEQALQAARQGLDVLRLSLHAAGLLGPSLRDPLDALPVHHAARAGKLHCLRLFVEEAA	60
Rat	ESPN	LPAVSRARNGATPAHDAATGYLSCLQWLLTQGGCRVQEKDNGSATVLHLAARFGHPDVV	120
Mouse	ESPN	LPAVSRARNGATPAHDAATGYLSCLQWLLTQGGCRVQEKDNGSATVLHLAARFGHPDVV	120
Human	ESPN	LPAAARARNGATPAHDAATGHLACLQWLLSQGGCRVQDKDNGSATVLHLAARFGHPEVV	120
Rat	ESPN	NWLLYQGGANSATTTDTGALPIHYAAAKGDLPSMKLLVGHYPEGVNAQTNGATPLYLAC	180
Mouse	ESPN	KWLLYQGGANSATTTDTGALPIHYAAAKGDLPSLKLVLGHYPEGVNAQTNGATPLYLAC	180
Human	ESPN	NWLLHHGGGDPATAATDMGALPIHYAAAKGDFPSLRLLVEHYPEGVNAQTNGATPLYLAC	180
Rat	ESPN	QEGHLEVTKYLVQECSADPHLRAQDGMTPLHAAQMGHNPVLVWLVSFADVSF-EQDHDG	239
Mouse	ESPN	QEGHLEVTKYLVQECSADPHLRAQDGMTPLHAAQMGHNPVLVWLVSFADVSFSEQDHDG	240
Human	ESPN	QEGHLEVTKYLVQECSADPHLRAQDGMTPLHAAQMGHSPVIVWLVSCTDVSLSEQDKDG	240
Rat	ESPN	ATAMHFAASRGHTKVLSWLLHGAETSDQLWGGTPLHDAEENGELECCQILAVNGAGLDV	299
Mouse	ESPN	ATAMHFAASRGHTKVLSWLLHGAETSDQLWGGTPLHDAEENGELECCQILAVNGAGLDV	300
Human	ESPN	ATAMHFAASRGHTKVLSWLLHGGETISADLWGGTPLHDAEENGELECCQILVVGNAELDV	300
Rat	ESPN	RDHDGYTAADLADFNGHTHCSRYLRVTQTLSEHRVLSRDPMSDLEAKQPDGMSNPNTT	359
Mouse	ESPN	RDHDGYTAADLAEFNGHTHCSRYLRVTQTLSEHRVLSRDPQSDMLEAKQLDSGMSNPNTT	360
Human	ESPN	RDRDGYTAADLADFNGHSHCTRYLRVTENLSVEHRVLSRDPSEAELEAKQPDGMSNPNTT	360
Rat	ESPN	MSVQPPNFDLGSPTSTLSNYDSCSSSHSSSKQQRSTR---GARSSDLQS YMDMLNPE---	413
Mouse	ESPN	MSVQPMTFDLGSPTSTLSNYDSCSSSHSSSKQQRNRGIPGARAADLQS YMDMLNPEKSL	420
Human	ESPN	VSVQPLNFDLGSPTSTLSNYDSCSSSHSSIKGQHPPCGLSSARAADIQS YMDMLNPELGL	420
Rat	ESPN	PRSKQKPPSSLLPPPP--PSFPPPPPGTQLPPPPPGYPAPNPVGLHLDNIYMQTKNK	470
Mouse	ESPN	PRGKLGPSPPPPPPPPPSPFPPPPPGTQPPPPPGYPAPNPVGLHLNNIYMQTKNK	480
Human	ESPN	PRGTIGP--TPPPPP--PSFPPPPPGTQLPPPPPGYPAPKPEVGPQAADIYMQTKNK	476
Rat	ESPN	LRHVEVDSLKK-----EPSSGDGYSGLRQDSGLLRQDSELL-----LRHNTGLR	515
Mouse	ESPN	LRHVEVDSLKEPKVELNDQFAQPSGGDGHSLHRQDSGLLRQDSELLHRQELLRHSTGLR	540
Human	ESPN	LRHVETEALKK-----ELSSCDGHD-----GLR	499
Rat	ESPN	RQSDSRKQSFQSKQPSFGDYRQLGRSPGEPLAARPGMAHSEE-----	558
Mouse	ESPN	RQSDSRKQSFQSKQPSFGDYRQMGSRSPGEPLAARPGMAHSEE-----	583
Human	ESPN	RQSDSRKPRAFQSKQPSFGDYRQLGRCPGETLAARPGMAHSEEVRAARQPARAGCPRLGPA	559
Rat	ESPN	-----AALLPGNHVHNGCSADSKASREL--PPPPPPPLPEALSSPPAPPL	603
Mouse	ESPN	-----AALLPGNHVHNGCSADSKASREL--PPPPPPPLPEALSSPPAPPL	628
Human	ESPN	ARGSLEGPSAPPQAALLPGNHVPNGCAADPKASRELPLPPPPPPPLPEAASSPPAPPL	619
Rat	ESPN	PIEGAGAACGQRRSSSSTG-----STKSFNMMSPTGDNSELLAEIKAGKSLKPTPQ	654
Mouse	ESPN	PIEGAGAACGQRRSSSSTGKVRVLRHRKSTKSFNMMSPTGDNSELLAEIKAGKSLKPTPQ	688
Human	ESPN	PIESAGPGCGQRRSSSSTG-----STKSFNMMSPTGDNSELLAEIKAGKSLKPTPQ	670
Rat	ESPN	SKGLTTVFSGSGQPASQPESQPAVSPGSPRARSPTPPASGQPLNGSLVPAPPATLAP	714
Mouse	ESPN	SKGLTTVFSGSGQPASQPESQPLVSPAPSTRPTPPASGQPLNGSVVPAPPATPAP	748
Human	ESPN	SKGLTTVFSGIGQPAFQPDSELPVSPALS PVRSPTPPAAGFQPLNGSLVPVPPTTAP	730
Rat	ESPN	GVHLDVEALIPTLDEQGRPIPEWKQVMVRKLQKMQEEEEQRRKEEEEEEARLASLPAPWR	774
Mouse	ESPN	GVHLDVEALIPTLDEQGRPIPEWKQVMVRKLQKMQEEEEQRRKEEEEEEARLASLPAPWR	808
Human	ESPN	GVQLDVEALIPTHDEQGRPIPEWKQVMVRKMLKMQEEEEQRRKEEEEEEARLASMPAPWR	790
Rat	ESPN	RDILRKKLEEEEREQKRKEEERQKLEETQRAKEQSEKLRTLGYDEAKLAPWQRQVILKKGE	834
Mouse	ESPN	RDILRKKLEEEEREQKRKEEERQKLEETQRAKEQSEKLRTLGYDEAKLAPWQRQVILKKGE	868
Human	ESPN	RDLLRKKLEEEEREQKRKEEERQKQELRREKEQSEKLRTLGYDESKLAPWQRQVILKKGD	850
Rat	ESPN	IPK- 837	
Mouse	ESPN	IPK- 871	
Human	ESPN	IAY- 854	

Figure 3 Espin sequence alignment. ClustalW multiple protein alignment of rat, mouse, and human ESPN. Dark shaded residues denote identical amino acids. Light grey shading represents conserved amino acid substitutions and “-” indicates a gap in the alignment. Bars on top of amino acid residues at the N-terminus indicate the location of ankyrin repeats. Two proline rich regions are boxed. A dashed line indicates actin-binding amino acids (xAB). Asterisks mark the amino acids corresponding to the P-loop. Amino acids forming the WH2 domain are indicated by a bracket. Amino acids constituting the actin-bundling module, ABM,⁵ are underlined. The first amino acid affected by 1988delAGAG or 2469delGTCA is indicated by “Δ”. The amino acids shared by mouse and human espin proteins exhibit 83% identity and 88% similarity. Human and rat ESPN are 86% identical and 90% similar.

there are many other cytoskeletal proteins that are necessary for the development and maintenance of stereocilia. Mutations of *ACTG1* encoding γ -actin were recently reported to cause progressive hearing loss in humans.¹⁵ Both β - and γ -actin are present in the stereocilia of auditory hair cells in chicken,¹⁶ raising the possibility that espin interacts with either β -actin, γ -actin, or both in the stereocilia. Moreover, espin has multiple sites for protein–protein interactions, which may serve as a scaffold for assembly of macromolecular complexes important for structure and function of the stereocilia.

Our findings indicate that espin is essential for both hearing and balance in humans. The association of profound deafness and vestibular dysfunction in the absence of other associated phenotypes is unusual, although vestibular dysfunction is often not carefully evaluated or documented for most non-syndromic recessive deafness loci. Vestibular dysfunction has been excluded for 11 DFNB loci: *B1*, *B6*, *B7/11*, *B12*, *B17*, *B18*, *B21*, *B23*, *B26*, *B29*, and *B30*¹⁷ (see Homepage Hereditary Hearing Loss Homepage for individual references: <http://dnalab-www.uia.ac.be/dnalab/hhh/>). Affected individuals with mutations of *MYO7A* linked to *DFNB2* have a vestibular phenotype¹⁷ comparable to that associated with *DFNB36* and affected individuals with mutations of *MYO15A* also exhibit signs and symptoms of vestibular dysfunction.¹⁷ Vestibular dysfunction was also reported in a few deaf individuals who have recessive mutations of *MYO6*.¹⁸ The abnormal vestibular phenotype associated with *ESPN* mutations is unusual and will be a useful clinical marker for refining the differential diagnosis of non-syndromic deafness.

ACKNOWLEDGEMENTS

We are grateful to the families for their participation and the staff of Children's Hospital, Lahore, Pakistan, for their cooperation and help in providing facilities for ENG examinations. RNA for cDNA synthesis was received from C C Morton, who obtained human fetal ear tissue under IRB approval by the Human Research Committee of Brigham and Women's Hospital. We thank James Bartles, Inna Belyantseva, Erich Boger, Rafael Caruso, Dennis Drayna, Robert Morell, Barbara Ploplis, and Susan Sullivan for their comments and expertise.

Authors' affiliations

S Naz, S Riazuddin, E R Wilcox, T B Friedman, Section on Human Genetics, LMG, NIDCD, NIH, Rockville, MD 20850, USA
A J Griffith, Section on Gene Structure and Function, LMG, NIDCD, NIH, Rockville, MD, USA
A J Griffith, Hearing Section, NIDCD, NIH, Rockville, MD, USA
L L Hampton, J F Battey Jr, Section on G-protein Coupled Receptors, NINDS, NIH, Bethesda, MD, USA
S N Khan, S Riazuddin, National Center of Excellence in Molecular Biology, University of the Punjab, Lahore, Pakistan

Part of the study carried out in Pakistan was supported by a grant from the Higher Education Commission in Islamabad. Intramural funds from the National Institute on Deafness and Other Communication Disorders

(Z01 DC000035-07, Z01 DC000064-03, Z01 DC000039-07) supported this work.

Conflict of interest: none declared.

GenBank accession numbers: mouse *Espn* AF239886, rat *Espn* NM_019622.1, human *ESPN* AL136880, *ESPN* (LOC284729) AL035288; *ESPN* genomic sequence AL031848 and AL158217; genomic sequence for *ESPN* AL021920; EST for *ESPN* CB987978.

Correspondence to: Dr T B Friedman, Laboratory of Molecular Genetics, NIDCD, NIH, Rockville, MD 20850, USA; friedman@nidcd.nih.gov

Revised version received 8 March 2004

Accepted for publication 10 March 2004

REFERENCES

- Morton NE. Genetic epidemiology of hearing impairment. *Ann NY Acad Sci* 1991;**630**:16–31.
- Friedman TB, Griffith AJ. Human nonsyndromic sensorineural deafness. *Annu Rev Genomics Hum Genet* 2003;**4**:341–402.
- Smith RJ, Berlin CI, Hejtmancik JF, Keats BJ, Kimberling WJ, Lewis RA, Moller CG, Peltas MZ, Tranebjaerg L. Clinical diagnosis of the Usher syndromes. *Am J Med Genet* 1994;**50**:32–8.
- Zheng L, Sekerkova G, Vranich K, Tilney LG, Mugnaini E, Bartles JR. The deaf jerker mouse has a mutation in the gene encoding the espin actin-bundling proteins of hair cell stereocilia and lacks espins. *Cell* 2000;**102**:377–85.
- Bartles JR, Zheng L, Li A, Wierda A, Chen B. Small espin: a third actin-bundling protein and potential forked protein ortholog in brush border microvilli. *J Cell Biol* 1998;**143**:107–19.
- Bartles JR, Wierda A, Zheng L. Identification and characterization of espin, an actin-binding protein localized to the F-actin-rich junctional plaques of Sertoli cell ectoplasmic specializations. *J Cell Sci* 1996;**109**:1229–39.
- Petersen NS, Lankenau DH, Mitchell HK, Young P, Corcos VG. Forked proteins are components of fiber bundles present in developing bristles of *Drosophila melanogaster*. *Genetics* 1994;**136**:173–82.
- Chen B, Li A, Wang D, Wang M, Zheng L, Bartles JR. Espin contains an additional actin-binding site in its N terminus and is a major actin-bundling protein of the Sertoli cell-spermatid ectoplasmic specialization junctional plaque. *Mol Biol Cell* 1999;**10**:4327–39.
- Sekerkova G, Loomis PA, Changyaleket B, Zheng L, Eytan R, Chen B, Mugnaini E, Bartles JR. Novel espin actin-bundling proteins are localized to Purkinje cell dendritic spines and bind the Src homology 3 adapter protein insulin receptor substrate p53. *J Neurosci* 2003;**23**:1310–19.
- Tilney LG, Tilney MS, DeRosier DJ. Actin filaments, stereocilia, and hair cells: how cells count and measure. *Annu Rev Cell Biol* 1992;**8**:257–74.
- Bartles JR. Parallel actin bundles and their multiple actin-bundling proteins. *Curr Opin Cell Biol* 2000;**12**:72–8.
- Daudet N, Lebart MC. Transient expression of the t-isoform of plastins/fimbrin in the stereocilia of developing auditory hair cells. *Cell Motil Cytoskeleton* 2002;**53**:326–36.
- Loomis PA, Zheng L, Sekerkova G, Changyaleket B, Mugnaini E, Bartles JR. Espin cross-links cause the elongation of microvillus-type parallel actin bundles in vivo. *J Cell Biol* 2003;**163**:1045–55.
- Sjostrom B, Anniko M. Genetically induced inner ear degeneration. A structural and functional study. *Acta Otolaryngol Suppl* 1992;**493**:141–46.
- Zhu M, Yang T, Wei S, DeWan AT, Morell RJ, Ellenbein JL, Fisher RA, Leal SM, Smith RJ, Friderici KH. Mutations in the gamma-actin gene (*ACTG1*) are associated with dominant progressive deafness (DFNA20/26). *Am J Hum Genet* 2003;**73**:1082–91.
- Hofer D, Ness W, Drenckhahn D. Sorting of actin isoforms in chicken auditory hair cells. *J Cell Sci* 1997;**110**:765–70.
- Cremers CWRJ, Smith RJH, eds. Genetic hearing impairment, its clinical presentations. *Adv Otorhinolaryngol* 2002;**61**:113–56.
- Ahmed ZM, Morell RJ, Riazuddin S, Gropman A, Shaukat S, Ahmad MM, Mohiddin SA, Fananapazir L, Caruso RC, Husnain T, Khan SN, Griffith AJ, Riazuddin S, Friedman TB, Wilcox ER. Mutations of *MYO6* are associated with recessive deafness, DFNB37. *Am J Hum Genet* 2003;**72**:1315–22.