ELECTRONIC LETTER

LKB1 exonic and whole gene deletions are a common cause of Peutz-Jeghers syndrome

E Volikos,* J Robinson,* K Aittomäki, J-P Mecklin, H Järvinen, A M Westerman, F W M de Rooij, T Vogel, G Moeslein, V Launonen, I P M Tomlinson, A R J Silver, L A Aaltonen

This article is available free on JMG online via the **JMG Unlocked** open access trial, funded by the Joint Information Systems Committee. For further information, see <http://jmg.bmjjournals.com/cgi/content/full/42/2/97>

J Med Genet 2006;**43**:e18 (<http://www.jmedgenet.com/cgi/content/full/43/5/e18>). doi: 10.1136/jmg.2005.039875

Background: *LKB1/STK11* germline mutations cause Peutz-Jeghers syndrome (PJS). The existence of a second *PJS* locus is controversial, the evidence in its favour being families unlinked to *LKB1* and the low frequency of *LKB1* mutations found using conventional methods in several studies. Exonic and whole gene deletion or duplication events cannot be detected by routine mutation screening methods.

Objective: To seek evidence for *LKB1* germline deletions or duplications by screening patients meeting clinical criteria for *PJS* but without detected mutations on conventional screening.

Methods: From an original cohort of 76 patients, 48 were found to have a germline mutation by direct sequencing; the remaining 28 were examined using multiplex ligation dependent probe amplification (MLPA) analysis to detect *LKB1* copy number changes.

Results: Deletions were found in 11 of the 28 patients (39%)—that is, 14% of all *PJS* patients (11/76). Five patients had whole gene deletions, two had the promoter and exon 1 deleted, and in one patient exon 8 was deleted. Other deletions events involved: loss of exons 2–10; deletion of the promoter and exons 1–3; and loss of part of the promoter. No duplications were detected. Nine samples with deletions were sequenced at reported single nucleotide polymorphisms to exclude heterozygosity; homozygosity was found in all cases. No MLPA copy number changes were detected in 22 healthy individuals.

Conclusions: These results lessen the possibility of a second *PJS* locus, as the detection rate of germline mutations in *PJS* patients was about 80% (59/76). It is suggested that MLPA, or a suitable alternative, should be used for routine genetic testing of *PJS* patients in clinical practice.

Peutz-Jeghers syndrome (PJS) is diagnosed by the presence of multiple hamartomatous polyps of characteristic morphology, usually accompanied by mucocutaneous “freckling”. *PJS* patients often present in their first decade of life with intussusception, obstruction, or volvulus caused by small bowel polyps. The main complication of *PJS* later in life is a greatly increased risk of colorectal and upper gastrointestinal cancers, and of tumours at multiple other sites. *PJS* is caused by germline mutations in the *LKB1/STK11* gene.¹ *LKB1* acts as a tumour suppressor in humans, and loss of function may cause multiple defects, including those of cell polarity,² activation of the hypoxia pathway through AMP kinase and hypoxia inducible factor 1 α ,^{3,4} and increased Wnt signalling.⁵

The original report of *LKB1* mutations in *PJS* demonstrated mutations in about 90% of cases (11/12) using direct sequencing of constitutional DNA and mRNA.¹ Most subsequent screens for mutations in *PJS* have produced a similar, although slightly lower, frequency of patients with *LKB1* mutations, although the methods used for mutation detection have varied. Two studies, however, found mutations in only 10%⁶ and 18%⁷ of *PJS* cases, although the proportion of familial cases in the latter study was low. These findings have led to claims that a single *PJS* locus cannot explain all cases of *PJS*. Support for this contention has been provided by linkage studies in *PJS* families,^{8–10} which have reported exclusion of the *LKB1* locus (chromosome 19p13.3). Countervailing arguments have been that many tumour susceptibility genes harbour a large proportion of mutations which cannot be detected by routine screening methods, and that several conditions—even including normal variants—may mimic the pigmentation of *PJS*, leading to diagnostic misclassification.

Whole gene and exonic deletions and duplications cannot be detected using routine mutation screening methods. Given the high frequency of such changes in diseases such as von Hippel-Lindau syndrome,¹¹ and the increasing realisation that conditions such as hereditary non-polyposis colon cancer (HNPCC) are often caused by such changes,^{12–14} we screened for gene and exon scale mutations in a set of *PJS* patients without detected *LKB1* mutations. The results have implications for diagnostic genetic testing in *PJS* patients and for the hypothesised second *PJS* locus.

METHODS

A diagnosis of *PJS* was made when the patient presented with two or more *PJS* polyps, or one polyp and typical pigmented lesions, or one polyp and a family history of *PJS*.¹⁵ A cohort of 76 patients meeting these criteria had been screened previously for germline *LKB1* mutations using direct sequencing of each exon plus intron–exon boundaries. In 48 patients, small scale mutations had been found, including protein truncating changes, missense variants targeting phosphorylation sites or other highly conserved residues, and splice site variants.^{16–19} In all, 28 patients without detected mutations (13 familial, 13 isolated cases, two unknown) were identified for the present study. The multiplex ligation dependent probe amplification (MLPA) *LKB1/STK11* kit, designed to search for *LKB1* deletions or duplications at exon or whole gene level, was obtained from MRC-Holland (Amsterdam, Netherlands). MLPA reactions were

Abbreviations: MLPA, multiplex ligation dependent probe amplification; *PJS*, Peutz-Jeghers syndrome; SNP, single nucleotide polymorphism

Table 1 MLPA results for patients with <i>LKB1</i> changes		
Patient ID(s)	Result	Family history
11200	Whole gene deletion	Sporadic
11199	Whole gene deletion	Sporadic
11668	Whole gene deletion	Sporadic
24/1/1	Whole gene deletion	Unknown
02-551	Whole gene deletion	Familial
000/1/1; 52/1/1	Promoter and exon 1 deletion	Familial
59/1/1; 11181	Promoter and exon 1 deletion	Familial
13/1/1	Exon 8 deletion	Sporadic
7282	Exon 2–10 deletion	Familial
AXY	Promoter deletion	Sporadic
LEH	Promoter and exon 1–3 deletion	Familial

carried out according to the manufacturer’s instructions and the products were analysed using the Applied Biosystems PRISM™ 3100 DNA sequencer. MLPA data were analysed using the Applied Biosystems Genotyper software (PE Applied Biosystems, Foster City, California, USA). Finally, the MLPA data were evaluated by a Microsoft Excel based program originally designed to detect *APC* copy number changes (www.ngsl.org.uk/ManchesterPages/Downloads/Dosage) and adapted to evaluate copy number changes with the MLPA *STK11* test kit. Samples were analysed alongside 22 normal healthy individuals to exclude detection of false positives. All samples with possible MLPA changes were tested at least twice to confirm the change. Single nucleotide polymorphism (SNP) analysis was undertaken by sequencing to exclude heterozygosity within deletions reported by MLPA.

RESULTS AND COMMENT

Eleven of the 28 PJS patients tested (39%) had germline *LKB1* mutations detected by MLPA (table 1). Five patients had whole gene deletions, two had the promoter region and exon 1 deleted, and in one patient exon 8 was deleted. The remaining detected deletions, each in a single patient, involved the following: loss of exons 2–10; deletion of the promoter region and exons 1–3; and loss of part of the promoter region. No duplications of any part of the gene were detected. Twenty two healthy individuals were screened by MLPA and no copy number changes detected.

For nine samples with deletions (75%), the appropriate region of the gene was sequenced in the patients concerned, so as to confirm the MLPA data by excluding heterozygosity at all reported SNPs. Two SNPs (rs3764640, rs3764641) lying near to exon 1 of *LKB1* were found to be homozygous in the three patients with the promoter region and exon 1 deletions. Five SNPs (rs3795063, rs2075604, rs2075606, rs741764, and rs2075608), spread through *LKB1* from intron one to intron eight, were homozygous in the five patients with loss of the entire copy of *LKB1*. Absence of polymorphic sites underlying the MLPA detection oligonucleotides was additionally confirmed using the latest version of dbSNP (BUILD 125, 29 September 2005). Given the original sample of 76 patients, we estimate the proportion of PJS cases with large deletions at 14% (11/76). These patients had no clinicopathological features which distinguished them from the other patients with or without *LKB1* mutations in terms of family history, number of polyps, presence of pigmentation, or development of cancer (details not shown).

LKB1 is a highly conserved protein, especially within the kinase domain (codons 48–309), and deletion of the promoter region or single or multiple exons are all predicted to have profound effects on kinase activity. Exon 8 encodes amino acids 308–369, distal to the kinase domain, but functional effects of such C-terminal mutations have been

shown using in vitro assays. Sapkota *et al*²⁰ found that phosphorylation of residue 431 was essential for cell growth suppression by *LKB1* and Forcet *et al*²¹ showed that C-terminal *LKB1* mutations reduced activation of AMPK and diminished the ability of *LKB1* to induce and maintain cell polarity.

Large scale germline deletions have very rarely been reported in PJS, probably because methods suitable for their detection have not been used. Jiang *et al*⁶ found one PJS patient with a deletion spanning exons 2–7 of *LKB1*. One whole gene deletion and one genomic rearrangement (a ~2 kb deletion) has been reported in two PJS patients.^{19 22} Another study⁷ used long range polymerase chain reaction to amplify from exon 3 to exon 8 of *LKB1* in order to search for products of novel sizes, but found no changes in their PJS patients. However, this method would not have detected any of the cases in our study with whole gene deletions or deletion of the promoter region and exon 1.

The finding of relatively frequent, large scale germline *LKB1* mutations in PJS patients strongly suggests that MLPA testing (or some suitable alternative) should be introduced into the diagnostic genetics laboratory. Our findings also have consequences for the likelihood that there exists a second uncharacterised PJS locus. Analysis of published *LKB1* mutations shows that a mean of 59% of PJS cases (median = 58%, range = 10%–100%) have harboured a germline mutation, which was detectable using standard DNA based screening methods such as single stranded conformational polymorphism (SSCP) analysis, conformation specific gel electrophoresis (CSGE), dHPLC, and direct sequencing of coding regions and intron–exon boundaries.^{1 6 7 9 10 18 23–30} In addition, we estimate that approximately 10% of cases appear to have mutations which are detectable using RNA based methods only.^{1 23 30} Therefore, allowing for a mutation detection rate of about 40% by MLPA in the remaining 30% of PJS cases, the frequency of PJS patients with detectable mutations becomes about 80–85%. Given that the mutation detection rate in familial PJS cases has been consistently higher in families than in isolated cases, it is likely that a proportion of the latter have only had a presumptive or possible diagnosis of PJS.

In our view, therefore, the proportion of unambiguous PJS cases with *LKB1* mutations is sufficiently high for there to be no good case for a second PJS locus based on these data alone. During the preparation of this paper, another study on germline *LKB1* deletions³¹ reached identical conclusions. Seventeen of 34 PJS patients without small scale *LKB1* mutations were found to have germline deletions by MLPA. The authors found that when only patients who met the clinical criteria for PJS were considered, the overall mutation detection rate was 94% (64% point mutations and 30% large deletions), very close to our estimate from this study. We therefore conclude that the remaining evidence for the existence of an uncharacterised minor PJS locus therefore comes solely from families whose disease is unlinked to *LKB1*.

ACKNOWLEDGEMENTS

The study was supported by grants from the Academy of Finland (44870/Finnish Centre of Excellence Programme 2000–2005, 76227, 77547), the Sigrid Juselius Foundation, the Cancer Society of Finland, and Helsinki University Central Hospital. The support of Cancer Research UK is also acknowledged

Authors’ affiliations

E Volikos, J Robinson, A Silver, Cancer Research UK Colorectal Cancer Unit, St Mark’s Hospital, Harrow, Middlesex, UK
K Aittomäki, Clinical Genetics, Haartman Institute, University of Helsinki, Helsinki, Finland

J-P Mecklin, Department of Surgery, Jyväskylä Central Hospital, Jyväskylä, Finland
H Järvinen, Laboratory of Cancer Genetics, Institute of Medical Technology, University of Tampere and Tampere University Hospital, Tampere, Finland
A M Westerman, **F W M de Rooij**, Laboratory of Vascular and Metabolic Diseases, Department of Internal Medicine, Erasmus Medical Centre, Rotterdam, Netherlands
T Vogel, Unfallchirurgie, Universitätsklinikum, Heinrich Heine Universität, Düsseldorf, Germany
V Launonen, **L Aaltonen**, Department of Medical Genetics, Haartman Institute, University of Helsinki, Helsinki, Finland
G Moeslein, Allgemein-und Viszeralchirurgie, Universitätsklinikum, Heinrich Heine Universität, Düsseldorf, Germany
I Tomlinson, Molecular and Population Genetics Laboratory, London Research Institute, Cancer Research UK, London, UK, and Cancer Research UK Colorectal Cancer Unit, St Mark's Hospital, Harrow, Middlesex, UK

Conflicts of interest: none declared

*EV and JR contributed equally to this work.

Correspondence to: Ian Tomlinson, Molecular and Population Genetics Laboratory, London Research Institute, Cancer Research UK, London WC2A 3PX, UK; ian.tomlinson@cancer.org.uk

Received 25 November 2005

Revised version received 13 January 2006

Accepted for publication 16 January 2006

REFERENCES

- Hemminki A, Markie D, Tomlinson I, Avizienyte E, Roth S, Loukola A, Bignell G, Warren W, Aminoff M, Hoglund P, Jarvinen H, Kristo P, Pelin K, Ridanpaa M, Salovaara R, Toro T, Bodmer W, Olschwang S, Olsen AS, Stratton MR, de la Chapelle A, Aaltonen LA. A serine/threonine kinase gene defective in Peutz-Jeghers syndrome. *Nature* 1998;**391**:184-7.
- Marignani PA. LKB1, the multitasking tumour suppressor kinase. *J Clin Pathol* 2005;**58**:15-19.
- Brugarolas J, Kaelin WG. Dysregulation of HIF and VEGF is a unifying feature of the familial hamartoma syndromes. *Cancer Cell* 2004;**6**:7-10.
- Hardie DG. New roles for the LKB1→AMPK pathway. *Curr Opin Cell Biol* 2005;**17**:167-73.
- Lin-Marq N, Borel C, Antonarakis SE. Peutz-Jeghers LKB1 mutants fail to activate GSK-3beta, preventing it from inhibiting Wnt signaling. *Mol Genet Genomics* 2005;**273**:184-96.
- Jiang CY, Esufali S, Berk T, Gallinger S, Cohen Z, Tobi M, Redston M, Bapat B. STK11/LKB1 germline mutations are not identified in most Peutz-Jeghers syndrome patients. *Clin Genet* 1999;**56**:136-41.
- Boardman LA, Couch FJ, Burgart LJ, Schwartz D, Berry R, McDonnell SK, Schaid DJ, Hartmann LC, Schroeder JJ, Stratakis CA, Thibodeau SN. Genetic heterogeneity in Peutz-Jeghers syndrome. *Hum Mutat* 2000;**16**:23-30.
- Buchet-Poyau K, Mehenni H, Radhakrishna U, Antonarakis SE. Search for the second Peutz-Jeghers syndrome locus: exclusion of the STK13, PRKCG, KLK10, and PSCD2 genes on chromosome 19 and the STK11IP gene on chromosome 2. *Cytogenet Genome Res* 2002;**97**:171-8.
- Mehenni H, Gehrig C, Nezu J, Oku A, Shimane M, Rossier C, Guex N, Blouin JL, Scott HS, Antonarakis SE. Loss of LKB1 kinase activity in Peutz-Jeghers syndrome, and evidence for allelic and locus heterogeneity. *Am J Hum Genet* 1998;**63**:1641-50.
- Olschwang S, Boisson C, Thomas G. Peutz-Jeghers families unlinked to STK11/LKB1 gene mutations are highly predisposed to primitive biliary adenocarcinoma. *J Med Genet* 2001;**38**:356-60.
- Richards FM, Crossey PA, Phipps ME, Foster K, Latif F, Evans G, Sampson J, Lerman MI, Zbar B, Affara NA, et al. Detailed mapping of germline deletions of the von Hippel-Lindau disease tumour suppressor gene. *Hum Mol Genet* 1994;**3**:595-8.
- Wijnen J, van der Klift H, Vasen H, Khan PM, Menko F, Tops C, Meijers Heijboer H, Lindhout D, Moller P, Fodde R. MSH2 genomic deletions are a frequent cause of HNPCC. *Nat Genet* 1998;**20**:326-8.
- Taylor CF, Charlton RS, Burn J, Sheridan E, Taylor GR. Genomic deletions in MSH2 or MLH1 are a frequent cause of hereditary non-polyposis colorectal cancer: identification of novel and recurrent deletions by MLPA. *Hum Mutat* 2003;**22**:428-33.
- Wang Y, Friedl W, Lamberti C, Jungck M, Mathiak M, Pagenstecher C, Propping P, Mangold E. Hereditary nonpolyposis colorectal cancer: frequent occurrence of large genomic deletions in MSH2 and MLH1 genes. *Int J Cancer* 2003;**103**:636-41.
- Tomlinson IP, Houlston RS. Peutz-Jeghers syndrome. *J Med Genet* 1997;**34**:1007-11.
- Lim W, Hearle N, Shah B, Murday V, Hodgson SV, Lucassen A, Eccles D, Talbot I, Neale K, Lim AG, O'Donohue J, Donaldson A, Macdonald RC, Young ID, Robinson MH, Lee PW, Stoodley BJ, Tomlinson I, Alderson D, Holbrook AG, Vyas S, Swarbrick ET, Lewis AA, Phillips RK, Houlston RS. Further observations on LKB1/STK11 status and cancer risk in Peutz-Jeghers syndrome. *Br J Cancer* 2003;**89**:308-13.
- Sweet K, Willis J, Zhou XP, Gallione C, Sawada T, Alhopuro P, Khoo SK, Patocs A, Martin C, Bridgeman S, Heinz J, Pilarski R, Lehtonen R, Prior TW, Frebourg T, Teh BT, Marchuk DA, Aaltonen LA, Eng C. Molecular classification of patients with unexplained hamartomatous and hyperplastic polyposis. *JAMA* 2005;**294**:2465-73.
- Wang ZJ, Churchman M, Avizienyte E, McKeown C, Davies S, Evans DG, Ferguson A, Ellis I, Xu WH, Yan ZY, Aaltonen LA, Tomlinson IP. Germline mutations of the LKB1 (STK11) gene in Peutz-Jeghers patients. *J Med Genet* 1999;**36**:365-8.
- Ylikorkala A, Avizienyte E, Tomlinson IP, Tiiainen M, Roth S, Loukola A, Hemminki A, Johansson M, Sistonen P, Markie D, Neale K, Phillips R, Zaubner P, Twama T, Sampson J, Jarvinen H, Makela TP, Aaltonen LA. Mutations and impaired function of LKB1 in familial and non-familial Peutz-Jeghers syndrome and a sporadic testicular cancer. *Hum Mol Genet* 1999;**8**:45-51.
- Sapkota GP, Kieloch A, Lizcano JM, Lain S, Arthur JS, Williams MR, Morrice N, Deak M, Alessi DR. Phosphorylation of the protein kinase mutated in Peutz-Jeghers cancer syndrome, LKB1/STK11, at Ser431 by p90(RSK) and cAMP-dependent protein kinase, but not its farnesylation at Cys(433), is essential for LKB1 to suppress cell growth. *J Biol Chem* 2001;**276**:19469-82.
- Forcet C, Etienne-Manneville S, Gaude H, Fournier L, Debilly S, Salmi M, Baas A, Olschwang S, Clevers H, Billaud M. Functional analysis of Peutz-Jeghers mutations reveals that the LKB1 C-terminal region exerts a crucial role in regulating both the AMPK pathway and the cell polarity. *Hum Mol Genet* 2005;**14**:1283-92.
- Le Meur N, Martin C, Saugier-Verber P, Joly G, Lemoine F, Moirot H, Rossi A, Bachy B, Cabot A, Joly P, Frebourg T. Complete germline deletion of the STK11 gene in a family with Peutz-Jeghers syndrome. *Eur J Hum Genet* 2004;**12**:415-18.
- Abad AA, Gunther K, Kraus C, Hohenberger W, Ballhausen WG. Mutation screening at the RNA level of the STK11/LKB1 gene in Peutz-Jeghers syndrome reveals complex splicing abnormalities and a novel mRNA isoform (STK11 c.597(insertion mark)598insIVS4). *Hum Mutat* 2001;**18**:397-410.
- Kruse R, Uhlhaas S, Lamberti C, Keller KM, Jackisch C, Steinhard J, Knopfle G, Loff S, Back W, Stolte M, Jungck M, Propping P, Friedl W, Jenne DE. Peutz-Jeghers syndrome: four novel inactivating germline mutations in the STK11 gene. *Mutations in brief*, No 227. Online. *Hum Mutat* 1999;**13**:257-8.
- Nakagawa H, Koyama K, Miyoshi Y, Ando H, Baba S, Watanabe M, Yasutomi M, Matsuura N, Monden M, Nakamura Y. Nine novel germline mutations of STK11 in ten families with Peutz-Jeghers syndrome. *Hum Genet* 1998;**103**:168-72.
- Resta N, Simone C, Marenzi C, Montera M, Gentile M, Susca F, Gristina R, Pozzi S, Bertario L, Bufo P, Carlomagno N, Ingrosso M, Rossini FP, Tenconi R, Guanti G. STK11 mutations in Peutz-Jeghers syndrome and sporadic colon cancer. *Cancer Res* 1998;**58**:4799-801.
- Resta N, Stella A, Susca FC, Di Giacomo M, Forleo G, Miccalis I, Rossini FP, Genuardi M, Piepoli A, Grammatico P, Guanti G. Two novel mutations and a new STK11/LKB1 gene isoform in Peutz-Jeghers patients. *Hum Mutat* 2002;**20**:78-9.
- Scott RJ, Crooks R, Meldrum CJ, Thomas L, Smith CJ, Mowat D, McPhillips M, Spigelman AD. Mutation analysis of the STK11/LKB1 gene and clinical characteristics of an Australian series of Peutz-Jeghers syndrome patients. *Clin Genet* 2002;**62**:282-7.
- Westerman AM, Entius MM, Boor PP, Koole R, de Baar E, Offerhaus GJ, Lubinski J, Lindhout D, Halley DJ, de Rooij FW, Wilson JH. Novel mutations in the LKB1/STK11 gene in Dutch Peutz-Jeghers families. *Hum Mutat* 1999;**13**:476-81.
- Yoon KA, Ku JL, Choi HS, Heo SC, Jeong SY, Park YJ, Kim NK, Kim JC, Jung PM, Park JG. Germline mutations of the STK11 gene in Korean Peutz-Jeghers syndrome patients. *Br J Cancer* 2000;**82**:1403-6.
- Aretz S, Stienen D, Uhlhaas S, Loff S, Back W, Pagenstecher C, McLeod DR, Graham GE, Mangold E, Santer R, Propping P, Friedl W. High proportion of large genomic STK11 deletions in Peutz-Jeghers syndrome. *Hum Mutat* 2005;**26**:513-19.