Inheritance of autoinflammatory diseases: shifting paradigms and nomenclature

Isabelle Touitou

Correspondence to

Professor Isabelle Touitou,
CHRU Montpellier, UM1,
INSERMU844, UMAI,
laboratoire de génétique,
Hopital A de Villeneuve,
34295, Montpellier Cedex 15,
France;
isabelle.touitou@inserm.fr

Received 31 January 2013

Revised 28 February 2013

Accepted 1 March 2013

Published Online First

27 March 2013

ABSTRACT

Over 15 years have passed since the discovery of the first autoinflammatory gene, *MEFV*, responsible for familial Mediterranean fever. The identification of another gene, *TNFRSF1A*, in 1999 led to the concept of autoinflammation which characterises rheumatological conditions triggered by a defective innate immunity. Substantive progress has been made since then with the identification of 18 autoinflammatory genes accounting for up to 24 disease entities showing overlapping symptoms. The accumulation of studies reporting patients with missing or excess mutations as compared with expected numbers favours the hypothesis that these diseases are distributed along a continuum ranging from monogenic to multifactorial conditions, rather than featuring only classical modes of inheritance. Moreover, the probable interactions of environmental and epigenetic factors further obscure our understanding of the mechanisms underlying the phenotypic expression of patients. This review explores the history of autoinflammatory gene discovery, discusses the nosological disparities stemming from the clinical versus pathophysiological definition of autoinflammatory diseases and summarises various inheritance patterns. This review calls for a consistent disease nomenclature and presents a reconciling hypothesis which places different sequence variants within the autoinflammatory disease continuum. Integrating these new concepts should help to facilitate communication between health professionals and promote personalised patient care.

INTRODUCTION

The name autoinflammation was proposed in 1999 to echo that of autoimmunity.^{1 2} However, the rather artificial boundaries between autoinflammatory diseases (innate immunity) and autoimmune diseases (adaptive immunity) are gradually disappearing to give rise to the currently preferred concept of immunological disease continuum.³ Behçet's disease is a good example of a multifactorial condition with features of both autoinflammation and autoimmunity.⁴ 'Pure' autoinflammatory diseases are classically defined as conditions without high titre auto-antibodies or antigen-specific T cells. While initially focused on hereditary recurrent fevers (HRFs), autoinflammatory diseases now encompass a wide spectrum of disorders ranging from rare monogenic, for example, the prototype familial Mediterranean fever (FMF), to more frequent multifactorial diseases, for example, Crohn's disease. This explains why the team which originally proposed the term 'autoinflammatory disease' now suggests that its definition should be reassessed to include all clinical disorders marked by abnormally increased inflammation, mediated predominantly by

the innate immune system, with a significant host predisposition.⁵ Indeed, most cases of autoinflammatory disease are characterised by attacks of systemic inflammation resulting in recurrent fever, acute arthritis and increased acute phase reactants. Frequent accompanying symptoms are skin lesions or rashes, myalgia, bone deformation or sensorineural defects. The key symptoms of autoinflammatory diseases are listed in table 1. The appropriate diagnosis of a disease can prove challenging due to overlapping or mimicking symptoms, or even association with, autoimmune diseases. Although several classifications of autoinflammatory diseases have been proposed in order to help the clinician, not one of these has been consensually adopted (box 1).

Our understanding of the pathophysiology of autoinflammatory diseases has dramatically increased due to the unravelling of defective signalling pathways. The discovery of the FMF gene in 1997 shed light on a new death domain (the pyrin domain⁶) that presents high homology with proteins involved in inflammation and apoptosis.^{7 8} A second gene containing a death domain was found in 1999 to be responsible for a dominant form of recurrent fever: Tumour Necrosis Factor (TNF) Receptor Associated Periodic Syndrome or TRAPS.² Since then, 16 other autoinflammatory genes have been discovered (table 1), including more than 1000 sequence variants.^{9–11} Their clinical significance is often unclear, and the mode of transmission of these diseases is becoming difficult to interpret as the number of families with unexpected genetic associations or patterns of inheritance is growing.

This review has compiled comprehensive genetic data on autoinflammatory diseases in order to clarify the current complexity and controversy surrounding their inheritance, and calls for a consistent disease nomenclature. A list of current disease and gene acronyms is provided in box 2.

LESSONS FROM HRFs

HRFs are the prototype of autoinflammatory diseases as they include the first disease whose gene was identified, that is, FMF.^{7 8} While HRFs are classically monogenic families with unexpected genetic patterns have been reported (box 3) and are discussed below. Large series of patients with recurrent fevers and biological inflammation, in which other common inflammatory diseases have been ruled out, are now available enabling the dissection of genetic and non-genetic factors affecting the patient phenotype.

Missing mutations

This chapter includes the discovery of either no mutations in the suspected gene or of a single

Editor's choice
Scan to access more
free content

To cite: Touitou I. *J Med Genet* 2013;50:349–359.

Table 1 Genes responsible for monogenic autoinflammatory diseases and their corresponding mode of transmission, disease names and key symptoms

Gene name	Transmission	Disease names			Key symptoms
		Historical	Clinical	Physiological	
<i>CARD14</i> *	Dominant		PRP		Follicular papules, scaly red-orange patches, palmoplantar hyperkeratosis
<i>IL10</i>	Recessive		IBD		Early-onset severe enterocolitis
<i>IL10RA</i>	Recessive		IBD		Early-onset severe enterocolitis
<i>IL10RB</i>	Recessive		IBD		Early-onset severe enterocolitis
<i>IL1RN</i>	Recessive			DIRA	Sterile multifocal osteomyelitis, periostitis, pustular psoriasis
<i>IL36RN</i>	Recessive			DITRA	Psoriasis-associated pustular phenotypes
<i>LPIN2</i>	Recessive	Majeeds syndrome	CRMO (subtype of)		Multifocal osteomyelitis, dyserythropoietic anaemia, neutrophilic dermatosis
<i>MEFV</i>	Recessive	Mamou and Cattani	FMF (periodic fever)		Recurrent fever, peritonitis, arthralgia, erythema
<i>MVK</i>	Dominant†		FMF		Recurrent fever, peritonitis, arthralgia, erythema
	Recessive	Dutch-type fever	HIDS	MKD, mild	Recurrent fever, vomiting, diarrhoea, arthralgias, skin signs, lymphadenopathy
	Recessive		MEVA	MKD, severe	Mental and growth retardation, ataxia
	Dominant		DSAP		Keratoses
<i>NLRP12</i>	Dominant	Guadeloupean fever	FCAS2	NAPS12	Cold-induced urticaria, recurrent fever, arthralgia
<i>NLRP3</i>	Dominant		FCAS	CAPS (cryopyrinopathy)	Cold-induced urticaria, recurrent fever, arthralgia
	Dominant	MWS		CAPS (cryopyrinopathy)	Urticaria, sensorineural impairment
	Dominant		CINCA/NOMID	CAPS (cryopyrinopathy)	Urticaria, neonatal meningitis
<i>NOD2</i>	Dominant	Blau (Jabs) syndrome			Granulomatous arthritis, uveitis, camptodactyly
<i>PLCγ2</i> ‡	Dominant			APLAID	Recurrent blistering skin lesions, arthralgia, enterocolitis, mild immunodeficiency
<i>PSMB8</i>	Recessive	NNS	JMP, CANDLE		Recurrent fever, lipodystrophy
<i>PSTPIP1</i>	Dominant		PAPA		Pyoderma gangrenosum, acne, pyogenic arthritis
<i>RBCK1</i> §	Recessive				Chronic autoinflammation, bacterial infections, muscular amylopectinosis
<i>SH3BP2</i>	Dominant		Cherubism		Fibro-osseous lesions
<i>TNFRSF1A</i>	Dominant	Hibernian fever		TRAPS	Recurrent fever, abdominal pain, arthralgia, myalgia, skin lesions

Genes responsible for the four first described hereditary recurrent fevers are underlined.

*The other *CARD14* associated phenotype (PSORS2) was not included because it is currently considered that although innate immunity through NFKB activation contributes to the onset of psoriasis, the ensuing chronic inflammation is mostly driven by T cells.

†Not consensual.

‡The phenotypically distinct PLAID (PLCG2-associated Antibody deficiency and Immune Dysregulation)/FCAS3 syndrome associated with this gene does not include autoinflammatory features.

§No disease name is yet associated with mutations in this gene.

CINCA, cutaneous and articular syndrome; NOMID, neonatal-onset multisystem inflammatory disease.

For further acronyms see box 2.

mutation in recessive diseases. Such patients are hereafter referred to as 'genetically orphan'. Several scientific and medical reasons could account for the 20%–85% of genetically orphan patients, including the large number of autoinflammatory genes presumed to be unidentified, the lack of functional tests (except for mevalonate kinase deficiency (MKD)) and the rather unspecific main clinical feature (recurrent fever at the paediatric age) which characterises most autoinflammatory patients.^{12–19} However, this significant proportion could also be due to the evolution in how many of the autoinflammatory diseases are defined, which remains a matter of debate and depends on the habits and discipline of health professionals.

Familial Mediterranean fever

FMF was known as periodic disease or recurrent polyserositis, prior to being called 'Familial Mediterranean fever' by Heller in 1955,²⁰ a name now universally used. All of these names clearly refer to the key symptoms of the disease. However, fever and

polyserositis are also common to the other HRFs. Livneh and others defined the clinical criteria for FMF.^{21–24} Although very useful to the clinicians, they could not be totally specific. As numerous patients with FMF clinical criteria but without *MEFV* mutations were reported, genetic heterogeneity was proposed.²⁵ However, since then a *MEFV2* gene has not been identified (figure 1), and the proportion of genetically orphan FMF patients depends on patient ancestry. It is low in Mediterranean countries (20%–25%), although very high in other populations (80%–85%).^{16–26} This high rate may also be due to wrong clinical overdiagnosis. FMF is classically recessively transmitted, yet cases of dominant fever were reported which justified the creation of a new entity named 'dominant periodic fever' as no genetic test was available until 1997. After this milestone, some families were indeed related to the *MEFV* gene since a dominant segregation of a heterozygous *MEFV* mutation was proven, and series of FMF patients with a single *MEFV* mutation were described.^{27–33} After an unsuccessful search for the second

Box 1 Evolving classification of autoinflammatory diseases

- ▶ The first classifications of autoinflammatory diseases were based on the patients' phenotype. Hence, hereditary recurrent fevers including the prototype familial Mediterranean fever define monogenic disorders with recurrent fever as the key symptom.
- ▶ More recently, McGonagle and colleagues³ suggested an integrated classification using the autoimmune versus autoinflammatory ratio involved in the condition as a means to rank each autoinflammatory condition within the continuum of immunological diseases.
- ▶ At the same time, the NIH group suggested a classification scheme based on the molecular insights gathered over the past decade.⁸⁸
- ▶ No authoritative classification has emerged in 2012, most likely due to the extending identification of phenotype overlaps and genetic heterogeneity.

mutation, it was speculated that a true dominant transmission of certain severe mutations could exist. A recent study derived from the largest European cohort, Eurofever, a shared online registry for patients with autoinflammatory diseases,^{34–35} supported gene dosage.³⁶ Gershoni and colleagues suggested that complex alleles can lead to a more severe form of the disease.^{37–38} Among other hypotheses are as yet unknown molecular mechanisms, such as regulation of transcription,³⁹ non-exonic mutations^{40–41} or, more likely, mutations in another gene.^{25–42–45}

TNF Receptor Associated Periodic Syndrome

In 1968, there were reports of a Swedish family with recurrent fever, abdominal pain that persisted for 7–14 days and renal deposition of amyloid identical to that seen in FMF.⁴⁶ In 1982, another family with an autosomal dominant periodic fever was described in Ireland, and the term 'Hibernian fever' was created on this occasion.⁴⁷ It was revealed that these families and others identified later^{48–49} suffered from genetic conditions other than FMF. Upon the 1999 discovery of the causative gene, *TNFRSF1A*, McDermott *et al*² preferred to associate the phenotype with the mutated protein, and created the TRAPS acronym that stands for 'TNF Receptor Associated Periodic Syndrome'. 'Hibernian' or 'dominant periodic' fevers are no longer in use. As a *TNFRSF1A* mutation was not identified in a proportion of patients with the TRAPS-like phenotype, Borghini *et al*⁴⁴ analysed the sequence of genes known to play a pivotal role in the mechanisms downregulating the TNF-induced inflammatory response such as receptor shedding, receptor internalisation and negative regulation of nuclear factor- κ B transcription. This work revealed only a few unreported variants, apparently neutral; therefore, the authors concluded that the gene(s) responsible for TRAPS-like syndrome still remain to be identified.

Cryopyrin Associated Periodic Syndrome

A similar logic was applied to name the three clinical entities caused by dominant mutations within the same gene, *NLRP3*, encoding cryopyrin. 'Cryopyrin Associated Periodic Syndrome' (CAPS⁵⁰ or cryopyrinopathies⁵¹) was generated as a unifying term for Familial Cold Autoinflammatory Syndrome (FCAS), Muckle-Wells Syndrome and Chronic Infantile Neurological Cutaneous and Articular syndrome, also known as

Box 2 Acronyms in autoinflammatory diseases

- ▶ APLAID, Autoinflammatory PLCG2-associated Antibody deficiency and Immune Dysregulation
- ▶ CANDLE, Chronic Atypical Neutrophilic Dermatitis with Lipodystrophy and Elevated temperature syndrome
- ▶ CAPS, Cryopyrin Associated Periodic Syndrome
- ▶ *CARD14*, Caspase Recruitment Domain Protein 14
- ▶ DIRA, Deficiency of Interleukin 1 Receptor antagonist
- ▶ DITRA, Deficiency of Interleukin 36 Receptor antagonist
- ▶ DSAP, Disseminated Superficial Actinic Porokeratosis
- ▶ FCAS, Familial Cold Urticaria
- ▶ FMF, Familial Mediterranean Fever
- ▶ HIDS, Hyper IgD Syndrome
- ▶ *HOIL-1*, Haeme-Oxidised IRP2 ubiquitin Ligase 1
- ▶ HRF, Hereditary Recurrent Fever
- ▶ IBD, Inflammatory Bowel Disease
- ▶ IL10, Interleukin 10
- ▶ IL10RA, Interleukin 10 Receptor α
- ▶ IL10RB, Interleukin 10 Receptor β
- ▶ IL1RN, Interleukin 1 Receptor Antagonist
- ▶ IL36RN, Interleukin 36 Receptor Antagonist
- ▶ JMP, Joint Contractures, Muscular Atrophy, Microcytic Anaemia and Panniculitis-Induced Lipodystrophy
- ▶ LPIN2, Lipin2
- ▶ MEVA, Mevalonic Aciduria
- ▶ *MEFV*, MEditerranean FeVer
- ▶ MKD, Mevalonate Kinase Deficiency
- ▶ *MVK*, MeValonate Kinase
- ▶ NAPS12, NLRP12-Associated Periodic Syndrome
- ▶ *NLRP12*, NOD-Like-Receptor Family, Pyrin Domain-Containing 12
- ▶ *NLRP3*, NOD-Like-Receptor Family, Pyrin Domain-Containing 3
- ▶ NNS, Nakajo-Nishimura Syndrome
- ▶ NOD2, Nucleotide-binding Oligomerisation Domain Protein 2
- ▶ PAPA, Pyogenic sterile Arthritis, Pyoderma gangrenosum, and Acne
- ▶ PLAID, PLCG2-associated Antibody deficiency and Immune Dysregulation
- ▶ *PLCG2*, Phospholipase C, γ -2
- ▶ PRP, Pityriasis Rubra Pilaris
- ▶ *PSMB8*, Proteasome Subunit, β -Type, 8
- ▶ PSORS, Psoriasis
- ▶ *PSTPIP1*, Proline/Serine/Threonine Phosphatase-Interacting Protein 1
- ▶ *RBCK1*, Ranbp-Type and C3HC4-Type Zinc Finger-Containing 1
- ▶ *SH3BP2*, SH3 Domain-Binding Protein 2
- ▶ *TNFRSF1A*, Tumour Necrosis Factor Receptor Super Family 1A
- ▶ TRAPS, TNF Receptor Associated Periodic Syndrome
- ▶ VUS, Variant of uncertain significance

Neonatal-Onset Multisystem Inflammatory Disease. These conditions are no longer considered as three subphenotypes with increasing severity because of the heterogeneous presentation of this monogenic disease. Instead, the indepth examination of patients with *NLRP3* mutations demonstrated that they display a continuum of symptoms.⁵¹ Consistently, although some mutations were associated with only one of the three CAPS diseases,

Box 3 Genetic mechanisms involved in the inheritance of hereditary recurrent fevers

- ▶ Complex allele: the combination of two or more mutations occurring in cis within the same gene
- ▶ Gene dosage: effect on the phenotype is proportional to the number of mutated alleles
- ▶ Genetic heterogeneity: mutation(s) in one of alternative genes with a strong effect causing the disease
- ▶ Genetic homogeneity: mutation(s) in a single known gene with a strong effect causing the disease
- ▶ Modifier genes: genes responsible for variable expression of the disease
- ▶ Oligogenism: mutation(s) in a limited number of genes with an intermediate effect causing the disease (two genes acting together cause digenism)
- ▶ Pseudodominance: vertical segregation of recessive diseases due to abnormally high mutation frequency usually related to consanguinity or founder effect
- ▶ Susceptibility genetic factors: several genes with a mild effect act together to cause the disease

for example, L305P in FCAS, several sequence variants did not clearly correlate with the phenotype.⁵¹ The absence of clear dominant segregation was explained in some families by the existence of *de novo* mutations⁴⁵ (also found in TRAPS⁵²) and mosaicism.^{53–55} It was also revealed that some of the orphan CAPS patients had mutations in other genes, such as *NLRP12*⁵⁶ or *IL1RN*.⁵⁷

Mevalonate kinase deficiency

'Hyperimmunoglobulinemia D and periodic fever Syndrome' (HIDS, formerly Periodic Fever, Dutch Type⁵⁸) owes its name to increased IgD titres reported in patients suffering from recurrent fever, macula-papular rashes and arthralgia lasting for 3–7 days. Active gene hunting unexpectedly revealed that this recessive condition was caused by mutations in the mevalonate kinase gene⁵⁹ responsible for severe mevalonic aciduria. The switch to the pathophysiological name, MKD, is progressing slowly.⁵⁸ All autoinflammatory experts agree that the term 'HIDS' should be dropped since high IgD sera levels are not specific and can be found in other inflammatory conditions^{60–62} or can be absent in MKD.^{59 61 63 64} MKD is classically recessively inherited and true orphan cases either do not exist or are scarce, since the rate of genetic confirmation (homozygosity or compound heterozygosity) nearly reaches 100% in patients with decreased mevalonate enzymatic activity.⁶⁵ Unexpectedly, dominant mutations in this same gene were recently associated with 'disseminate superficial actinic porokeratosis' (DSAP).⁶⁶ Although DSAP patients had no clinical features of MKD, or abnormalities in their serum IgD concentrations, it remains unclear whether they had decreased enzymatic activity. However, the mutations c.417_418insC and c.604G>A found in two DSAP patients were also reported in two different MKD patients in a compound heterozygous state.

Excess mutations

More than two mutations in recessive diseases

Consanguinity, as well as founder effect, is a well-recognised cause of pseudodominance in recessive diseases, as they increase

the frequency of rare pathogenic alleles. It is not unusual that three mutations segregate in a single family suffering from FMF.⁶⁷ One such case was also identified in a MKD family.⁶⁸ Four mutations were found in Druze FMF patients.⁶⁹

Two mutations in dominant diseases

A handful of reports describe patients bearing two mutations in either the gene *TNFRSF1A* or *NLRP3*, responsible for dominant HRFs.^{70–73} In most cases, one of the two sequence variants is of unclear significance (VUS), such as p.Arg121Gln (usual name R92Q, c.362G>A,) or p.Val198Met (V198M, c.592G>A), in these two genes respectively. Three R92Q homozygotes were identified (ref⁷⁰ and personal communication), and one p.Pro75Leu (usual name P46L, c.224C>T) homozygote (personal communication). Other combinations include V198M and p.Leu353Pro (L353P, c.1058T>C) within *NLRP3* in a patient with a FCAS phenotype,⁷³ and p.Cys102Trp (usual name C73W, c.306C>G) and R92Q within *TNFRSF1A* in a patient with a TRAPS phenotype (personal communication). It is difficult to determine whether there is a gene dose response as first, the variants are usually not phased and may well be located on the same chromosome (complex allele), and second, the reported cases are too scarce to allow statistical evaluation.

Mutations in more than one gene

Cases in which several mutated genes cluster are increasingly reported (figure 2), confusing the clinician while sometimes regrettably delaying accurate and relevant patient care. Up to four mutations have been found. The most likely cause of association is a fortuitous coexistence of highly prevalent mutations in a single individual. Indeed, in most patients one of the two mutations was common in the general population. Digenism was also investigated;^{29 72} however, this was only confirmed in one case through clearly demonstrating cosegregation of the R92Q and V198M variants with the phenotype in a multiplex family suffering from a dominant form of autoinflammatory disease with features of TRAPS, CAPS and Behçet's disease.⁷²

Modifying factors

Genetic factors

Gender and certain mutations are known to confer a higher risk of renal amyloidosis to patients. p.Met694Val (M694V, c.2080A>G) homozygosity has been repeatedly associated with severe disease and kidney complications in FMF patients.²⁶ In addition, cysteine mutations in the TRAPS gene are thought to lead to renal amyloidosis, although large series are lacking.⁷⁴ Polymorphisms in the major histocompatibility complex class I chain-related gene A (MICA)^{75 76} and the serum amyloid-A genes^{75 77–79} have both been reported as bad prognostic variables in FMF.

Environmental factors

Environment has been suspected as a modifying factor since the initial study of Armenian FMF patients living in California.⁸⁰ The authors found that migrants had less amyloidosis than the other patients. In a very large collaborative study,⁸¹ we assessed several factors that may play a role in renal amyloidosis, and showed that the country of recruitment, rather than *MEFV* genotype, was the key risk factor. This risk paralleled infant mortality rates, a highly sensitive indicator of population health which reflects economic development, general living conditions, social well-being and quality of environment.⁸¹ Ozen *et al*⁸² showed that 78% of patients living in Turkey had a severe course of FMF compared with 34% living in Germany. Based

Figure 1 Clinical versus pathophysiological definition of autoinflammatory diseases; discussion on the example of familial Mediterranean fever (FMF). (A) Clinical definition: FMF currently has a clinical name. As it is a recessive disease, two mutations in the causative gene *MEFV* are expected. Provided that the gene has been totally explored and rare mutations excluded, which is never routinely carried out, there are two hypotheses accounting for those patients with nil or only one mutation: existence of another gene (*MEFV2*) or reassignment of the patients to another hereditary recurrent fever (HRF). Other genetic, environmental or epigenetic factors may modify the phenotype although are not mandatory for the disease to occur. Multifactorial diseases mimicking FMF are not discussed. (B) Pathophysiological definition: A naming proposition matching other HRFs. Pyrin Associated Periodic Syndrome would cover all patients with two non-allelic *MEFV* mutations. Phenotype 3 defines asymptomatic individuals.¹¹⁴ This figure is only reproduced in colour in the online version.

on the concordance in clinical manifestations in Israeli monozygote and dizygote twins, Ben-Zvi *et al*⁸³ calculated that the environmental effect on the phenotype of FMF was estimated at around 12%.

Epigenetic factors

Very little is known about the involvement of epigenetics in autoinflammatory disorders. In the sole available study, the methylation level of *MEFV* exon 2 within tended to be slightly higher in FMF patients compared with healthy controls.⁸⁴ This needs further confirmation in larger patient series and in other populations.

FUTURE DIRECTIONS

How to deal with new autoinflammatory diseases?

Issue: phenotype overlaps, name complexity

The tremendous insights gained into the pathophysiology of autoinflammatory diseases fuelled candidate gene screening and exome sequencing in genetically orphan patients. This research confirmed that a single gene could be involved in various phenotypes and, conversely, many different genes could be involved in overlapping phenotypes (table 1).

Three disease names were developed for a group of phenotypes caused by mutations in the Proteasome Subunit, β -Type 8 (*PMSB8*) gene. These names reflect the main clinical features, that is, 'joint contractures, muscle atrophy, microcytic anaemia and panniculitis-induced lipodystrophy syndrome',⁸⁵ or 'chronic atypical neutrophilic dermatosis with lipodystrophy and elevated temperature syndrome',⁸⁶ or the name of the physicians who described the disease at that time, 'Nakajo-Nishimura syndrome'.⁸⁷

Families with cold-induced urticaria and fever yet no *NLRP3* mutations (orphan CAPS) were investigated. Two of them were

related to mutations in *NLRP12*. This condition has no consensual name. It was still unnamed upon the gene discovery,⁵⁶ and was agreed as *NLRP12* Associated Periodic Syndrome to parallel TRAPS and CAPS in a database dedicated to autoinflammatory mutations, Infervers (<http://fmf.igh.cnrs.fr/ISSAID/infervers/>).⁹ Later, it was labelled FCAS2 or Guadeloupe variant periodic fever in a review by Masters and coworkers,⁸⁸ as well as *NLRP12* associated disorders by the gene discoverer in 2011,⁸⁹ and alternatively FCAS2, *NLRP12* associated disease and *NLRP12*-related periodic fever in two recent papers by the Eurofever consortium.^{35 90}

The pathophysiological rule was adopted to name two recently recognised recessive autoinflammatory conditions with severe skin lesions, deficiency of interleukin 1 receptor antagonist (*DIRA*)⁵⁷ and deficiency of interleukin 36 receptor antagonist (*DITRA*).⁹¹ However, a naming issue could arise if mutations in another member of the interleukin receptor antagonist family also happened to be responsible for an autoinflammatory disease. The *DIRA* gene was identified after a candidate approach in three families negative for *NLRP3* mutations, but fully responsive to anti-interleukin 1 therapies, while *DITRA* was discovered after exome sequencing of consanguineous Tunisian families with generalised pustular psoriasis. A recent paper showed that the *DITRA* spectrum also covers palmar-plantar pustulosis and acrodermatitis continua of Hallopeau.⁹²

The psoriasis susceptibility locus 2 (*PSORS2*) gene was revealed to be responsible for a second monogenic form of psoriasis caused by mutations in the dominant gene, caspase recruitment domain family, member 14 (*CARD14*).^{93 94} *CARD14* is also responsible for familial pityriasis rubra pilaris.⁹⁵

Autoinflammatory phospholipase C, γ 2 (*PLC γ 2*) associated antibody deficiency and immune dysregulation (*APLAID*) is caused by dominant mutations in the *PLC γ 2* gene.⁹⁶ This

Figure 2 Dual mutations in hereditary recurrent fever (HRF) genes. A colour code is used to represent the four main HRF genes: Blue: *MEFV*; Green: *TNFRSF1A*; Red: *MVK*; Orange: *NLRP3*. Darker backgrounds indicate patients with confirmed genetic diagnosis: two clearly pathogenic mutations for recessive diseases (familial Mediterranean fever (FMF) and mevalonate kinase deficiency (MKD)), and one clearly pathogenic mutation for dominant diseases (TNF Receptor Associated Periodic Syndrome and Cryopyrin Associated Periodic Syndrome). The number of patients recorded with a given genotype is proportional to the surface of the corresponding circle. Overlaps depict the coexisting variants reported in the two genes. The most frequent combinations tend to include variants of unclear significance such as E148Q (*MEFV*) and R92Q (*TNFRSF1A*) as defined in the guidelines for the genetic diagnosis of HRFs.¹¹⁵ M694V and p.Val377Ile (V377I, c.1129G>A) are the two most frequent causative mutations in FMF and MKD, respectively. This figure is only reproduced in colour in the online version.

syndrome is the autoinflammatory form of PLC γ 2 associated antibody deficiency and immune dysregulation (PLAID, also called FCAS3).⁹⁷

The latest hereditary autoinflammatory syndrome which has been linked to a gene has not been named.^{98–99} The patients with this unnamed syndrome presented with chronic inflammation, bacterial infections and muscular amylopecinosis. They were found to carry biallelic loss-of-expression and loss-of-function mutations in *RBCK1* (*HOIL1*), a component of the linear ubiquitination chain assembly complex.

Solution: disease nomenclature consistency

Ben-Chetrit and colleagues suggested the adoption of the general name 'pyrin associated disorders' for FMF.¹⁰⁰ An approach consistent with TRAPS and CAPS would call for 'PAPS', standing for Pyrin Associated Periodic Syndrome (figure 1). There are pros and cons in generalising the nomenclature in this way.

The advantages are that all patients diagnosed with protein associated disease names would de facto be genetically confirmed. The notion of genetic heterogeneity would no longer have justification. Any existing clinical criteria would be supportive, but dispensable when unavailable. The best example is TRAPS, the HRF with the largest clinical spectrum for which there have been no attempts to delineate clinical criteria. Interestingly, this rule unifies the genetic aspect and catalyses the development of targeted and therefore more efficient therapeutic molecules.

Drawbacks of using this paradigm include the mathematically higher proportion of genetically orphan patients with a convincing autoinflammatory phenotype, coined with the inelegant yet tempting diagnosis of 'autoinflammatory disease-like patients'. More of an issue is the fact that the definition of 'mutation' is becoming vague for sequence variants that were thought to be causative upon the gene discovery, yet are now believed to be at

best functional polymorphisms. Classical examples are p.Glu148Gln (E148Q, c.442G>C) in the *MEFV* gene and R92Q in the *TNFRSF1A* gene. Guidelines have been assembled to help genetic interpretation of these variants. The group of patients with a 'low penetrance' genotype would also increase, especially if mutational screening of asymptomatic or atypical relatives is boosted with the development of high throughput genetic techniques.

A consensus conference proved both successful and helpful for the nomenclature of proteins involved in innate immunity regulation.¹⁰¹ For example, the key inflammasome protein was given the alternative names of cryopyrin, CIAS1, NALP3, PYPAF1 and CTRL1.1 before it was definitively fixed as NLRP3. Such a meeting would be highly recommended to harmonise the names of current and future autoinflammatory monogenic conditions.

How to deal with VUS?

Issue: monogenic versus multifactorial diseases

An increasing number of studies support the involvement of sequence variants of genes responsible for Mendelian autoinflammatory diseases in either autoinflammatory or autoimmune complex conditions (figure 3). Crohn's disease is one of the first multifactorial diseases in which an innate immunity gene named *NOD2/CARD15*,¹⁰² responsible for a dominant autoinflammatory disorder named Blau,¹⁰³ was identified as a major inflammatory bowel disease (IBD1) gene. The FMF, TRAPS and CAPS genes were also inconsistently associated with Crohn's disease.^{104–107} *MEFV* and *TNFRSF1A* may have a role in Behçet's disease.^{43–108} *MEFV*, *NLRP3* and *TNFRSF1A* were suggested as susceptibility factors for multiple sclerosis. The variant R92Q in the latter gene was the most convincing as it was confirmed in multiple genome-wide association studies.¹⁰⁹ Nevertheless, this mutation did not appear to be a severity marker of the disease neither modifying the clinical progression

Figure 3 Association of genes responsible for monogenic autoinflammatory diseases with multifactorial diseases. To obtain the most relevant figures and avoid duplicated data, studies were only retained which described significantly large series. Reviews and case reports were not included. Studies consistently demonstrating significant association of the gene with the disease are depicted in red; studies showing a trend or contradictory results are shown in purple; blue is used for absence of association. AS, ankylosing spondylitis; BD, Behçet's disease; CAD, coronary artery disease; CD, Crohn's disease; CeliacD, celiac disease; FMS, fibromyalgia syndrome; HSP, Henoch-Schönlein purpura and polyarteritis nodosa; IBD, inflammatory bowel disease; JIA, juvenile idiopathic arthritis; MI, myocardial infarction; MS, multiple sclerosis; PBC, primary biliary cirrhosis; PFAPA, periodic fever with adenitis pharyngitis and aphthosis; PR, palindromic arthritis; PsA, psoriasis; PsoJIA, psoriatic JIA; RA, rheumatoid arthritis; SLE, systemic lupus erythematosus; UC, ulcerative colitis. This figure is only reproduced in colour in the online version.

of multiple sclerosis nor its therapeutic response. Although the NLRP3 inflammasome has a critical role in monosodium urate crystal-induced inflammatory gout attacks,¹¹⁰ no study demonstrated *NLRP3* variants associated with gout. All in all, this review revealed numerous conflicting results (figure 3), probably due to different study designs and populations.

Solution: integrating patient background

The coexistence of complex inflammatory diseases and monogenic autoinflammatory gene variants common in the general population is likely to be fortuitous. An appealing hypothesis is that 'severe mutations' (eg, M694V in *MEFV*) could segregate with specific monogenic disorders, while other 'mild/low penetrance mutations' (eg, R92Q within *TNFRSF1A*) could favour non-specific inflammation (figure 4). It is tempting to speculate that the accumulation of such VUS associated with a non-genetic background could either trigger multifactorial diseases or at least modify their expression or severity. Depending on circumstances, a sequence variant could behave as a true causative

mutation, a functional polymorphism or remain silent. This hypothesis reconciles the current controversies surrounding the clinical significance of several sequence variants such as E148Q (*MEFV*) or Q703K (*NLRP3*), and accounts for why they may be found at the same frequency in both patients and the general population.

Current genetic diagnostic strategies which are restricted to the screening of a limited number of genes or exons will not definitively resolve the issue of dealing with VUS. The hypothesis proposed will undoubtedly be confirmed with the development of exome sequencing approaches which will provide phenotype associated genetic signatures.

CONCLUSIONS

Increasing knowledge of the inheritance of autoinflammatory diseases switched simple paradigms into puzzling genetic situations. FMF, first described as a recessive disease due to a handful of clear causative mutations, was revealed to be possibly dominantly inherited and its gene to harbour a constellation of

Figure 4 Inheritance of autoinflammatory diseases, a hypothesis bridging monogenic and multifactorial conditions. Autoinflammatory diseases are classically classified as either monogenic or multifactorial; however, many families do not fit into these models. This hypothesis proposes that the modes of inheritance of autoinflammatory diseases are distributed along a continuum involving mutation type, number of genes and environmental factors. VUS, variants of unknown significance. The thickness of the line depicts the strength of the environmental effect. This figure is only reproduced in colour in the online version.

VUS (more than 20 publications for E148Q) associated with other inflammatory disorders such as juvenile idiopathic arthritis.^{111–113} This review demonstrated that at least part of the controversy relies on nomenclature. Prior to gene discovery, a clinical definition based on established criteria was generally assigned for most of the autoinflammatory diseases (FMF, FCAS, etc). However, the history of the names of the diseases reflects the trend towards a nomenclature which includes the name of the mutated protein. In some cases, a pathophysiological definition has been (re)assigned subsequent to gene discovery (eg, TRAPS, MKD). In parallel, large scale collaborative and country-specific studies confirmed recurrent unproved hypotheses, for example, the role of environment. Other questions such as the true existence of dominant FMF, the influence of epigenetic factors and oligogenism remain unanswered.

While the discovery of these genes opened up new ways of diagnosing these diseases through the molecular screening of mutations, the large accumulation of sequence data demonstrates that interpretation of genetic diagnosis and counselling is becoming increasingly difficult. This advocates the continuous efforts toward a network of multidisciplinary specialists, including molecular biologists, epidemiologists, clinicians, statisticians and web scientists, in order to: (i) elaborate registries devoted to gathering a critical mass of genetic and clinical data such as Infevers^{9–11} or Eurofever^{34,35} and (ii) promote high throughput sequencing which will improve patient care through better understanding of disease-specific defective signalling pathways and subsequent adaptation of patient therapies.

Acknowledgements The author wishes to thank Joanne Strawbridge (Angloscribe) and V Macioce for English language editing and Drs L Cuisset, S Grandemange and M Barat for constructive reviewing of the paper.

Contributors IT is responsible for the overall content as both author and guarantor. IT had the idea for the article, performed the literature search and wrote the article.

Funding This work was supported by the French Ministry of Health and the Montpellier University Hospital Center.

Competing interests None.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- McDermott MF, Aksentjevich I. The autoinflammatory syndromes. *Curr Opin Allergy Clin Immunol* 2002;2:511–16.
- McDermott MF, Aksentjevich I, Galon J, McDermott EM, Ogunkolade BW, Centola M, Mansfield E, Gadina M, Karenko L, Pettersson T, McCarthy J, Frucht DM, Aringer M, Torosyan Y, Teppo AM, Wilson M, Karaarslan HM, Wan Y, Todd I, Wood G, Schlimgen R, Kumarajeewa TR, Cooper SM, Vella JP, Kastner DL. Germline mutations in the extracellular domains of the 55 kDa TNF receptor, TNFR1, define a family of dominantly inherited autoinflammatory syndromes. *Cell* 1999;97:133–44.
- McGonagle D, Aziz A, Dickie LJ, McDermott MF. An integrated classification of pediatric inflammatory diseases, based on the concepts of autoinflammation and the immunological disease continuum. *Pediatr Res* 2009;65(5 Pt 2):38R–45R.
- Gul A. Behcet's disease as an autoinflammatory disorder. *Curr Drug Targets Inflamm Allergy* 2005;4:81–3.
- Kastner DL, Aksentjevich I, Goldbach-Mansky R. Autoinflammatory disease reloaded: a clinical perspective. *Cell* 2010;140:784–90.
- Martinon F, Hofmann K, Tschopp J. The pyrin domain: a possible member of the death domain-fold family implicated in apoptosis and inflammation. *Curr Biol* 2001;11:R118–20.
- Consortium TFF. A candidate gene for familial Mediterranean fever. *Nat Genet* 1997;17:25–31.
- The International FMF Consortium. Ancient missense mutations in a new member of the RoRet gene family are likely to cause familial Mediterranean fever. *Cell* 1997;90:797–807.
- Milhavet F, Cuisset L, Hoffman HM, Slim R, El-Shanti H, Aksentjevich I, Lesage S, Waterham H, Wise C, de Menthier C Sarrauste, Toutou I. The infevers autoinflammatory mutation online registry: update with new genes and functions. *Hum Mutat* 2008;29:803–8.
- Sarrauste de Menthier C, Terriere S, Pugnere D, Ruiz M, Demaille J, Toutou I. INFEVERS: the registry for FMF and hereditary inflammatory disorders mutations. *Nucleic Acids Res* 2003;31:282–5.
- Toutou I, Lesage S, McDermott M, Cuisset L, Hoffman H, Dode C, Shoham N, Aganna E, Hugot JP, Wise C, Waterham H, Pugnere D, Demaille J, de Menthier C Sarrauste. Infevers: an evolving mutation database for auto-inflammatory syndromes. *Hum Mutat* 2004;24:194–8.
- Cuisset L, Jeru I, Dumont B, Fabre A, Cochet E, Le Bozec J, Delpech M, Amselem S, Toutou I. Mutations in the autoinflammatory cryopyrin-associated periodic syndrome gene: epidemiological study and lessons from eight years of genetic analysis in France. *Ann Rheum Dis* 2011;70:495–9.
- Federici L, Rittore-Domingo C, Kone-Paut I, Jorgensen C, Rodiere M, Le Quellec A, Toutou I. A decision tree for genetic diagnosis of hereditary periodic fever in unselected patients. *Ann Rheum Dis* 2006;65:1427–32.
- Grateau G. Clinical and genetic aspects of the hereditary periodic fever syndromes. *Rheumatology (Oxford)* 2004;43:410–15.
- Lainka E, Bielak M, Lohse P, Timmann C, Stojanov S, von Kries R, Niehues T, Neudorf U. Familial mediterranean fever in Germany: epidemiological, clinical, and genetic characteristics of a pediatric population. *Eur J Pediatr* 2012.
- Lainka E, Neudorf U, Lohse P, Timmann C, Bielak M, Stojanov S, Huss K, von Kries R, Niehues T. Analysis of cryopyrin-associated periodic syndromes (CAPS) in German children: epidemiological, clinical and genetic characteristics. *Klin Padiatr* 2012;171:1775–85.
- Lainka E, Neudorf U, Lohse P, Timmann C, Bielak M, Stojanov S, Huss K, von Kries R, Niehues T. Incidence and clinical features of hyperimmunoglobulinemia D and periodic fever syndrome (HIDS) and spectrum of mevalonate kinase (MKV) mutations in German children. *Rheumatol Int* 2011;32:3253–60.
- Lainka E, Neudorf U, Lohse P, Timmann C, Stojanov S, Huss K, von Kries R, Niehues T. Incidence of TNFRSF1A mutations in German children: epidemiological, clinical and genetic characteristics. *Rheumatology (Oxford)* 2009;48:987–91.
- Samuels J, Aksentjevich I, Torosyan Y, Centola M, Deng Z, Sood R, Kastner DL. Familial Mediterranean fever at the millennium. Clinical spectrum, ancient mutations, and a survey of 100 American referrals to the National Institutes of Health. *Medicine (Baltimore)* 1998;77:268–97.
- Heller H, Kariv J, Sherf L, Sohar E. Familial Mediterranean fever. *Harefuah* 1955;48:91–4.
- Kalkan G, Demirkaya E, Ozen S. Severity scoring system for paediatric FMF. *Nat Rev Rheumatol* 2012;8:621.
- Kondi A, Hentgen V, Piram M, Letierce A, Guillaume-Czitrom S, Kone-Paut I. Validation of the new paediatric criteria for the diagnosis of familial Mediterranean fever: data from a mixed population of 100 children from the French reference centre for auto-inflammatory disorders. *Rheumatology (Oxford)* 2010;49:2200–3.
- Livneh A, Langevitz P, Zemer D, Zaks N, Kees S, Lidar T, Migdal A, Padeh S, Pras M. Criteria for the diagnosis of familial Mediterranean fever. *Arthritis Rheum* 1997;40:1879–85.
- Yalcinkaya F, Ozen S, Ozcarar ZB, Aktay N, Cakar N, Duzova A, Kasapcopur O, Elhan AH, Doganay B, Ekim M, Kara N, Uncu N, Bakkaloglu A. A new set of criteria for the diagnosis of familial Mediterranean fever in childhood. *Rheumatology (Oxford)* 2009;48:395–8.
- Akarsu AN, Saatci U, Ozen S, Bakkaloglu A, Besbas N, Sarfarazi M. Genetic linkage study of familial Mediterranean fever (FMF) to 16p13.3 and evidence for genetic heterogeneity in the Turkish population. *J Med Genet* 1997;34:573–8.
- Toutou I. The spectrum of Familial Mediterranean Fever (FMF) mutations. *Eur J Hum Genet* 2001;9:473–83.
- Moradian MM, Sarkisian T, Ajrapetyan H, Avanesian N. Genotype-phenotype studies in a large cohort of Armenian patients with familial Mediterranean fever suggest clinical disease with heterozygous MEFV mutations. *J Hum Genet* 2010;55:389–93.
- Marek-Yagel D, Berkun Y, Padeh S, Abu A, Reznik-Wolf H, Livneh A, Pras M, Pras E. Clinical disease among patients heterozygous for familial Mediterranean fever. *Arthritis Rheum* 2009;60:1862–6.
- Booth MG, Chae JJ, Masters SL, Remmers EF, Barham B, Le JM, Barron KS, Holland SM, Kastner DL, Aksentjevich I. Familial Mediterranean fever with a single MEFV mutation: where is the second hit? *Arthritis Rheum* 2009;60:1851–61.
- Kone-Paut I, Hentgen V, Guillaume-Czitrom S, Compeyrot-Lacassagne S, Tran TA, Toutou I. The clinical spectrum of 94 patients carrying a single mutated MEFV allele. *Rheumatology (Oxford)* 2009;48:840–2.
- Aldea A, Campistol JM, Arostegui JJ, Rius J, Maso M, Vives J, Yague J. A severe autosomal-dominant periodic inflammatory disorder with renal AA amyloidosis and colchicine resistance associated to the MEFV H478Y variant in a Spanish kindred: an unusual familial Mediterranean fever phenotype or another MEFV-associated periodic inflammatory disorder? *Am J Med Genet A* 2004;124:67–73.
- Booth DR, Gillmore JD, Lachmann HJ, Booth SE, Bybee A, Soyuturk M, Akar S, Pepys MB, Tunca M, Hawkins PN. The genetic basis of autosomal dominant familial Mediterranean fever. *QJM* 2000;93:217–21.
- Livneh A, Aksentjevich I, Langevitz P, Torosyan Y, N GS, Shinar Y, Pras E, Zaks N, Padeh S, Kastner DL, Pras M. A single mutated MEFV allele in Israeli patients

- suffering from familial Mediterranean fever and Behcet's disease (FMF-BD). *Eur J Hum Genet* 2001;9:191–6.
- 34 Ozen S, Frenkel J, Ruperto N, Gattorno M. The Eurofever project: towards better care for autoinflammatory diseases. *Eur J Pediatr* 2011;170:445–52.
 - 35 Toplak N, Frenkel J, Ozen S, Lachmann HJ, Woo P, Kone-Paut I, De Benedetti F, Neven B, Hofer M, Dolezalova P, Kummerle-Deschner J, Toutou I, Hentgen V, Simon A, Girschick H, Rose C, Wouters C, Vesely R, Arostegui J, Stojanov S, Ozgocan H, Martini A, Ruperto N, Gattorno M. An international registry on autoinflammatory diseases: the Eurofever experience. *Ann Rheum Dis* 2012;71:1177–82.
 - 36 Federici S, Calcagno G, Finetti M, Gallizzi R, Meini A, Vitale A, Caroli F, Cattalini M, Caorsi R, Zulian F, Tommasini A, Insalaco A, Sormani MP, Baldi M, Ceccherini I, Martini A, Gattorno M. Clinical impact of MEV mutations in children with periodic fever in a prevalent western European Caucasian population. *Ann Rheum Dis* 2012;71:1961–5.
 - 37 Gershoni-Baruch R, Broza Y, Brik R. Prevalence and significance of mutations in the familial Mediterranean fever gene in Henoch-Schönlein purpura. *J Pediatr* 2003;143:658–61.
 - 38 Gershoni-Baruch R, Shinawi M, Shamaly H, Katsinetz L, Brik R. Familial Mediterranean fever: the segregation of four different mutations in 13 individuals from one inbred family: genotype-phenotype correlation and intrafamilial variability. *Am J Med Genet* 2002;109:198–201.
 - 39 Grandemange S, Soler S, Toutou I. Expression of the familial Mediterranean fever gene is regulated by nonsense-mediated decay. *Hum Mol Genet* 2009;18:4746–55.
 - 40 Ustek D, Ekmekci C, Oku B, Cosan F, Cakiris A, Abaci N, Celik S, Kamali S, Hatemi G, Kasapcopur O, Ozdogan H, Gul A. MEV gene 3'-UTR Alu repeat polymorphisms in patients with familial Mediterranean fever. *Clin Exp Rheumatol* 2008;26(4 Suppl 50):S72–76.
 - 41 Notaricola C, Boizet-Bonhoure B, de Santa Barbara P, Osta MA, Cattani D, Toutou I. Characterization of new mutations in the 5'-flanking region of the familial Mediterranean fever gene. *Genes Immun* 2009;10:273–9.
 - 42 Aganna E, Hammond L, Hawkins PN, Aldea A, McKee SA, van Amstel HK, Mischung C, Kusuhara K, Saulsbury FT, Lachmann HJ, Bybee A, McDermott EM, La Regina M, Arostegui JI, Campistol JM, Worthington S, High KP, Molloy MG, Baker N, Bidwell JL, Castaner JL, Whiteford ML, Janssens-Korpola PL, Manna R, Powell RJ, Woo P, Solis P, Minden K, Frenkel J, Yague J, Mirakian RM, Hitman GA, McDermott MF. Heterogeneity among patients with tumor necrosis factor receptor-associated periodic syndrome phenotypes. *Arthritis Rheum* 2003;48:2632–44.
 - 43 Kone-Paut I, Sanchez E, Le Quellec A, Manna R, Toutou I. Autoinflammatory gene mutations in Behcet's disease. *Ann Rheum Dis* 2007;66:832–4.
 - 44 Borghini S, Fiore M, Di Duca M, Caroli F, Finetti M, Santamaria G, Ferlito F, Bua F, Picco P, Obici L, Martini A, Gattorno M, Ceccherini I. Candidate genes in patients with autoinflammatory syndrome resembling tumor necrosis factor receptor-associated periodic syndrome without mutations in the TNFRSF1A gene. *J Rheumatol* 2011;38:1378–84.
 - 45 Aksentijevich I, Nowak M, Mallah M, Chae JJ, Watford WT, Hofmann SR, Stein L, Russo R, Goldsmith D, Dent P, Rosenberg HF, Austin F, Remmers EF, Balow JE Jr, Rosenzweig S, Komarow H, Shoham NG, Wood G, Jones J, Mangra N, Carrero H, Adams BS, Moore TL, Schikler K, Hoffman H, Lovell DJ, Lipnick R, Barron K, O'Shea JJ, Kastner DL, Goldbach-Mansky R. De novo CIAS1 mutations, cytokine activation, and evidence for genetic heterogeneity in patients with neonatal-onset multisystem inflammatory disease (NOMID): a new member of the expanding family of pyrin-associated autoinflammatory diseases. *Arthritis Rheum* 2002;46:3340–8.
 - 46 Bergman F, Warmenius S. Familial perireticular amyloidosis in a Swedish family. *Am J Med* 1968;45:601–6.
 - 47 Williamson LM, Hull D, Mehta R, Reeves WG, Robinson BH, Toghiani PJ. Familial Hibernian fever. *Q J Med* 1982;51:469–80.
 - 48 Gertz MA, Petitt RM, Perrault J, Kyle RA. Autosomal dominant familial Mediterranean fever-like syndrome with amyloidosis. *Mayo Clin Proc* 1987;62:1095–100.
 - 49 Karenko L, Pettersson T, Roberts P. Autosomal dominant 'Mediterranean fever' in a Finnish family. *J Intern Med* 1992;232:365–9.
 - 50 Houten SM, Frenkel J, Waterham HR. Isoprenoid biosynthesis in hereditary periodic fever syndromes and inflammation. *Cell Mol Life Sci* 2003;60:1118–34.
 - 51 Aksentijevich I, Putnam CD, Remmers EF, Mueller JL, Le J, Kolodner RD, Moak Z, Chuang M, Austin F, Goldbach-Mansky R, Hoffman HM, Kastner DL. The clinical continuum of cryopyrinopathies: novel CIAS1 mutations in North American patients and a new cryopyrin model. *Arthritis Rheum* 2007;56:1273–85.
 - 52 Dode C, Cuisset L, Delpech M, Grateau G. TNFRSF1A-associated periodic syndrome (TRAPS), Muckle-Wells syndrome (MWS) and renal amyloidosis. *J Nephrol* 2003;16:435–7.
 - 53 Izawa K, Hijikata A, Tanaka N, Kawai T, Saito MK, Goldbach-Mansky R, Aksentijevich I, Yasumi T, Nakahata T, Heike T, Nishikomori R, Ohara O. Detection of base substitution-type somatic mosaicism of the NLRP3 gene with >99.9% statistical confidence by massively parallel sequencing. *DNA Res* 2012;19:143–52.
 - 54 Tanaka N, Izawa K, Saito MK, Sakuma M, Oshima K, Ohara O, Nishikomori R, Morimoto T, Kambe N, Goldbach-Mansky R, Aksentijevich I, de Saint Basile G, Neven B, van Gijn M, Frenkel J, Arostegui JI, Yague J, Merino R, Ibanez M, Pontillo A, Takada H, Imagawa T, Kawai T, Yasumi T, Nakahata T, Heike T. High incidence of NLRP3 somatic mosaicism in patients with chronic infantile neurologic, cutaneous, articular syndrome: results of an International Multicenter Collaborative Study. *Arthritis Rheum* 2011;63:3625–32.
 - 55 Arostegui JI, Saldana MD Lopez, Pascal M, Clemente D, Aymerich M, Balaguer F, Goel A, del Castillo C, Fournier, Rius J, Plaza S, Robledillo JC Lopez, Juan M, Ibanez M, Yague J. A somatic NLRP3 mutation as a cause of a sporadic case of chronic infantile neurologic, cutaneous, articular syndrome/neonatal-onset multisystem inflammatory disease: novel evidence of the role of low-level mosaicism as the pathophysiologic mechanism underlying mendelian inherited diseases. *Arthritis Rheum* 2010;62:1158–66.
 - 56 Jeru I, Duquesnoy P, Fernandes-Alnemri T, Cochet E, Yu JW, Lackmy-Port-Lis M, Grimprel E, Landman-Parker J, Hentgen V, Marlin S, McElreavey K, Sarkisian T, Grateau G, Alnemri ES, Amselem S. Mutations in NALP12 cause hereditary periodic fever syndromes. *Proc Natl Acad Sci USA* 2008;105:1614–19.
 - 57 Aksentijevich I, Masters SL, Ferguson PJ, Dancy P, Frenkel J, van Royen-Kerkhoff A, Laxer R, Tedgard U, Cowen EW, Pham TH, Booty M, Estes JD, Sandler NG, Plass N, Stone DL, Turner ML, Hill S, Butman JA, Schneider R, Babyn P, El-Shanti H, Pope E, Barron K, Bing X, Laurence A, Lee CC, Chapelle D, Clarke GI, Ohson K, Nicholson M, Gadina M, Yang B, Korman BD, Gregersen PK, van Hagen PM, Hak AE, Huizing M, Rahman P, Douek DC, Remmers EF, Kastner DL, Goldbach-Mansky R. An autoinflammatory disease with deficiency of the interleukin-1-receptor antagonist. *N Engl J Med* 2009;360:2426–37.
 - 58 Frenkel J, Houten SM, Waterham HR, Wanders RJ, Rijkers GT, Kimpen JL, Duran R, Poll-The BT, Kuis W. Mevalonate kinase deficiency and Dutch type periodic fever. *Clin Exp Rheumatol* 2000;18:525–32.
 - 59 Houten SM, Kuis W, Duran R, de Koning TJ, van Royen-Kerkhof A, Romeijn GJ, Frenkel J, Dorland L, de Barse MM, Huijbers WA, Rijkers GT, Waterham HR, Wanders RJ, Poll-The BT. Mutations in MVK, encoding mevalonate kinase, cause hyperimmunoglobulinemia D and periodic fever syndrome. *Nat Genet* 1999;22:175–7.
 - 60 Chen K, Cerutti A. New insights into the enigma of immunoglobulin D. *Immunol Rev* 2010;237:160–79.
 - 61 Di Rocco M, Caruso U, Waterham HR, Picco P, Loy A, Wanders RJ. Mevalonate kinase deficiency in a child with periodic fever and without hyperimmunoglobulinemia D. *J Inher Metab Dis* 2001;24:411–12.
 - 62 Simon A, van Deuren M, Tighe PJ, van der Meer JW, Drenth JP. Genetic analysis as a valuable key to diagnosis and treatment of periodic fever. *Arch Intern Med* 2001;161:2491–3.
 - 63 Ammouri W, Cuisset L, Rouaghe S, Rolland MO, Delpech M, Grateau G, Ravet N. Diagnostic value of serum immunoglobulinemia D level in patients with a clinical suspicion of hyper IgD syndrome. *Rheumatology (Oxford)* 2007;46:1597–600.
 - 64 Saulsbury FT. Hyperimmunoglobulinemia D and periodic fever syndrome (HIDS) in a child with normal serum IgD, but increased serum IgA concentration. *J Pediatr* 2003;143:127–9.
 - 65 Cuisset L, Drenth JP, Simon A, Vincent MF, van der Velde, Visser, van der Meer S, Grateau JW, Delpech GM. Molecular analysis of MVK mutations and enzymatic activity in hyper-IgD and periodic fever syndrome. *Eur J Hum Genet* 2001;9:260–6.
 - 66 Zhang SQ, Jiang T, Li M, Zhang X, Ren YQ, Wei SC, Sun LD, Cheng H, Li Y, Yin XY, Hu ZM, Wang ZY, Liu Y, Guo BR, Tang HY, Tang XF, Ding YT, Wang JB, Li P, Wu BY, Wang W, Yuan XF, Hou JS, Ha WW, Wang WJ, Zhai YJ, Wang J, Qian FF, Zhou FS, Chen G, Zuo XB, Zheng XD, Sheng YJ, Gao JP, Liang B, Zhu J, Xiao FL, Wang PG, Cui Y, Li H, Liu SX, Gao M, Fan X, Shen SK, Zeng M, Sun GQ, Xu Y, Hu JC, He TT, Li YR, Yang HM, Yu ZY, Zhang HF, Hu X, Yang K, Zhao SX, Zhou YW, Liu JJ, Du WD, Zhang L, Xia K, Yang S, Zhang XJ. Exome sequencing identifies MVK mutations in disseminated superficial actinic porokeratosis. *Nat Genet* 2012;44:1156–60.
 - 67 Tarp BD, Obel N. Familial Mediterranean fever with pseudodominant inheritance. *Ugeskr Laeger* 2000;162:6261–2.
 - 68 Hospach T, Lohse P, Heilbronner H, Dannecker GE. Pseudodominant inheritance of the hyperimmunoglobulinemia D with periodic fever syndrome in a mother and her two monozygotic twins. *Arthritis Rheum* 2005;52:3606–10.
 - 69 Toutou I, Ben-Chetrit E, Notaricola C, Domingo C, Dewalle M, Dross C, Dupont M, Demaille J, Rosner I, Rozenbaum M. Familial Mediterranean fever clinical and genetic features in Druzes and in Iraqi Jews: a preliminary study. *J Rheumatol* 1998;25:916–19.
 - 70 Ravet N, Rouaghe S, Dode C, Bienvenu J, Stirnemann J, Levy P, Delpech M, Grateau G. Clinical significance of P46L and R92Q substitutions in the tumour necrosis factor superfamily 1A gene. *Ann Rheum Dis* 2006;65:1158–62.
 - 71 Toutou I, Notaricola C, Grandemange S. Identifying mutations in autoinflammatory diseases: towards novel genetic tests and therapies? *Am J Pharmacogenomics* 2004;4:109–18.
 - 72 Toutou I, Perez C, Dumont B, Federici L, Jorgensen C. Refractory auto-inflammatory syndrome associated with digenic transmission of low-penetrance

- tumour necrosis factor receptor-associated periodic syndrome and cryopyrin-associated periodic syndrome mutations. *Ann Rheum Dis* 2006;65:1530–1.
- 73 Kaipainen-Seppanen O, Punnonen K, van Gijn ME, Mononen T. Two pathogenic CIAS1 mutations and plasma cytokine profile in a Finnish patient with familial cold autoinflammatory syndrome responsive to anakinra. *Scand J Rheumatol* 2008;37:75–6.
 - 74 Aksentijevich I, Galon J, Soares M, Mansfield E, Hull K, Oh HH, Goldbach-Mansky R, Dean J, Athreya B, Reginato AJ, Henrickson M, Pons-Estel B, O'Shea JJ, Kastner DL. The tumor-necrosis-factor receptor-associated periodic syndrome: new mutations in TNFRSF1A, ancestral origins, genotype-phenotype studies, and evidence for further genetic heterogeneity of periodic fevers. *Am J Hum Genet* 2001;69:301–14.
 - 75 Medlej-Hashim M, Delague V, Chouery E, Salem N, Rawashdeh M, Lefranc G, Loiselet J, Megarbane A. Amyloidosis in familial Mediterranean fever patients: correlation with MEFV genotype and SAA1 and MICA polymorphisms effects. *BMC Med Genet* 2004;5:4.
 - 76 Turkcapar N, Tuncali T, Kutlay S, Burhan BY, Kinikli G, Erturk S, Duman M. The contribution of genotypes at the MICA gene triplet repeat polymorphisms and MEFV mutations to amyloidosis and course of the disease in the patients with familial Mediterranean fever. *Rheumatol Int* 2007;27:545–51.
 - 77 Cazeneuve C, Ajrapetyan H, Papin S, Roudot-Thoraval F, Genevieve D, Mndjoyan E, Papazian M, Sarkisian A, Babloyan A, Boissier B, Duquesnoy P, Kouyoumdjian JC, Girodon-Boulant E, Grateau G, Sarkisian T, Amselem S. Identification of MEFV-independent modifying genetic factors for familial Mediterranean fever. *Am J Hum Genet* 2000;67:1136–43.
 - 78 Gershoni-Baruch R, Brik R, Zacks N, Shinawi M, Lidar M, Livneh A. The contribution of genotypes at the MEFV and SAA1 loci to amyloidosis and disease severity in patients with familial Mediterranean fever. *Arthritis Rheum* 2003;48:1149–55.
 - 79 Bakaloglu A, Duzova A, Ozen S, Balci B, Besbas N, Topaloglu R, Ozaltin F, Yilmaz E. Influence of Serum Amyloid A (SAA1) and SAA2 gene polymorphisms on renal amyloidosis, and on SAA/C-reactive protein values in patients with familial mediterranean fever in the Turkish population. *J Rheumatol* 2004;31:1139–42.
 - 80 Schwabe AD, Peters RS. Familial Mediterranean Fever in Armenians. Analysis of 100 cases. *Medicine (Baltimore)* 1974;53:453–62.
 - 81 Toutou I, Sarkisian T, Medlej-Hashim M, Tunca M, Livneh A, Cattani D, Yalcinkaya F, Ozen S, Majeed H, Ozdogan H, Kastner D, Booth D, Ben-Chetrit E, Pugner D, Michelon C, Seguret F, Gershoni-Baruch R. Country as the primary risk factor for renal amyloidosis in familial mediterranean fever. *Arthritis Rheum* 2007;56:1706–12.
 - 82 Ozen S, Aktay N, Laina E, Duzova A, Bakaloglu A, Kallinich T. Disease severity in children and adolescents with familial Mediterranean fever: a comparative study to explore environmental effects on a monogenic disease. *Ann Rheum Dis* 2009;68:246–8.
 - 83 Ben-Zvi I, Brandt B, Berkun Y, Lidar M, Livneh A. The relative contribution of environmental and genetic factors to phenotypic variation in familial Mediterranean fever (FMF). *Gene* 491:260–3.
 - 84 Kiretepe AK, Kasapcopur O, Arisoy N, Erdem G, Celikyapi, Hatemi G, Ozdogan H, Turanlı E Tahir. Analysis of MEFV exon methylation and expression patterns in familial Mediterranean fever. *BMC Med Genet* 2011;12:105.
 - 85 Agarwal AK, Xing C, DeMartino GN, Mizrahi D, Hernandez MD, Sousa AB, de Villarreal L, Martinez, dos Santos HG, Garg A. PSMB8 encoding the beta5i proteasome subunit is mutated in joint contractures, muscle atrophy, microcytic anemia, and panniculitis-induced lipodystrophy syndrome. *Am J Hum Genet* 2010;87:866–72.
 - 86 Liu Y, Ramot Y, Torrello A, Paller AS, Si N, Babay S, Kim PW, Sheikh A, Lee CC, Chen Y, Vera A, Zhang X, Goldbach-Mansky R, Zlotogorski A. Mutations in proteasome subunit beta type 8 cause chronic atypical neutrophilic dermatosis with lipodystrophy and elevated temperature with evidence of genetic and phenotypic heterogeneity. *Arthritis Rheum* 2011;64:895–907.
 - 87 Arima K, Kinoshita A, Mishima H, Kanazawa N, Kaneko T, Mizushima T, Ichinose K, Nakamura H, Tsujino A, Kawakami A, Matsunaka M, Kasagi S, Kawano S, Kumagai S, Ohmura K, Mimori T, Hirano M, Ueno S, Tanaka K, Tanaka M, Toyoshima I, Sugino H, Yamakawa A, Niikawa N, Furukawa F, Murata S, Eguchi K, Ida H, Yoshiura K. Proteasome assembly defect due to a proteasome subunit beta type 8 (PSMB8) mutation causes the autoinflammatory disorder, Nakajo-Nishimura syndrome. *Proc Natl Acad Sci USA* 2011;108:14914–19.
 - 88 Masters SL, Simon A, Aksentijevich I, Kastner DL. Horror autoinflammaticus: the molecular pathophysiology of autoinflammatory disease (*). *Annu Rev Immunol* 2009;27:621–68.
 - 89 Jeru I, Hentgen V, Normand S, Duquesnoy P, Cochet E, Delwail A, Grateau G, Marlin S, Amselem S, Lecron JC. Role of interleukin-1beta in NLRP12-associated autoinflammatory disorders and resistance to anti-interleukin-1 therapy. *Arthritis Rheum* 2011;63:2142–8.
 - 90 Ter Haar N, Lachmann H, Ozen S, Woo P, Uziel Y, Modesto C, Kone-Paut I, Cantarini L, Insalaco A, Neven B, Hofer M, Rigante D, Al-Mayouf S, Toutou I, Gallizzi R, Papadopoulou-Alataki E, Martino S, Kuemmerle-Deschner J, Obici L, lagaru N, Simon A, Nielsen S, Martini A, Ruperto N, Gattorno M, Frenkel J. Treatment of autoinflammatory diseases: results from the Eurofever Registry and a literature review. *Ann Rheum Dis* Published Online First: 29 June 2012. doi:10.1136/annrheumdis-2011-201268
 - 91 Marrakchi S, Guigue P, Renshaw BR, Puel A, Pei XY, Fraitaig S, Zribi J, Bal E, Cluzeau C, Chrabieh M, Towne JE, Douangpanya J, Pons C, Mansour S, Serre V, Makni H, Mahfoudh N, Fakhfakh F, Bodemer C, Feingold J, Hadj-Rabia S, Favre M, Genin E, Sahbatou M, Munnich A, Casanova JL, Sims JE, Turki H, Bachelez H, Smahi A. Interleukin-36-receptor antagonist deficiency and generalized pustular psoriasis. *N Engl J Med* 2011;365:620–8.
 - 92 Setta-Kaffetzi N, Navarini AA, Patel VM, Pullabhatla V, Pink AE, Choon SE, Allen MA, Burden AD, Griffiths CE, Seyger MM, Kirby B, Trembath RC, Simpson MA, Smith CH, Capon F, Barker JN. Rare pathogenic variants in IL36RN underlie a spectrum of psoriasis-associated pustular phenotypes. *J Invest Dermatol* 2013 Jan 10. doi:10.1038/jid.2012.490. [Epub ahead of print].
 - 93 Jordan CT, Cao L, Roberson ED, Duan S, Helms CA, Nair RP, Duffin KC, Stuart PE, Goldgar D, Hayashi G, Olsson EH, Feng BJ, Pullinger CR, Kane JP, Wise CA, Goldbach-Mansky R, Lowes MA, Peddle L, Chandran V, Liao W, Rahman P, Krueger GG, Gladman D, Elder JT, Menter A, Bowcock AM. Rare and common variants in CARD14, encoding an epidermal regulator of NF-kappaB, in psoriasis. *Am J Hum Genet* 2012;90:796–808.
 - 94 Jordan CT, Cao L, Roberson ED, Pierson KC, Yang CF, Joyce CE, Ryan C, Duan S, Helms CA, Liu Y, Chen Y, McBride AA, Hwu WL, Wu JY, Chen YT, Menter A, Goldbach-Mansky R, Lowes MA, Bowcock AM. PSORS2 is due to mutations in CARD14. *Am J Hum Genet* 2012;90:784–95.
 - 95 Fuchs-Telem D, Sarig O, van Steensel MA, Isakov O, Israeli S, Nussbeck J, Richard K, Winpenningckx V, Vernooij M, Shomron N, Uitto J, Fleckman P, Richard G, Sprecher E. Familial pityriasis rubra pilaris is caused by mutations in CARD14. *Am J Hum Genet* 2012;91:163–70.
 - 96 Zhou Q, Lee GS, Brady J, Datta S, Katam M, Sheikh A, Martins MS, Bunney TD, Santich BH, Moir S, Kuhns DB, Priel DA, Ombrello A, Stone D, Ombrello MJ, Khan J, Milner JD, Kastner DL, Aksentijevich I. A hypermorphic missense mutation in PLCG2, encoding phospholipase Cgamma2, causes a dominantly inherited autoinflammatory disease with immunodeficiency. *Am J Hum Genet* 2012;91:713–20.
 - 97 Ombrello MJ, Remmers EF, Sun G, Freeman AF, Datta S, Torabi-Parizi P, Subramanian N, Bunney TD, Baxendale RW, Martins MS, Romberg N, Komarow H, Aksentijevich I, Kim HS, Ho J, Cruse G, Jung MY, Gilfillan AM, Metcalfe DD, Nelson C, O'Brien M, Wisch L, Stone K, Douek DC, Gandhi C, Wanderer AA, Lee H, Nelson SF, Shianna KV, Cirulli ET, Goldstein DB, Long EO, Moir S, Meffre E, Holland SM, Kastner DL, Katam M, Hoffman HM, Milner JD. Cold urticaria, immunodeficiency, and autoimmunity related to PLCG2 deletions. *N Engl J Med* 2012;366:330–8.
 - 98 Ombrello MJ, Kastner DL, Milner JD. HOIL and water: the two faces of HOIL-1 deficiency. *Nat Immunol* 2012;13:1133–5.
 - 99 Boisson B, Laplantine E, Prando C, Giliani S, Israelsson E, Xu Z, Abhyankar A, Israel L, Trevejo-Nunez G, Bogunovic D, Cepika AM, Macduff D, Chrabieh M, Hubeau M, Bajolle F, Debre M, Mazzolari E, Vairo D, Agou F, Virgin HW, Bossuyt X, Rambaud C, Facchetti F, Bonnet D, Quartier P, Fournet JC, Pascual V, Chaussabel D, Notarangelo LD, Puel A, Israel A, Casanova JL, Picard C. Immunodeficiency, autoinflammation and amylopectinosis in humans with inherited HOIL-1 and LUBAC deficiency. *Nat Immunol* 2012;13:1178–86.
 - 100 Ben-Chetrit E, Peleg H, Aamar S, Heyman SN. The spectrum of MEFV clinical presentations—is it familial Mediterranean fever only? *Rheumatology (Oxford)* 2009;48:1455–9.
 - 101 Ting JP, Lovering RC, Alnemri ES, Bertin J, Boss JM, Davis BK, Flavell RA, Girardin SE, Godzik A, Harton JA, Hoffman HM, Hugot JP, Inohara N, Mackenzie A, Maltas LJ, Nunez G, Ogura Y, Otten LA, Philpott D, Reed JC, Reith W, Schreiber S, Steimle V, Ward PA. The NLR gene family: a standard nomenclature. *Immunity* 2008;28:285–7.
 - 102 Hugot JP, Chamaillard M, Zouali H, Lesage S, Cezard JP, Belaiche J, Almer S, Tysk C, O'Morain CA, Gassull M, Binder V, Finkel Y, Cortot A, Modigliani R, Laurent-Puig P, Gower-Rousseau C, Macry J, Colombel JF, Sahbatou M, Thomas G. Association of NOD2 leucine-rich repeat variants with susceptibility to Crohn's disease. *Nature* 2001;411:599–603.
 - 103 Miceli-Richard C, Lesage S, Rybojad M, Prieur AM, Manouvrier-Hanu S, Hafner R, Chamaillard M, Zouali H, Thomas G, Hugot JP. CARD15 mutations in Blau syndrome. *Nat Genet* 2001;29:19–20.
 - 104 Cattani D. MEFV mutation carriers and diseases other than familial mediterranean fever: proved and non-proved associations; putative biological advantage. *Curr Drug Targets Inflamm Allergy* 2005;4:105–12.
 - 105 Uslu N, Yuce A, Demir H, Saltik-Temizel IN, Usta Y, Yilmaz E, Besbas N, Gurakan F, Ozen H, Ozen S. The association of inflammatory bowel disease and mediterranean fever gene (MEFV) mutations in Turkish children. *Dig Dis Sci* 2010;55:3488–94.
 - 106 Villani AC, Lemire M, Fortin G, Louis E, Silverberg MS, Collette C, Baba N, Libouille C, Belaiche J, Bitton A, Gaudet D, Cohen A, Langelier D, Fortin PR, Wither JE, Sarfati M, Rutgeerts P, Rioux JD, Vermeire S, Hudson TJ,

- Franchimont D. Common variants in the NLRP3 region contribute to Crohn's disease susceptibility. *Nat Genet* 2009;41:71–6.
- 107 Villani AC, Lemire M, Louis E, Silverberg MS, Collette C, Fortin G, Nimmo ER, Renaud Y, Brunet S, Libioulle C, Belaiche J, Bitton A, Gaudet D, Cohen A, Langelier D, Rioux JD, Arnott ID, Wild GE, Rutgeerts P, Satsangi J, Vermeire S, Hudson TJ, Franchimont D. Genetic variation in the familial Mediterranean fever gene (MEFV) and risk for Crohn's disease and ulcerative colitis. *PLoS One* 2009;4:e7154.
- 108 Amoura Z, Dode C, Hue S, Caillaud-Zucman S, Bahram S, Delpech M, Grateau G, Wechsler B, Piette JC. Association of the R92Q TNFRSF1A mutation and extracranial deep vein thrombosis in patients with Behcet's disease. *Arthritis Rheum* 2005;52:608–11.
- 109 Caminero A, Comabella M, Montalban X. Role of tumour necrosis factor (TNF)-alpha and TNFRSF1A R92Q mutation in the pathogenesis of TNF receptor-associated periodic syndrome and multiple sclerosis. *Clin Exp Immunol* 2011;166:338–45.
- 110 Kingsbury SR, Conaghan PG, McDermott MF. The role of the NLRP3 inflammasome in gout. *J Inflamm Res* 2011;4:39–49.
- 111 Ayaz NA, Ozen S, Bilginer Y, Erguven M, Taskiran E, Yilmaz E, Besbas N, Topaloglu R, Bakaloglu A. MEFV mutations in systemic onset juvenile idiopathic arthritis. *Rheumatology (Oxford)* 2009;48:23–5.
- 112 Day TG, Ramanan AV, Hinks A, Lamb R, Packham J, Wise C, Punaro M, Donn RP. Autoinflammatory genes and susceptibility to psoriatic juvenile idiopathic arthritis. *Arthritis Rheum* 2008;58:2142–6.
- 113 Ozen S, Bakaloglu A, Yilmaz E, Duzova A, Balci B, Topaloglu R, Besbas N. Mutations in the gene for familial Mediterranean fever: do they predispose to inflammation? *J Rheumatol* 2003;30:2014–18.
- 114 Soriano A, Manna R. Familial Mediterranean fever: new phenotypes. *Autoimmun Rev* 2012;12:31–7.
- 115 Shinar Y, Obici L, Aksentijevich I, Bennetts B, Austrup F, Ceccherini I, Costa JM, De Leener A, Gattorno M, Kania U, Kone-Paut I, Lezer S, Livneh A, Moix I, Nishikomori R, Ozen S, Phylactou L, Risom L, Rowczenio D, Sarkisian T, van Gijn ME, Witsch-Baumgartner M, Morris M, Hoffman HM, Touitou I. Guidelines for the genetic diagnosis of hereditary recurrent fevers. *Ann Rheum Dis* 2012;71:1599–605.