Haemoglobin Q India ($\alpha 64(\text{E}13)$ Aspartic Acid $\rightarrow$ Histidine) Associated with β -Thalassaemia Observed in Three Sindhi Families

P. K. SUKUMARAN,* S. M. MERCHANT,† MENNA P. DESAI,‡
BARBARA G. WILTSHIRE,‡ and H. LEHMANN‡

Haemoglobin Q was first reported in a Chinese man in whom it was associated with Hb H disease, and therefore with α -thalassaemia (Vella *et al*, 1958). It has been discovered in other people, mostly in south-east Asia, and these reports have been summarized by Sagnet *et al* (1968). Although it was known that Hb Q was an α -chain abnormal haemoglobin with a mutation in the tryptic peptide No. 9 of the α -chain (αTpIX) which comprises residues 62–90 of the 141 residues of the α -chain, the exact amino-acid substitution had remained undetermined. However, 2 haemoglobins Q were characterized by Lorkin *et al* in 1970. One type came from both a Chinese and a Thai family (in which it was associated with α -thalassaemia) and had the substitution $\alpha 74 \text{ Asp} \rightarrow \text{His}$, and the other, $\alpha 75 \text{ Asp} \rightarrow \text{His}$, came from Iran. Independently, the first variant, described as Hb G Taichung, was found in a Chinese by Blackwell and Liu (1970), and in a Thai, where it was again associated with α -thalassaemia, as Hb Mahidol, by Pootrakul and Dixon (1970). In India only a few cases of Hb H disease have been found, indicating that α -thalassaemia is probably rare; β -thalassaemia is however quite common (Chatterjea, 1966; Chouhan, Sharma, and Parekh, 1970).

We have recently observed 3 Sindhi-speaking Hindu families with children suffering from β -thalassaemia major. Family studies showed some members with β -thalassaemia minor and one with β -thalassaemia major associated with Hb Q. This was shown to arise from an amino-acid substitution $\alpha 64 \text{ Asp} \rightarrow \text{His}$, and was therefore different both from the Q Thailand ($\alpha 74 \text{ Asp} \rightarrow \text{His}$) and Q Iran ($\alpha 75 \text{ Asp} \rightarrow \text{His}$). A preliminary report of this new Hb Q

India was presented in 1971 by Sukumaran, Wiltshire, and Lehmann.

Methods

Haematological studies were carried out using standard methods (Dacie and Lewis, 1968). Osmotic fragility tests were performed with Simmel's tyrode (Sanghvi, Sukumaran, and Lehmann, 1958), and Hb F levels estimated according to Singer, Chernoff, and Singer (1951). Paper electrophoresis of the haemoglobins was carried out at pH 8.9 using Tris buffer (Cradock-Watson, Fenton, and Lehmann, 1959), and starch gel electrophoresis with discontinuous Tris-citrate and borate buffer at pH 8.6 (Poulik, 1959). Haemoglobin A₂ levels were determined using cellulose acetate strips (Marengo-Rowe, 1965). The material available was sometimes so small that Hb Q₂ could not be quantitatively measured.

The abnormal haemoglobin was purified by repeated paper electrophoresis and by column chromatography using DEAE Sephadex (Huisman and Dozy, 1965). Preliminary studies for the identification of the mutant tryptic peptide were made using the methods summarized by Sick *et al* (1967). For thermolysin digestion the mutant peptide isolated from 55 mg of globin was eluted with 0.5 N NH₄OH and dissolved in 0.8 ml of 0.25 M ammonium acetate buffer, pH 8.6. Thermolysin (0.2 mg in 0.1 ml of 50 mM barium acetate) was added and the peptide incubated for 5 hr at 37°C. To stop the reaction, barium was precipitated with 0.1 ml of 0.1 N H₂SO₄. For pepsin digestion the mutant peptide from 20 mg globin was eluted with 0.05 N NH₄OH and dissolved in 0.9 ml of 0.05 N HCl. Pepsin (0.2 mg in 0.1 ml of water) was added and the peptide incubated for 5 hr at 37°C. The reaction was stopped by immersion in boiling water. After both thermolysin and pepsin digestions the supernatant was divided into 2 portions of 0.2 and 0.8 ml for diagnostic and preparative fingerprints, respectively.

Case Histories

Family A. A 1½-year-old boy (II.4) who had developed normally for one year, was admitted to the Jerbai Wadia Hospital for Children, with fever, increasing pallor, and distension of the abdomen. The child

Received 23 June 1972.

* Epidemiology Division, Cancer Research Institute, Tata Memorial Centre, Parel, Bombay 12, India.

† Jerbai Wadia Hospital for Children, Parel, Bombay 12, India.

‡ Medical Research Council Abnormal Haemoglobin Unit, University Department of Biochemistry, Cambridge, UK.

TABLE I
HAEMATOLOGICAL DATA (FAMILY A)

Case	Age (yr)	Hb (g/100 ml)	RBC ($\times 10^9/\mu$ l)	MVC (fl)	MCH (pg)	MCHC (%)	Reticulocytes (%)	Red Cell Morphology			Osmotic Fragility (% lysis)*	Hb F (%)	Electrophoretic Pattern (when quantitated % Q, A ₂ , and Q ₂ in parentheses),
								Vacuolation	Aniso Poikilo	Target Cells			
I.1	40	11.0	6.09	74.0	24.4	28.9	1.0	++	+	+	48.5	1.6	A + Q(8.6) + A(4.3) + Q ₂
I.2	33	11.0	5.28	72.0	20.8	29.0	0.6	++	+	+	62.0	1.5	A + A ₂ (5.2)
II.1	12	10.2	5.68	63.5	18.1	28.3	0.8	++	+	+	51.3	1.4	A + Q(8.0) + A ₂ (3.3) + Q ₂
II.2	9	12.5	4.55	83.5	27.5	32.8	1.4	—	—	—	96.5	0.5	A + A ₂ (2.6)
II.3	5	12.2	4.13	88.5	29.6	33.5	0.2	—	—	—	88.0	0.7	A + Q(19.6) + A ₂ (1.1) + Q ₂
II.4	1½	{ 4.8 7.6	—	—	—	—	3.2 1.8	+++ ++	++ ++	++ ++	71.5	32 24	A + F before transfusion A + F + A ₂ (2.4) after transfusion

* Percentage lysis in 0.4 tyrode solution (normal range 80–100%).
† Large number of nucleated red cells.
‡ One week after blood transfusion, occasional nucleated red cells.

FIG. 1. Electrophoretic pattern of haemoglobins from family A (starch gel). Right to left: Mother (I.2); child (II.4) post transfusion sample; father (I.1); Hb AL and Hb AS.

was dyspnoeic, weighed 7.5 kg and was 70 cm tall. His fontanelle was open and frontal and parietal bossing were evident. The spleen was enlarged to 7 cm and the liver to 5 cm below the costal margin. Haematological data of the boy and his family are given in Table I and are consistent with the diagnosis of the boy having β -thalassaemia major. The percentage of alkali resistant haemoglobin was 31.7% after a packed cell blood transfusion.

The father (I.1) was healthy, but blood examination revealed a raised Hb A₂ and other red cell stigmata consistent with β -thalassaemia minor. He also had a slow moving haemoglobin, and starch gel electrophoresis showed the presence of 2 haemoglobins A₂, which suggests that there was an α -chain variant present (Fig. 1). Its mobility on paper and on starch gel electrophoresis indicated that it was a Hb Q. The father was therefore a double heterozygote for β -thalassaemia and Hb Q.

The mother (I.2) was shown to have β -thalassaemia minor. Among the sibs, one sister (II.1) had both Hb Q and β -thalassaemia minor, while another sister (II.3) had 20% Hb Q and no β -thalassaemia (Fig. 2). The latter

FIG. 2. Pedigrees of families A, L, and T.

sister shows no abnormality at all indicating that this level of Hb Q causes no clinical symptoms.

Family L. A male child (II.4) aged 6 months was admitted to hospital looking pale and weak after having grown well up to the age of 4 months. He weighed 6 kg,

was 60 cm tall, and both liver and spleen were enlarged 6 cm below the costal margin. He was assumed to have β -thalassaemia major although x-rays of skull and long bones did not yet show any abnormality.

The haematological data of this child and his family are shown in Table II and can be seen to be in agreement

TABLE II
HAEMATOLOGICAL DATA (FAMILY L)

Case	Age (yr)	Hb (g/100 ml)	RBC ($\times 10^6/\mu\text{l}$)	MCV (fl)	MCH (pg)	MCHC (%)	Reticulocytes (%)	Red Cell Morphology			Osmotic Fragility (% lysis)*	Hb F (%)	Electrophoretic Pattern (when quantitated % Q, A ₂ , and Q ₂ in parentheses)
								Vacuolation	Aniso Poikilo	Target Cells			
I.1	40	13.0	4.47	85.0	29.1	34.2	0.6	—	—	—	100	1.2	A + A ₂ (2.2)
I.2	36	12.2	6.16	58.8	19.9	33.2	1.8	+	—	+	76.5	1.5	A + Q(14.2) + A ₂ (3.5) + Q ₂ (1.6)
I.3	38	11.6	5.45	72.5	21.3	29.4	1.6	+	+	+	31.7	1.4	A + Q(9.6) + A ₂ (3.4) + Q ₂ (0.7)
I.4	30	9.8	5.65	63.8	18.4	27.2	0.6	++	+	+	21.1	1.1	A + A ₂ (4.5)
II.1	8	9.9	5.25	66.5	18.8	28.3	0.6	+	+	Few	21.0	0.9	A + A ₂ (4.5)
II.2	6	10.4	5.20	68.2	20.0	29.3	0.6	++	+	+	15.4	1.7	A + Q(18.4) + A ₂ (3.8) + Q ₂ (1.0)
II.3	4	9.6	5.53	62.2	17.3	27.8	1.2	+	—	+	18.0	1.2	A + A ₂ (4.1)
II.4	$\frac{1}{12}$	5.5	2.92	61.6	18.9	30.6	4.6	+++	+++	+++†	48.5	56	F + Q _F

* Percentage haemolysis in 0.4 tyrode solution (normal range 80–100%).

† Large number of nucleated red cells.

FIG. 3. Electrophoretic pattern of haemoglobins from family L. **A** (starch gel): right to left, mother (I.4); child (II.4); father (I.3); father I.1 of family A; Hb AS. **B** (paper): right to left, thalassaemia major showing Hb F only; child II.4; father I.3; Hb AS.

with the child having β -thalassaemia major. The boy was transfused and given 2.5 mg folic acid by mouth and when discharged had a haemoglobin level of 11.5 g %. Electrophoresis of his haemoglobin on both starch gel and paper (Fig. 3) showed that a slow moving component was present as well as Hb F. It appears from the family studies that this haemoglobin consists of a fetal haemoglobin containing α^a chains instead of α^A chains, that is $\alpha_2^a\gamma_2$; it forms 6.9% of the total haemoglobin.

Both father (I.3) and mother (I.2) had β -thalassaemia minor, and the father's haemoglobin also had a slow moving component. The slow moving haemoglobins of both the father and the son can be seen in Fig. 3. Starch gel electrophoresis distinguishes both between normal Hb A and Hb F, and also between the 2 slow components $\alpha_2^a\beta_2$ and $\alpha_2^a\gamma_2$. The haemoglobin patterns of I.3 in family L and I.1 in family A can be seen to be identical, a double Hb A₂ being visible in both cases. The inheritance of α^a chains and β -thalassaemia in the son can be seen from the family tree (Fig. 2).

Family T. The 3rd child (III.2) was 4 months old and was admitted to hospital because he had been pale for 1 month. He weighed 5 kg, was 62 cm long, and had a spleen enlarged to 7.5 cm below the costal margin. The haematological data of the child and his

family are shown in Table III. These confirmed the diagnosis of β -thalassaemia major in the child, who was transfused, given 5 mg folic acid and discharged after 2 weeks with a haemoglobin level of 11.8 g %.

As Table III shows, both parents had β -thalassaemia minor. The father also had the slow moving haemoglobin Hb Q, which he appeared to have inherited from his mother (I.2). Both of them have a double Hb A₂, but the abnormal band is clearer in the case of the father, since β -thalassaemia has led to a raised Hb A₂ level. The grandmother had no sign of β -thalassaemia minor, but the grandfather (I.1) had β -thalassaemia minor but no slow moving haemoglobin (Fig. 4).

Examination of the Abnormality of the Globin

The haemoglobin Q was purified by paper electrophoresis at pH 8.9, eluted and concentrated. The globin was precipitated using acidic acetone and digested with trypsin. The fingerprints (Fig. 5) of the haemoglobin from I.3 (family L) and II.1 (family A) showed a new peptide containing both histidine and methionine as well as a reduced amount of peptides α^A TpXI and α^A TpVIII-IX. This suggested that the new peptide was a mutant α TpIX which was confirmed by amino-acid analysis. This showed that an aspartic acid or asparagine

TABLE III
HAEMATOLOGICAL DATA (FAMILY T)

Case	Age (yr)	Hb (g/ μ l)	RBC ($\times 10^6/\mu$ l)	MCV (fl)	MCH (pg)	MCHC (%)	Reticulocytes (%)	Red Cell Morphology			Osmotic Fragility (% lysis)*	Hb F (%)	Electrophoretic Pattern (when quantitated % Q, A ₂ , Q ₂ in parentheses)
								Vacuolation	Anisopoikilo	Target Cells			
I.1	55	9.6	4.28	77.0	22.4	29.0	0.1	+	+	+	56.3	1.6	A + A ₂ (5.4)
I.2	45	13.9	4.61	94.2	30.2	32.0	0.3	-	-	-	93.5	0.7	A + Q + A ₂ (2.7) + Q ₂
II.1	30	13.0	6.26	71.4	20.8	29.3	0.6	+	-	+	69.4	1.3	A + Q + A ₂ (4.4) + Q ₂
II.2	23	8.7	5.31	65.0	16.4	25.2	0.5	+	+	Few	72.8	1.6	A + A ₂ (5.1)
III.2	†	5.5	2.64	68.1	20.8	30.6	3.6	+++	++	++	83.3	78	F

* Percentage haemolysis in 0.4 tyrode solution (normal range 80-100%).

† Large number of nucleated red cells.

FIG. 4. Electrophoretic pattern of haemoglobins from family T (starch gel). Right to left: paternal grandfather (I.1); grandmother (I.2); father (II.1); child (III.2), mother (II.2).

had been replaced by histidine in both samples (Table IV). From Table V it can be seen that if there were a single charge change in α TpIX the electrophoretic RF would alter by about 0.1 unit. However, since the electrophoretic RF is increased from 0.0 to 0.2 units, it appears that there is a double charge change; indicating a mutation of aspartic acid $\rightarrow$ histidine as in Hb Q Iran and Hb Q Thailand, because histidine is positively charged under these conditions. The electrophoretic RF was determined taking lysine as unity and the neutral peptides

TABLE V
ELECTROPHORETIC AND CHROMATOGRAPHIC
MOBILITIES (RF VALUES)

Peptide	Electrophoretic RF	Chromatographic RF	Charge Change from α^A IX
α^A IX	0.00	0.14	0
α^A VIII-IX	0.09	0.11	+1
α^Q IranIX	0.19	0.13	+2
α^Q IndiaIX (I.3, family L)	0.21	0.13	+2
α^Q IndiaIX (II.1, family A)	0.22	0.13	+2

FIG. 5. Fingerprint (pH 6.4) of the soluble tryptic peptides of Hb Q India from father (I.3) of family L. (For details of interpretation see text.) * = Point of application.

TABLE IV
AMINO-ACID COMPOSITION OF α^Q TpIX (62-90)

Amino Acid	Residues Found		Expected in Hb A
	Sample (II.1, family A)	Sample (I.3, family L)	
Asp	5.01	5.07	6
Thr	1.04	0.90	1
Ser	1.94	1.77	2
Pro	0.94	1.00	1
Ala	7.35	6.70	7
Val	3.02	2.94	3
Met	0.73	0.51	1*
Leu	4.18	4.02	4
His	3.77	4.30	3
Lys	1.11	1.33	1
Yield (n-moles)	20.3	37.0	

* Methionine is partially destroyed on acid hydrolysis.

as zero; for the chromatographic RF the solvent front is defined as unity and the origin as zero.

The new peptide from I.3 (family L) was digested with thermolysin which hydrolyses α TpIX in the positions shown in Fig. 6. The thermolysin fingerprint (Fig. 7) had a new positively charged histidine-containing peptide which was α 62-65 with a histidine replacing the aspartic acid at α 64. In digests of α^A TpIX the peptide α 62-65 runs on top of α 83-85 but in the abnormal α^Q TpIX only the peptide α 83-85 is present in this position on the fingerprint. The only other peptide containing aspartic acid, α 73-79, was present in its normal position (Table VI). The mutation in this haemoglobin must therefore be α 64 aspartic acid $\rightarrow$ histidine.

In the case of I.1 (family A) a peptic digest of α^Q TpIX was attempted. The results were unsatisfactory but it appeared that the mutation Asp $\rightarrow$ His was probably at position α 64 (Table VII). The fingerprint (Fig. 8) showed a new histidine-containing spot which is not present in peptic digests of α^A TpIX. This spot was found to contain the peptide α 62-70 without aspartic acid.

Discussion

Hb Q India does not cause any haematological disorders, perhaps because the residue involved, α 64 (E13), is on the surface of the haemoglobin tetramer. Charge changes at such positions do not usually affect the properties of the haemoglobin. The only other mutant so far found at this position is Hb Aida (α 64 Asp $\rightarrow$ Asn) (Ramot, Kinderlerer, and Lehmann, 1972). This is also asymptomatic, and is present in a proportion of 27%.

The proportion of Hb Q varies from 8-20%, the proportions in the cases of β -thalassaemia minor tending to be lower than that in the normal case

FIG. 6. Amino-acid sequence of α^A TpIX. $\uparrow$ = hydrolysis by pepsin; $\uparrow$ = hydrolysis by thermolysin.

TABLE VI
AMINO-ACID ANALYSIS OF THERMOLYSIN DIGEST IN
I.3 (FAMILY L)

Amino Acid	62-65		73-79		83-85	
	Ratio	Expected	Ratio	Expected	Ratio	Expected
Asp	—	1	2.94	3	1.07	1
Ser	0.33	—	—	—	0.88	1
Pro	—	—	0.91	1	—	—
Gly	0.31	—	—	—	—	—
Ala	1.96	2	1.08	1	—	—
Val	1.00	1	1.06	1	—	—
Met	—	—	0.70	1	—	—
Leu	—	—	—	—	1.05	1
His	1.05	—	—	—	—	—
Yield (n moles)	35		29		79

FIG. 7. Comparison of the fingerprints of thermolysin peptides of α^A TpIX and α^Q TpIX (α 62-90). (For details of interpretation see text.) * = Point of application.

TABLE VII
AMINO-ACID ANALYSIS OF PEPTIC DIGEST IN
I.1 (FAMILY A)

Amino Acid	62-69		84-86	
	Ratio	Expected	Ratio	Expected
Asp	1.07	2	0.93	1
Thr	0.97	1	—	—
Ser	—	—	1.28	1
Ala	3.08	3	—	—
Val	0.97	1	—	—
Leu	0.91	1	0.85	1
His	0.30	—	—	—
Yield (n moles)	9.1		2.2

FIG. 8. Comparison of the fingerprints of peptic peptides of α^A TpIX and α^Q TpIX (α 62-90). (For details of interpretation see text.) * = Point of application.

(Tables I and II). The proportion of Hb $\alpha_2\gamma_2$ compared to $\alpha_2\gamma_2$ in II.4 (family L) is 6.9%, even lower than in the trait. It is possible that when there is a shortage of non α -chains the α^A chains are preferentially incorporated into haemoglobin while the precipitated α -chains contain a greater proportion of α^Q . However, without more data on the proportions of Hb Q India in non-thalassaemic people, it is difficult to judge whether the lowering of the proportion of Hb Q in cases of β -thalassaemia minor is significant.

Summary

A new haemoglobin, Hb Q India, α 64 Asp $\rightarrow$ His has been discovered. It was found in association both with β -thalassaemia major and β -thalassaemia minor. It seems that the proportion of this haemoglobin is lower when β -thalassaemia is present.

Our thanks are due to Dr L. D. Sanghvi MSc, PhD (Columbia), Head, Epidemiology Division and Dean, Cancer Research Institute, Tata Memorial Centre, Parel, Bombay 12, for his continued encouragement in this work and to the Superintendent, Jerbai Wadia Hospital for Children, for permission to publish these data.

REFERENCES

Blackwell, R. Q. and Liu, C. S. (1970). Haemoglobin G Taichung: α 74 Asp $\rightarrow$ His. *Biochimica et Biophysica Acta*, **200**, 70-75.
Chatterjee, J. B. (1966). Haemoglobinopathies, glucose-6-phosphate dehydrogenase deficiency and allied problems in the Indian sub continent. *Bulletin of the World Health Organization*, **35**, 837-856.
Chouhan, D. M., Sharma, R. S., and Parekh, J. G. (1970). Alpha-thalassaemia in India. *Journal of the Indian Medical Association*, **54**, 364-367.
Cradock-Watson, J. E., Fenton, J. C. B., and Lehmann, H. (1959). Tris buffer for the demonstration of haemoglobin A₂ by paper electrophoresis. *Journal of Clinical Pathology*, **12**, 372-373.
Dacie, J. V. and Lewis, S. M. (1968). *Practical Haematology*, 4th ed. Churchill, London.
Huisman, T. H. J. and Dozy, A. M. (1965). Studies on the heterogeneity of hemoglobin. IX. The use of Tris (hydroxymethyl) aminomethane HCl buffers in the anion-exchange chromatography of hemoglobins. *Journal of Chromatography*, **19**, 160-169.

- Lorkin, P. A., Charlesworth, D., Lehmann, H., Rahbar, S., Tuchinda, S., and Lie Injo Luan Eng. (1970). Two haemoglobins Q, $\alpha 74$ (EF3) and $\alpha 75$ (EF4) aspartic acid $\rightarrow$ histidine. *British Journal of Haematology*, **19**, 117-125.
- Marengo-Rowe, A. J. (1965). Rapid electrophoresis and quantitation of haemoglobins on cellulose acetate. *Journal of Clinical Pathology*, **18**, 790-792.
- Pootrakul, S. and Dixon, G. H. (1970). Hemoglobin Mahidol: A New hemoglobin α -chain mutant. *Canadian Journal of Biochemistry*, **48**, 1066-1078.
- Poulik, M. D. (1959). Starch gel electrophoresis in a discontinuous system of buffers. *Nature*, **180**, 1477-1479.
- Ramot, B., Kinderlerer, J., and Lehmann, H. (1972). Haemoglobin Aida $\alpha 64$ aspartic acid $\rightarrow$ asparagine. (In preparation.)
- Sagnet, H., Morineaud, J. P., Delprat, J., Revil, H., Thomas, J., Philibert, H., and Marfart, Y. (1968). Une hémoglobinoïse rare: l'association hémoglobine Q- α -thalassémie. *Médecine Tropicale*, **28**, 133-138.
- Sanghvi, L. D., Sukumaran, P. K., and Lehmann, H. (1958). Haemoglobin J trait in two Indian women associated with thalassaemia in one. *British Medical Journal*, **2**, 828-830.
- Sick, K., Beale, D., Irvine, D., Lehmann, H., Goodall, P. T., and McDougall, S. (1967). Haemoglobin G Copenhagen and Haemoglobin J Cambridge. Two new β -chain variants of Haemoglobin A. *Biochimica et Biophysica Acta*, **140**, 231-242.
- Singer, K., Chernoff, A. I., and Singer, L. (1951). Studies on abnormal hemoglobins. I. Their demonstration in sickle cell anemia and other hematologic disorders by means of alkali denaturation. *Blood*, **6**, 413-428.
- Sukumaran, P. K., Wiltshire, B. G., and Lehmann, H. (1971). A new haemoglobin $\alpha 64$ aspartic acid $\rightarrow$ histidine observed in two Sindhi families. In *Abstracts and Proceedings of the 2nd Meeting of the Asian-Pacific Division of the International Society of Haematology*, p. 14. Melbourne.
- Vella, F., Wells, R. H. C., Ager, J. A. M., and Lehmann, H. (1958). A haemoglobinopathy involving Haemoglobin H and a new (Q) Haemoglobin. *British Medical Journal*, **1**, 752-755.