A search for linkage in cystic fibrosis

MARY C. GOODCHILD, J. H. EDWARDS, KAREN P. GLENN, CLARE GRINDEY,
R. HARRIS, PAULINE MACKINTOSH, and J. WENTZEL

From the Institute of Child Health, University of Birmingham; Infant Development Unit, Maternity Hospital, Birmingham; Department of Medical Genetics, University of Manchester; and the Regional Blood Transfusion Service, Birmingham

Summary. Linkage between the locus for cystic fibrosis and other genetic markers was explored in 40 families from Birmingham and 20 from Manchester. No strong evidence was found for linkage with any of the markers examined. There was evidence against close linkage with ABO, HLA, and Rh.

Cystic fibrosis, known originally as cystic fibrosis of the pancreas, is the commonest recessive lethal disease in most Caucasian populations. Death usually results from chronic lung disease, and few cases survive beyond young adulthood. In addition, fertility is greatly impaired in the male and probably impaired in the female. The basic defect is unknown and the evidence for its recessive nature is based entirely on its incidence in sibs (Carter, 1952), in other relatives also (Danks, Allan, and Anderson, 1965), and on its varied incidence in different racial groups, even when living in the same community (Wright and Morton, 1968).

Linkage between two loci means that they are sited on the same chromosome. Establishment of a linkage between the locus, or one of the loci, for cystic fibrosis, and a locus determining some other phenotype, would provide information for chromosome mapping. Such data could indicate also whether cystic fibrosis is determined at one locus or at several loci; in addition, definition of the position of a locus close to a cystic fibrosis locus could permit prenatal diagnosis in some families.

A previous linkage study in cystic fibrosis (Steinberg *et al*, 1956; Steinberg and Morton, 1956), did not reveal strong evidence for linkage between cystic fibrosis and any of the eight blood group loci; however, Steinberg *et al* (1956) stated that 'the data for the P locus indicate that it may be profitably tested further'.

The present work concerns an attempt to define a locus for cystic fibrosis by means of a linkage study which included other markers, including the normal

chromosomal variants often evident with modern staining techniques.

Patients and methods

This study was based on the families of 40 cystic fibrosis children attending clinics at the Birmingham Children's Hospital and of 20 more cystic fibrosis children attending clinics at the Royal Manchester Children's Hospital, West Park Hospital Macclesfield, and the Blackburn Royal Infirmary. Each family included at least two children, of which at least one was affected, and two parents. Whenever possible grandparents were included.

A linkage study involving a recessive condition, such as cystic fibrosis, gives limited information if the heterozygous grandparent and sib cannot be identified. Unfortunately, so far, there is no reliable method for the detection of the carrier in cystic fibrosis.

Blood samples were taken, with parental permission, into lithium heparin tubes. In Birmingham, lymphocyte separations for HLA typing were made using a Ficoll–Triosil density gradient technique, cell samples were washed to remove platelets, and typing was carried out in Falcon plastic trays using a modification of the technique of Terasaki and McClelland (1964). In Manchester, samples were taken into Boots' preservative-free heparin and typed on Searle glass plates. At both centres, tests included six specificities at the first locus and eight at the second. This was sufficient for haplotyping all these families.

Lods, an acronym for 'logarithms of the odds ratios', provide a simple additive measure of the evidence for linkage at any defined recombination fraction (θ). Values within the range -1.0 to $+2.0$ give very limited evidence either against or for linkage at any defined value of θ . Lods were computed by a generalization of Morton's method (Morton, 1955; Edwards, 1972; Chautard-Freire-Maia, 1974). Data published previously from

Boston by Steinberg *et al* (1956) were analysed using our computer programme, as were data from eight families from Leeds, kindly supplied by Dr M. d'A. Crawford. Data from the latter families were examined for linkage with P, but on analysis the lod from these families almost cancelled out, providing virtually no information (lods -0.01 at both $\theta=0.1$ and $\theta=0.3$).

Results

Total lods for the informative families from Birmingham and Manchester are shown together in Table I, and lods derived from the published data of Steinberg *et al* (1956) in Table II. Total lods from

TABLE I

LODS FROM BIRMINGHAM AND MANCHESTER FAMILIES BETWEEN VARIOUS LOCI AND CYSTIC FIBROSIS, ASSUMING A SINGLE CF LOCUS AT RECOMBINATION FRACTIONS OF 0.1 AND 0.3

	$\theta = 0.1$	$\theta = 0.3$
ABO	-2.35	-0.37
MNSs	-0.33	-0.02
P	-0.05	0.18
Rh	-2.01	-0.08
Haptoglobin	0.07	0.04
Acid phosphatase 1	-1.02	-0.14
6 phosphogluconate dehydrogenase	0.00	0.00
Phosphoglucomutase 1	-0.84	-0.08
Adenylate kinase	-0.26	-0.02
Glutamic pyruvic transaminase	-0.89	-0.05
Adenosine deaminase	0.08	0.03
HLA	-2.73	0.16

TABLE II

LODS COMPUTED FROM DATA BY STEINBERG *et al* (1956)

	$\theta = 0.1$	$\theta = 0.3$
ABO	-1.87	-0.16
MNSs	-0.30	0.09
P	-0.04	0.05
Rh	-2.13	-0.35
Kell	-0.03	-0.01
Fy	-0.27	-0.07
Jk	-0.05	-0.01

TABLE III

TOTAL LODS FROM DATA FROM BIRMINGHAM, MANCHESTER, BOSTON, AND LEEDS

	$\theta = 0.1$	$\theta = 0.3$
ABO	-4.22	-0.53
MNSs	-0.63	0.07
P	-0.10	0.22
Rh	-4.14	-0.43
Kell	-0.03	-0.01
Fy	-0.27	-0.07
Jk	-0.05	-0.01
Haptoglobin	0.07	0.04
Acid phosphatase 1	-1.02	-0.14
6 phosphogluconate dehydrogenase	0.00	0.00
Phosphoglucomutase 1	-0.84	-0.08
Adenylate kinase	-0.26	-0.02
Glutamic pyruvic transaminase	-0.89	-0.05
Adenosine deaminase	0.08	0.03
HLA	-2.73	0.16

FIG. Family W, showing the segregation of the sixth chromosomes. There is no evident variation within individuals. The slight differences between individuals are the result of the chromosomes being at slightly different stages of division. It is not even possible to define which of the pair of parental chromosomes has been passed on to any child.

all sources are given in Table III. There is no strong evidence for close linkage between the loci for CF and any of the markers examined.

Chromosomal banding (illustrated in detail for chromosome six in Family W., Fig. 1) was uninformative due to a failure to detect any clearly segregating marker chromosomes in any of the 14 families studied. Particular attention was paid to chromosome 6, since the first analyses of our data were suggestive of linkage of cystic fibrosis to both HLA and P, and there is evidence that the HLA and P loci are linked (Fellous *et al*, 1971).

Discussion

The position of the locus, or if several, the major locus, carrying the allele or alleles determining cystic fibrosis remains unknown. Information provided in this study gives no strong evidence for

linkage with any of the markers investigated. Nevertheless, the fairly high negative lods for ABO and cystic fibrosis (-4.22 , $\theta=0.1$, Table III), for HLA and cystic fibrosis (-2.73 , $\theta=0.1$, Table III) and for Rh cystic fibrosis (-4.14 , $\theta=0.1$, Table III) exclude close linkages. The lods for linkage between P and cystic fibrosis were positive in the data of Steinberg *et al* (1956) and in our data at $\theta=0.3$. However the total lod for linkage with P of 0.22 at $\theta=0.3$ (Table III) is very weak evidence for loose linkage between P and cystic fibrosis. The P locus is probably on chromosome 6 (Fellous *et al*, 1971), though not near to the HLA locus (Edwards *et al*, 1972; Lamm *et al*, 1975).

The absence of any strong evidence of linkage in a study of 60 families with a recessive disorder is hardly surprising, since the chance of the locus being sufficiently close to any of the markers tested is small, and probably less than 1 in a 100. Though many recessive disorders may be the result of the locus being lost by a chromosomal deficiency, the likelihood of such a deficiency being within the range of optical microscopy is low. The chromosomal studies on parents probably exclude any deficiency of more than half a band or interband width. Very close linkage between a major cystic fibrosis locus and HLA had been excluded already by absence of association (Polymenidis, Ludwig, and Götz, 1973; Goodchild, Nelson, and Anderson, 1973).

We thank Dr M. d'A. Crawford for allowing us to analyse and use his data, the Medical Research Council Human Biochemical Genetics Unit for typing the biochemical markers, and Professor J. J. van Rood and Dr

Q. Lansberger for computing facilities at the University of Leiden. In the Manchester area, we are grateful for the co-operation of Dr N. M. Mann, Dr J. D. Allan, Dr P. D. Moss, and Mrs E. Quinn for skilled field work. M.C.G. is indebted to the Cystic Fibrosis Research Trust for financial support and C.G. to the Centenary Research Fund.

REFERENCES

- Carter, C. O. (1952). *Fibrocystic Disease of the Pancreas. A Congenital Disorder of Mucus Production—Mucosis*, pp. 50–64. Ed. by M. Bodian. Heinemann, London.
- Chautard-Freire-Maia, E. A. (1974). Linkage relationships between 22 autosomal markers. *Annals of Human Genetics*, **38**, 191–198.
- Danks, D. M., Allan, J., and Anderson, C. M. (1965). A genetic study of fibrocystic disease of the pancreas. *Annals of Human Genetics*, **28**, 323–356.
- Edwards, J. H. (1972). A marker algebra. *Clinical Genetics*, **3**, 371–380.
- Edwards, J. H., Allen, F. H., Glenn, K. P., Lamm, L. U., and Robson, E. B. (1972). The linkage relationships of HL-A. *Histocompatibility Testing*, **3**, 746–751.
- Fellous, M., Billardon, C., Dausset, J., and Frézal, J. (1971). Linkage probable entre les locus 'HL-A' et 'P'. *Comptes Rendus Académie Scientifique (Paris)*, **272**, 3356–3359.
- Goodchild, M. C., Nelson, R., and Anderson, C. M. (1973). Cystic fibrosis and coeliac disease: coexistence in two children. *Archives of Disease in Childhood*, **48**, 684–691.
- Lamm, L. U., Thorsen, I.-L., Petersen, G. B., Jørgensen, J., Henningsen, K., Bech, B., and Kissmeyer-Nielsen, F. (1975). Data on the HL-A linkage group. *Annals of Human Genetics*, **38**, 383–390.
- Morton, N. E. (1955). Sequential tests for the detection of linkage. *American Journal of Human Genetics*, **7**, 277–318.
- Polymenidis, Z., Ludwig, H., and Götz, M. (1973). Cystic fibrosis and HL-A antigens. *Lancet*, **2**, 1452.
- Steinberg, A. G. and Morton, N. E. (1956). Sequential test for linkage between cystic fibrosis of the pancreas and the MNS locus. *American Journal of Human Genetics*, **8**, 177–189.
- Steinberg, A. G., Shwachman, H., Allen, F. H., and Dooley, R. R. (1956). Linkage studies with cystic fibrosis of the pancreas. *American Journal of Human Genetics*, **8**, 162–176.
- Terasaki, P. I. and McClelland, J. D. (1964). Microdroplet assay of human serum cytotoxins. *Nature (London)*, **204**, 998–1000.
- Wright, S. W. and Morton, N. E. (1968). Genetic studies on cystic fibrosis in Hawaii. *American Journal of Human Genetics*, **20**, 157–169.