

susceptibility to a particular disorder. The data presented here on PGM and tuberculosis lead one to suspect that discovery of true associations of gene markers and disease must await establishment of more refined techniques of gene identification, such as that of isoelectric focusing used here or of DNA probes currently in process of development.

The authors are grateful to Professor D F Roberts for his suggestions during this work and his perusal of the manuscript.

#### References

- <sup>1</sup> Selby R, Barnard JM, Buehler SK, Crumley J, Larsen B, Marshall WH. Tuberculosis associated with HLA-B8, BfS in a Newfoundland community study. *Tissue Antigens* 1978;**11**:403-8.
- <sup>2</sup> Khomenko AG, Averbach MM, Kalankhodzaev AA, Malenko AF, Chukanova VP, Moroz AM. Distribution of HLA antigens in tuberculosis patients. *Ter Arkh* 1980;**53**:135-8.
- <sup>3</sup> Mourant AE, Kopec AA, Domaniewska-Sobczak K. *Blood groups and disease*. Oxford: Oxford University Press, 1978.
- <sup>4</sup> Kooptzoff O, Walsh RJ. Blood groups and diseases: pulmonary tuberculosis. *Aust Ann Med* 1957;**6**:53-5.
- <sup>5</sup> Milunicova A, Dominec M. Blood characteristics and PTC test in young patients suffering from pulmonary tuberculosis and diabetes mellitus. *Rozhl Tuberk* 1966;**26**:545-50.
- <sup>6</sup> Hever O. Relations entre les phénotype d'haptoglobine dans diverses maladies. *Presse Med* 1969;**77**:1081-2.
- <sup>7</sup> Sidhu LS, Singh J, Bhatnagar DP, Pahuja JK. Association of pulmonary tuberculosis with ABO and Rh(D) blood groups. In: Sanghvi LD *et al*, eds. *Population genetics in India*. India: Orient Longman, 1974;135-40.
- <sup>8</sup> Karim AKMB, Elfellah MS, Evans DAP. Human acetylase polymorphism: estimation of allele frequency in Libya and details of global distribution. *J Med Genet* 1981;**18**:325-30.
- <sup>9</sup> Constans J, Viau M. Group-specific component: evidence of two subtypes of Gc<sup>1</sup> gene. *Science* 1977;**198**:1070-1.
- <sup>10</sup> Bark JE, Harris MJ, Firth M. Typing of the common phosphoglucomutase variants using isoelectric focusing. A new interpretation of phosphoglucomutase system. *J Forensic Sci Soc* 1976;**16**:115-20.
- <sup>11</sup> Papiha SS. Isoelectric focusing in the detection of genetic variation of enzyme and proteins: method and gene frequencies of the group-specific component (Gc) in the Gaddi tribe of Chamba (Himachal Pradesh). *J Indian Anthropol Soc* 1981;**16**:75-87.
- <sup>12</sup> Spencer N, Hopkinson DA, Harris H. Phosphoglucomutase polymorphism in man. *Nature* 1964;**204**:742-5.
- <sup>13</sup> Papiha SS, Roberts DF, White I, Chahal SMS, Asefi AJ. Population genetics of the group-specific component and phosphoglucomutase (PGM<sub>1</sub>) studies by isoelectric focusing. *Am J Phys Anthropol* 1982;**59**:1-7.
- <sup>14</sup> Woolf B. On estimating the relation between blood group and disease. *Ann Hum Genet* 1955;**19**:251-3.
- <sup>15</sup> Haldane JBS. The estimation and significance of the logarithm of a ratio of frequencies. *Ann Hum Genet* 1955;**20**:309-11.

Correspondence and requests for reprints to Dr S S Papiha, Department of Human Genetics, 19 Claremont Place, Newcastle upon Tyne NE2 4AA.

#### Correction

In the paper by Price and Wilson on "Dissection of the aorta in Turner's syndrome" which was published in the February 1983 issue of the Journal (20:61-3), the last sentence of the first column on the first page should have read: "They all had the clinical features of TS and the following chromosome abnormalities: 45,X (83); 46,X,i(Xq) (12); 45,X/46,X,i(Xq) (19); 45,X/46,XY (13); 46,X,del(X)(p) (5); 45,X/46,X,+r (7); 45,X/46,X,mar(X) (4); 45,X/46,X,r(X) (5); other mosaics (9)".