- ³ Gray JE, Syrett JE, Ritchie KM, Elliot WD. An interstitial translocation: chromosome No 1p to 4q. *Lancet* 1972;ii:92-3.
- ⁴ Therkelsen AJ, Hultén M, Jouessan J, Lundsten J, Christensen NC, Iversen T. Presumptive direct insertion within chromosome 2 in man. *Ann Hum Genet* 1973;36:367-73.
- ⁵ Toomey KE, Mohandas T, Sparkes RS, Kabach MM, Rimoin DL. Segregation of an insertional chromosome rearrangement in 3 generations. *J Med Genet* 1978;15:382-7.
- ⁶ Grace E, Sutherland GR, Bain AD. Familial insertional translocation. *Lancet* 1972;ii:231.
- ⁷ Rethoré MO, Lejeune J, Carpentier S. Trisomie pour la partie distale du bras court du chromosome 3 chez trois germains. Premier exemple d'insertion chromosomique: ins (7;3)(q31;p21p26). *Ann Genet (Paris)* 1972;15:159-65.
- ⁸ Shapiro LR, Warburton D. Interstitial translocation in man. *Lancet* 1972;ii:712-3.
- ⁹ Berger R, Touati G, Derre J. "Cri du chat" syndrome with maternal insertional translocation. *Clin Genet* 1974;5:428-32.
- ¹⁰ Chudley AE, Bauder F, Ray M, McAlpine PJ, Pena SDJ, Hamerton JL. Familial mental retardation in a family with an inherited chromosome rearrangement. *J Med Genet* 1974;11:353-63.
- ¹¹ Fitzgerald MG. Complex five-break rearrangement. *Clin Genet* 1974;5:62-7.
- ¹² O'Donnell JJ, Hall BD, Conte FA, Romanowski JC, Epstein CJ. Down syndrome: localisation of locus to distal portion of long arm of chromosome 21. *Pediatr Res* 1975;9:315.
- ¹³ Sparkes RS, Salter WJ, Blaker RG, Muller HM. Insertional translocation into the X chromosome of a 46,XY male. *Clin Genet* 1977;12:114-8.
- ¹⁴ Pan SG, Fatora SR, Sorg R, Garver KL, Steele MW. Meiotic consequences of an intrachromosomal insertion of chromosome No 1: a family pedigree. *Clin Genet* 1977;12:303-13.
- ¹⁵ Grass FS, Schwartz RP, Deal JO, Parke JC Jr. Gonadal dysgenesis, ultra-X chromosome insertion, and possible position effect in an otherwise normal female. *Clin Genet* 1981;20:28-35.
- ¹⁶ Borgaonker DS, Shaffer R, Reed WC, Jackson LG. *Repository of chromosomal variants and anomalies in man. An international registry of abnormal karyotypes*. Philadelphia: Thomas Jefferson University, 1981.
- ¹⁷ Varley JM, Gorden J, Hultén M. Familial reciprocal translocation t(9;13)(p11;p12) investigated by silver staining and in situ hybridisation. *Hum Genet* 1981;59:422-8.
- ¹⁸ Spowart G. Reassessment of presumed Y/22 and Y/15 translocations in man using a new technique. *Cytogenet Cell Genet* 1979;23:90-4.
- ¹⁹ Howell WM, Black DA. A rapid technique for producing silver-stained nucleolus organizer regions and trypsin-giensa bands on human chromosomes. *Hum Genet* 1978;43:53-6.
- ²⁰ Denton TE, Brooke WR, Howell WM. A technique for the simultaneous staining of both nucleolar organiser regions and kinetochores of human chromosomes with silver. *Stain Technol* 1977;52:311-3.
- ²¹ Williams JD, Summitt RL, Martens R, Kimbrell RA. Familial Down syndrome due to t(10;21) translocation. Evidence that the Down phenotype is related to trisomy of a specific segment of chromosome 21. *Am J Hum Genet* 1975;27:478-85.
- ²² Moses MJ. The synaptonemal complex and meiosis. In: Sparkes RS, ed. *Molecular human cytogenetics*. New York: Academic Press, 1977:101-25.
- ²³ Miklos GL, Willcocks DA, Baverstock PR. Restriction endonuclease and molecular analysis of three rat genomes with special reference to chromosome rearrangement and speciation problems. *Chromosoma* 1980;76:339-63.
- ²⁴ Miklos GL, John B. Heterochromatin and satellite DNA in man: properties and prospects. *Am J Hum Genet* 1979;31:264-80.
- ²⁵ Aurias A, Prieur M, Dutrillaux B. Systematic analysis of 95 reciprocal translocations of autosomes. *Hum Genet* 1978;45:259-82.

Correspondence and requests for reprints to Dr Jessie L Watt, Department of Genetics, University of Aberdeen, Medical School Building, Foresterhill, Aberdeen AB9 2ZD.

Partial trisomy 16 as a result of familial 16;20 translocation

E V DAVISON* AND J R BEESLEY†

*Department of Human Genetics, 19 Claremont Place, Newcastle upon Tyne NE2 4AA; and †Department of Paediatrics, Newcastle General Hospital, Westgate Road, Newcastle upon Tyne NE4 6BE.

SUMMARY Although trisomy 16 is well recognised in spontaneous abortuses,¹ it is infrequent in livebirths and there is little information about the clinical effects.²⁻³ We report two sibs with partial trisomy 16q resulting in infant death. Both children were severely growth retarded with small elfin faces, prominent foreheads, low set ears, abnormal external genitalia, and intractable diarrhoea.

Received for publication 17 November 1983.
Accepted for publication 6 February 1984.

Case reports

The first child of healthy unrelated young parents was born after an uncomplicated pregnancy by normal delivery at term. The child weighed 2125 g at birth and her head circumference was 31.5 cm. She had distinctive facies, with a prominent forehead, small chin, and low set ears. She had bilateral fusion of the middle and fourth toes and very prominent labia majora. On the third day she developed watery diarrhoea which necessitated intravenous fluid therapy. Her condition improved briefly although

the undiagnosed diarrhoea persisted until the fifth week of life when she suddenly deteriorated and died. Necropsy showed a small VSD but no other macroscopic congenital anomalies and no histology was done.

The second pregnancy resulted in a premature small for dates but healthy female. The third pregnancy resulted in a 10 week miscarriage about which there is no information. A fourth uncomplicated pregnancy resulted in a normal delivery at 36 weeks' gestation of a male infant. He weighed 1875 g at birth, length 46 cm, and head circumference 31.7 cm. Gestational assessment confirmed his maturity. External congenital anomalies included a small elfin face with a prominent forehead and low set ears, giving him an almost identical appearance to the child who had died 8 years earlier (fig 1). He had severe hypospadias with a bifid scrotum, dystrophic nails on all digits, and rockerbottom feet. A soft systolic murmur compatible with a VSD was audible after the third day of life. At 48 hours he became pale and hypotonic and developed frequent watery motions but no vomiting. He was never hypoglycaemic and there was no evidence of infection. Faecal fluid contained 2% reducing substances which persisted despite giving only dextrose feeds. Serum electrolytes were normal although initially the urea was raised (11.3 mmol/l) and he had a moderate metabolic acidosis. He was parenterally fed but the diarrhoea persisted despite subsequent trials of various infant formulae and loperamide. Serum electrolytes remained normal with the exception of moderate hypokalaemia which was

easily corrected. No bacteria or viruses were isolated from stool specimens. A barium follow through showed a normal bowel and a chest x-ray showed a normal heart with slight prominence of the lung vessels. The skull x-ray showed defective ossification of the parietal and frontal bones. Analyses of blood and urine showed no evidence of a metabolic disorder. Diarrhoea persisted until he died at 18 days. Permission for necropsy was refused.

CYTOGENETIC FINDINGS

The karyotypes of both abnormal infants were 46,XX or XY,-20,+der(20),t(16;20)(q13;p13)pat. The breakpoint of the long arm of chromosome 16 appeared to be at band q13 just below the heterochromatic region and on chromosome 20 at band p13 (fig 2).

FAMILY STUDIES

Cytogenetic analysis performed in 1976 after the birth of the first infant identified the translocation. The parents at that time were both 16 years of age, the implications of this translocation were not clearly understood, family studies were not attempted, and subsequent pregnancies were not monitored by amniocentesis. After the birth of the second child with the unbalanced translocation, both parents had matured considerably and further family studies were

FIG 1 Physical appearance of second abnormal child.

FIG 2 Partial karyotypes of (a) the probands, and (b) the father. (c) Diagrammatic representation of breakpoints.

initiated. The translocation was found in other members of the family, although in no other part of the family was there any history of obstetric failure or abnormal livebirths. All the relevant members of the family have now received genetic counselling. The original family have recently completed a further pregnancy which was monitored by amniocentesis, and the infant is a carrier of the balanced form of the translocation.

Discussion

Yunis *et al*² reported the first case of a liveborn infant with a partial trisomy 16q-. This trisomy is almost identical to ours although the case report of Yunis *et al*² does not state at what age the infant died. Since then, data have been accumulating on the effect of partial monosomy and partial trisomy of chromosome 16.

Most of the chromosomal anomalies are the result of a balanced translocation of parental derivation, which also involves either monosomy or trisomy of other chromosomes in the karyotype. Although early reports of full trisomy 16 do exist⁴⁻⁶ they predated banding techniques and cannot be included in an attempt to define a syndrome. Full trisomy 16 is well reported in spontaneous abortion and generally results in disorganised embryos with very little evidence of development. Limited post-natal survival is possible with both partial trisomy 16p and 16q.^{3,7}

Roberts and Duckett³ reviewed published reports and included three cases of partial trisomy 16p, three cases of partial trisomy 16q, and one case with partial trisomy 16p and 16q. The most common features described in both of these trisomies were digital deformities, low birth weight, small malformed facies, which included low set abnormal ears, small palpebral fissures, and hypoplastic mandible. These anomalies are also consistent with anomalies found in many other chromosomal aberrations. Ridler and McKeown⁷ reported trisomy 16q arising from a maternal 15p;16q translocation. The breakpoint on the 16 was at q11 and the involvement of chromosomal material was similar to that in the two cases reported here. The anomalies found included a small and emaciated infant with a small elongated head, a narrow face, and a very low hair line; the child was described as having leprechaun-like facies. The skin was wrinkled, the limbs were

abnormally flexed, and the hands showed a transverse palmar crease on the right and clinodactyly of both fifth digits. The infant died after 12 days.

The two cases presented here suggest an association between the intractable diarrhoea, which was the apparent cause of death in both, and partial trisomy 16. There is no report of this in any of the other cases.

Although there are perhaps still insufficient data to attempt to define a syndrome for partial or full trisomy 16, the most notable characteristics are intrauterine growth retardation, particular abnormalities of the face with downward slanting palpebral fissures, abnormal nose and ears, micrognathia, flexion deformities of the limbs and digits, and heart defects. Although many of these anomalies are found in association with other chromosome aberrations it is the particular facial appearance which suggests the trisomy for this chromosome.

The authors would like to thank Dr W D Elliott and Mr J Wagget for permission to report their patient, and Dr A R Boon for her involvement in the genetic counselling.

References

- 1 Creasy MR, Crolla JA, Alberman ED. A cytogenetic study of human spontaneous abortions using banding techniques. *Hum Genet* 1976;**31**:177-96.
- 2 Yunis E, Gonzalez JT, Torres de Caballero OM. Partial trisomy 16q-. *Hum Genet* 1977;**38**:347-50.
- 3 Roberts SH, Duckett DP. Trisomy 16q in a liveborn infant and a review of partial and full trisomy 16. *J Med Genet* 1978;**15**:375-81.
- 4 Schmidt BH, Becak W, Becak ML, *et al*. Metaphyseal dystosis. Review of literature; study of a case with cytogenetic analysis. *J Pediatr* 1963;**63**:106-12.
- 5 Backus JA, Darien GG. Group E triple cell line mosaicism with Sprengel's deformity and other asymmetrical congenital anomalies. Report of a case. *Am J Clin Pathol* 1968;**49**:527-34.
- 6 Arakaki DT, Waxman SH. Trisomy 16 in a mosaic carrier father and his aborted fetus. *J Med Genet* 1969;**6**:85-8.
- 7 Ridler MAC, McKeown JA. Trisomy 16q arising from a maternal 15p;16q translocation. *J Med Genet* 1979;**16**:317-20.

Correspondence and requests for reprints to Dr E V Davison, Department of Human Genetics, University of Newcastle upon Tyne, 19 Claremont Place, Newcastle upon Tyne NE2 4AA.