Short report

High frequency of two mutations in codon 778 in exon 8 of the ATP7B gene in Taiwanese families with Wilson disease

Lee-Ming Chuang, Huey-Peir Wu, Mei-Huei Jang, Tso-Ren Wang, Whey-Chen Sue, Boniface J Lin, Diane W Cox, Tong-Yuan Tai

Abstract

The gene for Wilson disease (WD) has been cloned as a P type copper transporter gene (ATP7B). To elucidate the possible genetic mechanism for the diversity of clinical manifestations, we characterised 22 Taiwanese families with WD by microsatellite haplotyping of close DNA markers D13S314-D13S301-D13S316. We also screened for mutations of codon 778 in the transmembrane region. There were at least 15 haplotypes within eight broad subgroups observed among 44 WD chromosomes. Among the 22 unrelated patients, we found that six patients (27%) carried a codon 778 mutation. Nucleotide sequence analysis showed there were two different mutations: the previously reported Arg778Leu mutation in four patients and Arg778Gln, a new mutation, in two patients. The two different mutations of the same codon occurred in two distinct microsatellite haplotypes.

(J Med Genet 1996;33:521-523)

Key words: ATP7B gene; haplotypes; Taiwan.

Wilson disease (WD) is a rare autosomal recessive disorder of copper transport.¹ Genetic studies showed linkage of the disease locus to the esterase D and retinoblastoma genes^{2,3} and to DNA markers in the 13q14 region.⁴ Microsatellite markers closely flanking the locus have been reported.⁵⁻⁷ Recently, a putative copper transporting P type ATPase (ATP7B) gene has been cloned and is defective in WD.^{8,9} The gene shows remarkable sequence homology to the gene for Menkes disease (ATP7A), another disorder of copper transport.¹⁰⁻¹² The proteins encoded by these genes belong to the P type ATPase family of cation transporters.¹³ Subsequently, more than 25 mutations, including small insertions and deletions, missense, nonsense, and splice site mutations have been described.^{8,9,14} The prevalence of mutations in

the Chinese population has not been extensively studied, although an Arg778Leu* missense mutation has been reported in two Chinese pedigrees from Hong Kong.¹⁴ We have shown linkage of WD to the chromosome 13q14 region, as reported for other populations, suggesting the monogenic nature of the disease.¹⁵ Since the variability of clinical manifestation in the Taiwanese pedigrees with WD^{16,17} might possibly be explained by genetic heterogeneity,^{14,18} we examined the microsatellite haplotypes and screened for the reported Chinese mutation in Taiwanese families. DNA was isolated from leucocytes from 25 patients in 22 families. The diagnosis was established by the detection of Kayser-Fleischer rings or abnormal liver function tests in the presence of low caeruloplasmin levels.¹⁵ Microsatellite (CA) dinucleotide repeats at loci D13S301, D13S314, D13S316, and D13S133 were analysed according to previous reports.^{6,7} Screening for the previously described codon 778 mutation in exon 8 of the ATP7B gene was performed by polymerase chain reaction (PCR) based restriction fragment length polymorphism (RFLP) analysis.¹⁴ The mutations were confirmed by direct nucleotide sequence analysis with an AmpliCycle™ sequencing kit (Perkin Elmer, NJ). The haplotypes carried on the WD chromosomes are listed in the table in the order of prevalence. The A haplotype was significantly higher in patients with WD in Taiwan ($p < 0.015$, χ^2 test). Among the 25 patients tested, eight patients in six (27.3%) pedigrees showed an absence of the *MspI* restriction site in the amplified fragments, suggesting a mutation in codon 778. The nucleotide change was verified by direct sequence analysis. Interestingly, two different mutations were found. Five patients carried a G2273T mutation which resulted in the Arg778Leu mutation, while the other three patients carried a G2273A mutation which resulted in Arg778Gln (figure) in exon 8 of the ATP7B gene. These codon 778 mutations have not been reported in white populations. The frequency of the Arg778Leu mutation in a total of 44 independent chromosomes examined was 11.4%, and that of Arg778Gln was 9.1%. In addition to the missense mutation in exon 8,

Department of Internal Medicine, National Taiwan University Hospital, 7 Chung-Shan South Road, Taipei, Taiwan 10016, ROC

L-M Chuang
H-P Wu
M-H Jang
T-Y Tai

Department of Pediatrics, National Taiwan University, Taipei, Taiwan, ROC
T-R Wang
W-C Sue

Graduate Institute of Clinical Medicine, National Taiwan University Hospital, Taipei, Taiwan, ROC
B J Lin

Research Institute, The Hospital for Sick Children, Toronto, Ontario, Canada
D W Cox

Correspondence to: Dr Chuang.

Received 4 October 1995.
Revised version accepted for publication 25 January 1996.

* Previously described as Arg779Leu. Three erroneous nucleotides at positions 1039, 1044, and 1056 have been deleted from Thomas *et al*¹⁴ resulting in one less amino acid in the designation.

Nucleotide sequence analyses of mutations in codon 778 of the ATP7B gene. Substitutions of nucleotide 2273 (arrow) were studied in a normal subject and three patients, a heterozygote (WD-1) and homozygote (WD-2) for G2273T mutation, and a heterozygote (WD-3) for G2273A mutation, respectively.

Common haplotypes in 22 Taiwanese families with WD

	D13S314	D13S301	D13S316	WD chromosomes		Normal chromosomes		p
				No	%	No	%	
A	141 (11)	148 (2)	144 (6)	9		3		
	141 (11)	150 (1)	144 (6)	6		2		
	141 (11)	152 (11)	144 (6)	2		1		
Subtotal				17	38.6	6	13.6	<0.015
B	150 (7)	144 (4)	151 (2)	7		2		
	148 (8)	144 (4)	151 (2)	1		1		
Subtotal				8	18.2	3	6.8	
C	150 (7)	144 (4)	144 (6)	4		3		
	148 (8)	144 (4)	144 (6)	1		2		
	150 (7)	148 (2)	144 (6)	1		3		
	150 (7)	142 (5)	144 (6)	0		1		
Subtotal				6	13.6	9	20.5	
D	148 (8)	144 (4)	147 (4)	4		3		
	148 (8)	142 (5)	147 (4)	1		0		
Subtotal				5	11.4	3	6.8	
E	149 (8)	150 (1)	144 (6)	3	6.8	1	2.3	
F	141 (11)	150 (1)	151 (2)	2		0		
G	147 (9)	140 (6)	151 (2)	2	4.5	1	2.3	
H	148 (8)	150 (1)	151 (2)	1		0		
Others				0		21	47.8	<0.001

Brackets denote designation of alleles according to Thomas *et al.*⁷

there was a previously unreported conservative change (C2250G), suggesting that a normal polymorphism was present in this region. To correlate the haplotypes with mutations, we found that the Arg778Leu mutation was segregating with haplotype D (8-4-4 in D13S314-301-316), while the Arg778Gln was found with haplotype E (8-1-6 in D13S314-301-316). These data strongly support the occurrence of different haplotypes associated with different mutations as suggested previously.¹⁹ We have not yet found the mutation associated with haplotype A, the most frequent haplotype in the Chinese patients with WD. The amino acid changes of codon 778, which is located in the transmembrane domain (Tm 4), might cause disruption of the appropriate anchorage of the transporter in the membrane. Whether Arg778Leu or Arg778Gln play the same role in the pathogenesis of WD requires further biological studies. However, the changes from a basic amino acid (Arg) to a neutral (Leu) or

even to an acidic amino acid (Gln) found in this study predict a dramatic change in the primary and secondary structure of this protein, which could culminate in the impaired copper transport.

This work was supported by grants from the National Science Council, Republic of China (NSC 84-2331-B-002-224-M02) and from National Taiwan University Hospital (NTUH-84208-B41).

- 1 Danks DM. Disorders of copper transport. In: Scriver CR, Beaudet AL, Sly WS, Valle D, eds. *The metabolic basis of inherited disease*. 6th ed. New York: McGraw-Hill, 1989: 1411-31.
- 2 Frydman M, Bonne-Tamir B, Farrer LA, *et al.* Assignment of the gene for Wilson disease to chromosome 13: linkage to the esterase D locus. *Proc Natl Acad Sci USA* 1985;82: 1819-21.
- 3 Bonne-Tamir B, Farrer LA, Frydman M, Kanaiah H. Evidence for linkage between Wilson disease and esterase D in three kindreds: detection of linkage for an autosomal recessive disorder by the family study method. *Genet Epidemiol* 1986;3:201-9.
- 4 Bowcock AM, Farrer LA, Hebert JM, Bale AE, Cavalli-Sforza LL. A contiguous linkage map of chromosome 13q with 39 distinct loci separated on average by 5.1 centimorgans. *Genomics* 1991;11:517-29.

- 5 Thomas GR, Roberts EZ, Rosales TO, *et al.* Allelic association and linkage studies in Wilson disease. *Hum Molec Genet* 1993;2:1401-5.
- 6 Petrukhin K, Fischer SG, Pirastu M, *et al.* Mapping, cloning and genetic characterization of the region containing the Wilson disease gene. *Nature Genet* 1993;5:338-43.
- 7 Thomas GR, Bull PC, Roberts EA, Walshe JM, Cox DW. Haplotype studies in Wilson disease. *Am J Hum Genet* 1994;54:71-8.
- 8 Bull PC, Thomas GR, Rommens JM, Forbes JR, Cox DW. The Wilson disease gene is a putative copper transporting P-type ATPase similar to the Menkes gene. *Nature Genet* 1993;5:327-37.
- 9 Tanzi RE, Petrukhin K, Chernov I, *et al.* The Wilson disease gene is a copper transporting ATPase with homology to the Menkes disease gene. *Nature Genet* 1993;5:344-50.
- 10 Vulpe C, Levinson B, Whitney S, Packman S, Gitschier J. Isolation of a candidate gene for Menkes disease and evidence that it encodes a copper-transporting ATPase. *Nature Genet* 1993;3:7-13.
- 11 Mercer JF, Livingston J, Hall B, *et al.* Isolation of a partial candidate gene for Menkes disease by positional cloning. *Nature Genet* 1993;3:20-5.
- 12 Chelly J, Tumber A, Tonnesen T, *et al.* Isolation of a candidate gene for Menkes disease that encodes a potential heavy metal binding protein. *Nature Genet* 1993;3:14-19.
- 13 Green NM, Stokes DL. Structural modeling of P-type ion pumps. *Acta Physiol Scand* 1993;146:59-68.
- 14 Thomas GR, Forbes JR, Roberts EA, Walshe JM, Cox DW. The Wilson disease gene: spectrum of mutations and their consequences. *Nature Genet* 1995;9:210-17.
- 15 Chuang LM, Tai TY, Wang TR, *et al.* Esterase D and retinoblastoma gene loci are tightly linked to Wilson's disease in Chinese pedigrees from Taiwan. *Hum Genet* 1991;87:465-8.
- 16 Strickland GT, Frommer D, Leu ML, Pollard R, Sherlock S, Cummings JN. Wilson's disease in the United Kingdom and Taiwan. *Q J Med* 1973;42:619-38.
- 17 Strickland GT, Leu ML. Wilson's disease: clinical and laboratory manifestations in 40 patients. *Medicine* 1975;54:113-37.
- 18 Cox DW, Fraser FC, Sass-Kortsak A. A genetic study of Wilson's disease: evidence for heterogeneity. *Am J Hum Genet* 1972;24:646-66.
- 19 Bull PC, Cox DW. Wilson disease and Menkes disease: new handles on heavy-metal transport. *Trends Genet* 1994;10:246-52.