

Germline and somatic mosaicism in a female carrier of Hunter disease

Roseline Froissart, Irène Maire, Véronique Bonnet, Thierry Levade, Dominique Bozon

Abstract

Carrier detection in a mucopolysaccharidosis type II family (Hunter disease) allowed the identification of germline and somatic mosaicism in the patient's mother: the R443X mutation was found in a varying proportion in tested tissue (7% in leucocytes, lymphocytes, and lymphoblastoid cells, and 22% in fibroblasts). The proband's sister carries the at risk allele (determined by haplotype analysis), but not the mutation. In sporadic cases of X linked diseases, germline mosaicism of the proband's mother is difficult to exclude and should be considered in genetic counselling.

(J Med Genet 1997;34:137-140)

Keywords: mucopolysaccharidosis type II; mosaicism; mutation.

Hunter disease or mucopolysaccharidosis type II (MPS II) is an X linked lysosomal storage disease resulting from the deficiency of iduronate-2-sulphatase (IDS, EC 3.1.6.13).¹ The IDS gene has been mapped to Xq28.² A full length cDNA clone containing an open reading frame of 1650 bp has been isolated and sequenced.³ The IDS gene spans 24 kb and contains nine exons^{4,5} and has been completely sequenced⁶ (GenBank accession number L43581). A pseudogene located 20 kb distal to the active gene has been characterised⁶ and is involved in recombinations with the IDS gene.^{7,8} Various IDS locus alterations have been described,⁹⁻¹² such as large deletions and rearrangements, but about 80% of the mutations are point mutations, minor deletions, or insertions. Most of them are private except for some that more frequently occur on CpG dinucleotides.

Female carriers of MPS II are asymptomatic though they often show decreased levels of IDS activity in serum or leucocytes or both. Cloned fibroblasts¹³ or hair roots¹⁴ have also been studied for carrier detection. However, attempts to identify carriers by enzyme assays have often led to ambiguous results because of non-random X inactivation. Indirect genotype analysis using intragenic and closely linked flanking polymorphisms, combined with enzymatic and pedigree analyses, allows carrier detection in some but not all families.¹⁵ Identification of the mutation in a patient with MPS II allows fast and reliable carrier detection in related at risk females.

The carrier status of females in a family with an isolated case of Hunter disease was studied

by haplotype, enzyme analysis, and mutation identification. This study led to the identification of germline and somatic mosaicism in the proband's mother.

Material and methods

CASE REPORT

The patient was the first child of healthy parents. The diagnosis was suspected at 3 years but was assessed at 6 years; increased excretion of glycosaminoglycans (heparan sulphate and dermatan sulphate) in urine was found as well as IDS deficiency in serum. He is now 17 years old and presents with coarse facial features, enlarged tongue, kyphosis, joint stiffness that has required surgery, short stature, hepatosplenomegaly, cardiopathy (mitral and aortic insufficiency, mitral stenosis), deafness, and mental retardation. The pedigree of the family is shown in fig 1.

IDS ACTIVITY

IDS activity was determined using the reagent kit provided by The Hospital for Sick Children, Toronto, Canada as previously described.^{16,17}

POLYMORPHIC MARKERS

Genomic DNA was extracted from leucocytes, lymphocytes, fibroblasts, or lymphoblastoid cells according to Jeanpierre.¹⁸ Genomic DNA from leucocytes was digested by *StuI*, *TaqI*, and *BanI*. Southern blotting was carried out using standard procedures. DNA probes, radiolabelled with $\alpha^{32}\text{P}$ -dCTP by random priming, included: IDS cDNA (1.7 kb *NotI/XhoI* fragment from the cDNA clone pB2Sc17 containing the complete coding region of IDS), probe II-10 at locus DXS466,¹⁹ probe U6.2-20E at locus DXS304,²⁰ and probe VK21A at locus DXS296.²¹ Haplotypes were determined with the *StuI* intragenic polymorphism (in intron 8) and three extragenic polymorphisms, *BanI* (U6.2-20E) and *TaqI* (II-10 and VK21A).

A microsatellite at locus DXS1113, located 10 kb distal to the IDS gene, was analysed as described by Weber *et al.*²²

AMPLIFICATION OF GENOMIC DNA

Exon 9 was amplified as a 580 bp fragment with oligonucleotides 5'I: 5'-CCATTCTGCTCTGTCGCTTC-3' (nt 54371-54352 in intron 8) and V6: 5'-CAAACGACCAGCTCTAACTC-3' (nt 53792-53812 in exon 9). PCR conditions were: 94°C for 30 seconds, 61°C for 30 seconds, and 72°C for 45 seconds, for 30 cycles.

Centre d'Etudes des Maladies Métaboliques, Biochimie Bâtiment D, Hôpital Debrousse, 29 Rue Soeur Bouvier, 69322 Lyon Cedex 05, France
R Froissart
I Maire
V Bonnet
D Bozon

Laboratoire de Biochimie Maladies Métaboliques, CJF INSERM 9206, CHU Rangueil, 1 Avenue Jean Poulhès, 31054 Toulouse Cedex, France
T Levade

Correspondence to: Dr Froissart.

Received 15 May 1996
Revised version accepted for publication
9 September 1996




Figure 1 Family pedigree showing the segregation of the haplotypes and R443X mutation.

Amplification of genomic DNA from hair roots was performed as follows: each hair root was rinsed in distilled water and in 100% ethanol, then heated to 94°C for 30 minutes in PCR buffer. Amplification was performed with two sets of primers: 5'I/V6 and microsatellite DXS1113 primers²² for 30 seconds at 94°C, 45 seconds at 60°C, and one minute at 72°C, for 30 cycles.

EXON 9 SEQUENCING

After amplification, PCR products were purified on a 5% acrylamide gel and directly sequenced with the Thermosequense cycle sequencing kit (Amersham) using $\gamma^{33}\text{P-ATP}$.

R443X MUTATION ANALYSIS

Allele specific oligonucleotide (ASO) hybridisation

The PCR products were analysed by ASO hybridisation. ASO used for the detection of the normal and mutant R443X sequences were 5'-AGCATTTCGATTCCG-3' and 5'-AGCATTTCGATTCCG-3' respectively. Washing conditions were 5 × SSC at 37°C for 2 × 10 minutes, and 2 × SSC at 42°C for 3 × 15 minutes.

TaqI digestion

The R443X mutation abolishes a *TaqI* site. After digestion of the PCR products, the mutation is detected by the presence of an abnormal

580 bp band instead of the normal 410 and 170 bp bands.

Quantitative PCR

To assess the relative proportion of mutant (R443X) and normal DNA, exon 9 was amplified as described previously for 29 cycles, and 1 μCi of $\alpha^{32}\text{P-dCTP}$ was added before the last cycle. Radiolabelled PCR products were digested by *TaqI* and separated on an 8% acrylamide non-denaturing gel. The gel was dried and autoradiographed. Radioactivity of each band was quantified by scanning the film. Then the bands were cut from the dried gel and radioactivity content estimated by scintillation counting.

Results

IDS ACTIVITY

IDS activity was borderline in the mother's leucocytes and within the normal range in serum and lymphoblastoid cells (table 1). IDS activity was found to be normal in the grandmother's leucocytes and serum and in the sister's lymphoblastoid cells. Study of 25 fibroblast clones from the sister did not show any deficient clone (data not shown).

HAPLOTYPE ANALYSIS

The results of haplotype analysis are shown in fig 1: microsatellite DXS1113 was the only informative marker. Unexpectedly, the patient's sister had inherited from her mother the same X chromosome as her brother.

MUTATION R443X

Exon 9 was systematically sequenced (presence of several mutational hot spots^{9 10 12}), allowing the identification of the R443X mutation in the patient's genomic DNA. The presence of this mutation was tested in the mother's leucocytes by *TaqI* digestion of exon 9 PCR product: this showed an abnormal distribution of digested and undigested PCR products (not shown). The mother's genomic DNA extracted from various sources (leucocytes, lymphocytes, lymphoblastoid cells, fibroblasts, and hair roots) was analysed by ASO hybridisation to test the hypothesis of somatic mosaicism: the mutant allele was present with a low and varying intensity (fig 2). The mutation was not detected in the grandmother's leucocytes nor in the leucocytes of the sister who had inherited the at risk haplotype.

DNA from 35 separate hair roots of the mother was used as template for PCR and R443X analysis: only one of them hybridised with both normal and mutant probes. Microsatellite analysis (DXS1113) was performed on each hair root and confirmed the presence of the two alleles (mother's karyotype 46,XX).

The relative proportion of mutant and normal DNA in the mother's cells was assessed by quantitative PCR (fig 3). In control DNA, only the 410 and 170 bp bands were detected, and in the DNA from an obligate carrier radioactivity of the abnormal 580 bp band represented about 50% of the total radioactivity, as expected. For the patient's mother, the

Table 1 IDS activity in various samples of the patient and his family

	Serum ($\text{pmol.h}^{-1}.\text{ml}^{-1}$)	Leucocytes ($\text{pmol.h}^{-1}.\text{mg}^{-1}$)	Lymphoblastoid cells ($\text{pmol.h}^{-1}.\text{mg}^{-1}$)
Patient			0.5
Mother	1355	77	44
Sister			62
Grandmother	1785	152	
Control	1301	118	44
Normal range	1185-1900	75-158	20-65




Figure 2 Detection of the R443X mutation by allele specific oligonucleotide hybridisation. After amplification of exon 9 as a 580 bp fragment, the R443X mutation was tested for after hybridisation with the normal and the mutant probes. Mother's hair roots (A1 to C11). Patient's fibroblasts (D1). Mother's leucocytes (D2), lymphocytes (D3), lymphoblastoid cells (D4), fibroblasts (D5). Grandmother's leucocytes (D6), fibroblasts (D7). Sister's lymphoblastoid cells (D8), fibroblasts (D9). Control leucocytes (D10).




Figure 3 Quantitative PCR. Exon 9 was radiolabelled during the PCR and digested with *TaqI*. In normal DNA, the 580 bp fragment is cut into 410 and 170 bp fragments. The R443X mutation abolishes a *TaqI* site and is detected by the presence of a 580 bp fragment. The relative proportion of mutant and normal DNA was quantified by autoradiography and scintillation counting. Patient's fibroblasts (lane 1), mother's leucocytes (lane 2), mother's lymphocytes (lane 3), mother's lymphoblastoid cells (lane 4), mother's fibroblasts (lane 5), heterozygote's leucocytes (lane 6), control leucocytes (lane 7). DNA fragment sizes are given in base pairs.

proportion of the abnormal 580 bp band was about 7% in leucocytes, lymphocytes, and lymphoblastoid cells, and 22% in fibroblasts.

Discussion

Carrier detection in this family was first performed by measurement of IDS activity and haplotype analysis: the results of IDS activity were inconclusive for the mother, and the carrier status of the patient's sister could not be determined.

The R443X mutation was later identified in the patient. This mutation, which occurs on a CpG dinucleotide, introduces a stop codon at position 443 and creates a truncated, non-functional protein.²³ This mutation has been described in several patients of different ethnic origin.²³⁻²⁵ Haplotype and mutation analysis showed that the mother carries a mutant or

wild type sequence on the same chromosome. The low proportion of mutant chromosomes varied from tissue to tissue but remained lower than in a normal carrier (the highest proportion was observed in fibroblasts). These results suggest somatic mosaicism in the mother. Her daughter, who inherited the at risk haplotype, does not carry the R443X mutation and could be excluded as a carrier. Therefore, our findings strongly support the existence of both germinal and somatic mosaicism in the mother.

Somatic mosaicism can be easily detected when the proportion of mutant cells is very different from 50%. However, the finding of a classical heterozygote status in leucocytes does not rule out mosaicism and, in some instances, both haplotype and sequence analysis have been necessary to recognise mosaicism.²⁶

To date, no case of somatic and germinal mosaicism in Hunter disease has been reported. In a previous report, Ben Simon-Schiff *et al*²⁷ suggested the presence of germinal mosaicism in an obligate carrier, based on discordance of normal IDS activity in serum and random X inactivation, but no clear evidence of germline mosaicism was obtained.

Several mechanisms have been proposed for the occurrence of somatic and germline mosaicism. The most likely explanation for the existence of two populations of cells in the mother is that the mutation occurred as a result of a mitotic error at an early embryonic stage; the postzygotic mitotic mutation must have occurred early enough to be transmitted to her germline cells, most probably before differentiation of the three major germ layers (endoderm, mesoderm, ectoderm), and therefore after the first cell division and before the late blastocyst stage of development.²⁸ Alternatively, an uncorrected half chromatid mutation in one of the parental gametes could have established mosaicism after the first mitotic division after conception.²⁹ As a third possibility, the mother could be a chimera consisting of two fertilised eggs, one carrying the mutation³⁰; in this case, the grandmother could also be a carrier. In our study, the grandmother was not a carrier as she does not carry the at risk allele.

In the probands' mother, germline mosaicism, either isolated³¹⁻³⁴ or as part of somatic mosaicism,^{26 35-37} has been documented in a growing number of X linked disorders and germline mosaicism cannot be excluded in the non-carrier mothers of sporadic cases. Therefore, the risk for mosaicism in families with isolated cases might be higher than previously reported. Bakker *et al*²² estimated that the incidence of gonadal mosaicism could be as high as 14% for the at risk haplotype in new cases of DMD. Forty three families with a sporadic case of Hunter disease were studied in our laboratory. In four cases the proband's mother was not found to be a carrier of the mutation and germline mosaicism could not be excluded. Prenatal diagnosis should consequently be offered to every mother of a sporadic case.

We thank Drs J J Hopwood, J L Mandel, T J M Hulsebos, and G R Sutherland for providing probes pB2Sc17, U6.2-20E, II-10, and VK21A respectively, and Drs F Prieur and G Bourrouillou for karyotype analysis. This work was supported by "Vaincre les Maladies Lysosomales" (VML) and "Programme Hospitalier de Recherche Clinique" (PHRC).

- 1 Neufeld EF, Muenzer J. The mucopolysaccharidoses. In: Scriver CR, Beaudet AL, Sly WS, Valle D, eds. *The metabolic basis of inherited disease*. 7th ed. New York: McGraw-Hill: 1995:2465-94.
- 2 Wilson PJ, Suthers GK, Callen DF, et al. Frequent deletions at Xq28 indicate genetic heterogeneity in Hunter syndrome. *Hum Genet* 1991;86:505-8.
- 3 Wilson PJ, Morris DS, Anson DS, et al. Hunter syndrome: isolation of an iduronate-2-sulfatase cDNA clone and analysis of patient DNA. *Proc Natl Acad Sci USA* 1990;87:8531-5.
- 4 Flomen RH, Green EP, Green PM, Bentley DR, Giannelli F. Determination of the organisation of coding sequences within the iduronate sulphate sulphatase (IDS) gene. *Hum Mol Genet* 1993;2:5-10.
- 5 Wilson PJ, Meaney CA, Hopwood JJ, Morris CP. Sequence of the iduronate-2-sulfatase (IDS) gene. *Genomics* 1993;17:773-5.
- 6 Timms KM, Lu F, Shen Y, et al. 130 kb of DNA sequence reveals two new genes and a regional duplication distal to the human iduronate-2-sulfate sulfatase locus. *PCR Methods and Applications* 1995;5:71-8.
- 7 Bondeson ML, Dahl N, Malmgren H, et al. Inversion of the IDS gene resulting from recombination with IDS-related sequences is a common cause of the Hunter syndrome. *Hum Mol Genet* 1995;4:615-21.
- 8 Birot AM, Bouton O, Froissart R, Maire I, Bozon D. IDS gene-pseudogene exchange responsible for an intragenic deletion in a Hunter patient. *Hum Mutat* 1996;8:44-50.
- 9 Hopwood JJ, Bunge S, Morris CP, et al. Molecular basis of mucopolysaccharidosis type II: mutations in the iduronate-2-sulphatase gene. *Hum Mutat* 1993;2:435-42.
- 10 Jonsson JJ, Aronovich EL, Braun SE, Whitley CB. Molecular diagnosis of mucopolysaccharidosis type II (Hunter syndrome) by automated sequencing and computer-assisted interpretation: toward mutation mapping of the iduronate-2-sulfatase gene. *Am J Hum Genet* 1995;56:597-607.
- 11 Goldenfum SL, Young E, Michelakakis H, Tsagarakis S, Winchester B. Mutation analysis in 20 patients with Hunter disease. *Hum Mutat* 1996;7:76-8.
- 12 Olsen TC, Eiken HG, Knappskog PM, et al. Mutations in the iduronate-2-sulfatase gene in five Norwegians with Hunter syndrome. *Hum Genet* 1996;97:198-203.
- 13 Migeon BR, Sprengle JA, Liebaers I, Scott JF, Neufeld EF. X-linked Hunter syndrome: the heterozygous phenotype in cell culture. *Am J Hum Genet* 1977;29:448-54.
- 14 Yutaka T, Fluharty AL, Stevens RL, Kihara H. Iduronate sulfatase analysis of hair roots for identification of Hunter syndrome heterozygotes. *Am J Hum Genet* 1978;30:575-82.
- 15 Gal A, Beck M, Sewell AC, Morris CP, Schwinger E, Hopwood JJ. Gene diagnosis and carrier detection in Hunter syndrome by the iduronate-2-sulphatase cDNA probe. *J Inherited Metab Dis* 1992;15:342-6.
- 16 Hopwood JJ. α -L-iduronidase, β -D-glucuronidase and 2-sulfo-L-iduronate-2-sulfatase: preparation and characterization of radioactive substrates from heparin. *Carbohydr Res* 1979;69:203-16.
- 17 Wasteson A, Neufeld EF. Iduronate sulfatase from human plasma. *Methods Enzymol* 1982;43:573-8.
- 18 Jeanpierre M. A rapid method for the purification of DNA from blood. *Nucleic Acids Res* 1987;15:9611.
- 19 Hulsebos TJM, Oostra BA, Broersen S, Smits A, van Oost BA, Westerveld A. New distal marker closely linked to the fragile X locus. *Hum Genet* 1991;87:369-72.
- 20 Rousseau F, Vincent A, Oberle I, Mandel JL. New informative polymorphism at the DXS 304 locus, a close distal marker for the fragile X locus. *Hum Genet* 1990;84:263-6.
- 21 Suthers GK, Callen DF, Hyland VJ, et al. A new DNA marker tightly linked to the fragile X locus (FRAXA). *Science* 1989;246:1298-300.
- 22 Weber C, Oudet C, Johnson S, Schlessinger D, Hanauer A. Dinucleotide repeat polymorphism close to IDS gene in Xq27.3-q28 (DXS 1113). *Hum Mol Genet* 1993;2:612.
- 23 Sukegawa K, Tomatsu S, Tamai K, et al. Intermediate form of mucopolysaccharidosis type II (Hunter disease): a C¹³²⁷ to T substitution in the iduronate sulfatase gene. *Biochem Biophys Res Commun* 1992;183:809-13.
- 24 Bunge S, Steglich C, Beck M, et al. Mutation analysis of the iduronate-2-sulfatase gene in patients with mucopolysaccharidosis type II (Hunter syndrome). *Hum Mol Genet* 1992;1:335-9.
- 25 Froissart R, Blond JL, Maire I, et al. Hunter syndrome: gene deletions and rearrangements. *Hum Mutat* 1993;2:138-40.
- 26 Sommer SS, Knöll A, Greenberg CR, Ketterling RP. Germline mosaicism in a female who seemed to be a carrier by sequence analysis. *Hum Mol Genet* 1995;4:2181-2.
- 27 Ben Simon-Schiff E, Bach G, Zlotogora J, Abeliovich D. Combined enzymatic and linkage analysis for heterozygote detection in Hunter syndrome: identification of an apparent case of germinal mosaicism. *Am J Med Genet* 1993;47:837-42.
- 28 Kontusaari S, Tromp G, Kuivaniemi H, Stolle C, Pope FM, Prockop DJ. Substitution of aspartate for glycine 1018 in the type III procollagen (COL3A1) gene causes type IV Ehlers-Danlos syndrome: the mutated allele is present in most blood leukocytes of the asymptomatic and mosaic mother. *Am J Hum Genet* 1992;51:497-507.
- 29 Cantu JM, Rivas F, Rivera H, Ruiz C. The prezygotic origin of structural mosaicisms. *Ann Genet (Paris)* 1985;28:73-4.
- 30 Witkowski R. Germinal "mosaicism" - germline mutation or chimerism? *Hum Genet* 1992;88:359-60.
- 31 Darras BT, Blattner P, Harper JF, Spiro AJ, Alter S, Francke U. Intragenic deletions in 21 Duchenne muscular dystrophy (DMD)/Becker muscular dystrophy (BMD) families studied with the dystrophin cDNA: location of breakpoints on *HindIII* and *BglII* exon-containing fragment maps, meiotic and mitotic origin of the mutations. *Am J Hum Genet* 1988;43:620-9.
- 32 Bakker E, Veenema H, den Dunnen JT, et al. Germinal mosaicism increases the recurrence risk for "new" Duchenne muscular dystrophy mutations. *J Med Genet* 1989;26:553-9.
- 33 Gitschier J, Levinson B, Lehesjoki AE, de la Chapelle A. Mosaicism and sporadic haemophilia: implications for carrier detection. *Lancet* 1989;i:273-4.
- 34 Wilton SD, Chandler DC, Kakulas BA, Laing NG. Identification of a point mutation and germinal mosaicism in a Duchenne muscular dystrophy family. *Hum Mutat* 1994;3:133-40.
- 35 Bröcker-Vriends AHJT, Briët E, Dreesen JCFM, et al. Somatic origin of inherited haemophilia. *Hum Genet* 1990;85:288-92.
- 36 Voit T, Neuen-Jacob E, Mahler V, Jauch A, Cremer M. Somatic mosaicism for a deletion of the dystrophin gene in a carrier of Becker muscular dystrophy. *Eur J Pediatr* 1992;151:112-16.
- 37 Bunyan DJ, Robinson DO, Collins AL, Cockwell AE, Bullman HMS, Whittaker PA. Germline and somatic mosaicism in a female carrier of Duchenne muscular dystrophy. *Hum Genet* 1994;93:541-4.