

Review articles

- Defining the genetic contribution of type 2 diabetes mellitus *J van Tilburg, T W van Haften, P Pearson, C Wijmenga* 569
 Ectodermal dysplasias: a new clinical-genetic classification *M Priolo, C Laganà* 579

Original articles

- β -adrenergic antagonists improve sleep and behavioural disturbances in a circadian disorder, Smith-Magenis syndrome
H De Leersnyder, M de Blois, M Vekemans, D Sidi, E Villain, C Kindermans, A Munnich 586
 Breakpoint analysis of Turner patients with partial Xp deletions: implications for the lymphoedema gene location
C A Boucher, C A Sargent, T Ogata, N A Affara 591
 Classification and genetic features of neonatal haemochromatosis: a study of 27 affected pedigrees and molecular analysis
 of genes implicated in iron metabolism *A L Kelly, P W Lunt, F Rodrigues, P J Berry, D M Flynn, P J McKiernan, D A Kelly,
 G Mieli-Vergani, T M Cox* 599

Letters to the Editor

- Clustering and frequency of mutations in the retinal guanylate cyclase (*GUCY2D*) gene in patients with dominant cone-rod
 dystrophies *A M Payne, A G Morris, S M Downes, S Johnson, A C Bird, A T Moore, S S Bhattacharya, D M Hunt* 611
 A G339R mutation in the *CTNS* gene is a common cause of nephropathic cystinosis in the south western Ontario Amish
 Mennonite population *C A Rupa, D Matsell, S Surry, V Siu* 615
 De novo terminal deletion of chromosome 15q26.1 characterised by comparative genomic hybridisation and FISH with locus
 specific probes *H Tönnies, I Schulze, H-C Hennies, L M Neumann, R Keitzer, H Neitzel* 617
 Interstitial deletion of chromosome 11 (q22.3-q23.2) in a boy with mild developmental delay *M Syrrou, J-P Fryns* 621
 Microdeletion in the *FMR-1* gene: an apparent null allele using routine clinical PCR amplification *M R Hegde, B Chong,
 M Fawkner, N Lambiris, H Peters, A Kenneson, S T Warren, D R Love, J McGaughran* 624
 Non-invasive evaluation of arterial involvement in patients affected with Fabry disease *P Boutouyrie, S Laurent, B Laloux,
 O Lidove, J Grunfeld, D P Germain* 629
 Variation of iron loading expression in C282Y homozygous haemochromatosis probands and sib pairs *C Mura, G Le Gac,
 V Scotet, O Raguene, A-Y Mercier, C Férec* 632
 Haptoglobin genotype as a risk factor for postmenopausal osteoporosis *G P Pescarmona, P D'Amelio, E Morra, G C Isaia* 636
GDNF as a candidate modifier in a type 1 neurofibromatosis (NF1) enteric phenotype *M Bahau, A Pelet, D Vidaud,
 T Lamireau, B Le Bail, A Munnich, M Vidaud, S Lyonnet, D Lacombe* 638
 Six novel mutations in the *PRF1* gene in children with haemophagocytic lymphohistiocytosis *R Clementi, U zur Stadt,
 G Savoldi, S Varotto, V Conter, C De Fusco, L D Notarangelo, M Schneider, C Klersy, G Janka, C Danesino, M Aricò* 643
 A novel mutation in the endothelin B receptor gene in a patient with Shah-Waardenburg syndrome and Down syndrome
J P Boardman, P Syrris, S E Holder, N J Robertson, N Carter, K Lakhoo 646

Notice

Instructions for authors

Electronic letters

- Mitochondrial DNA inheritance in patients with deleted mtDNA *L Quintana-Murci, A Rötig, A Munnich, P Rustin, T Bourgeron* e28
 Screening families with endometrial and colorectal cancers for germline mutations *T Liu, J Chen, S Salahshor, S Kuismanen,
 E Holmberg, H Grönberg, P Peltomäki, A Lindblom* e29
 Novel mutations of *SOX10* suggest a dominant negative role in Waardenburg-Shah syndrome *M H Sham, V C H Lui,
 B L S Chen, M Fu, P K H Tam* e30
 Fibrinogen genotypes (α and β) are associated with plasma fibrinogen levels in Chinese *Y Liu, N Saha, C K Heng,
 S Hong, P S Low* e31
 A case of ring chromosome 2 with growth retardation, mild dysmorphism, and microdeletion of 2p detected using FISH
S L Dee, A T Clark, L R Willatt, J R W Yates e32

JMG

JOURNAL OF MEDICAL GENETICS

www.jmedgenet.com