

Deletion of the *SIM1* gene (6q16.2) in a patient with a Prader-Willi-like phenotype

L Faivre, V Cormier-Daire, J M Lapierre, L Colleaux, S Jacquemont, D Geneviève, P Saunier, A Munnich, C Turleau, S Romana, M Prieur, M C De Blois, M Vekemans

J Med Genet 2002;39:594–596

Apart from Prader-Willi syndrome, which is a well delineated imprinting disorder of the 15q11-q12 region, other chromosome anomalies have been described in a small number of patients with features reminiscent of Prader-Willi syndrome, including hypotonia, progressive obesity, small extremities, and delayed developmental milestones. Among these chromosome anomalies are some cases of interstitial deletion of chromosome 6q¹⁻⁵ and haploinsufficiency of the *SIM1* gene (6q16.2) has been proposed as a candidate gene for obesity.⁶ Here, we report a fifth case of Prader-Willi-like phenotype associated with an interstitial chromosome 6q deletion (6q16.1-q21) detected only by high resolution banding techniques. This suggests that a subgroup of patients with features reminiscent of Prader-Willi syndrome and an interstitial deletion of chromosome 6q16.2 could be delineated.

CASE REPORT

The proband was the only child of a 27 year old mother and a 32 year old father. Intrauterine growth retardation, oligohydramnios, and a left club foot were noted during the third trimester of pregnancy. He was born at term after a normal delivery. His growth parameters were weight 2350 g (–2.5 SD), length 47 cm (–1.5 SD), and OFC 33 cm (–1.5 SD). He was described as floppy and had feeding difficulties in early infancy. He sat at the age of 2 years, walked at 3½ years, and had no speech when we first saw him aged 5 years. Excessive weight gain began at 3 years, with a big appetite and food seeking behaviour. There were no sleep disturbances. His behaviour was hyperactive, with a short attention span and intolerance to frustration. Interaction with other children was non-existent.

At physical examination, weight was +5.5 SD, height was –1 SD, and OFC was +2.5 SD. He had generalised obesity, slightly dysmorphic features including a square and flat face, a large forehead with a protruding metopic ridge, small palpebral fissures, mild strabismus, a thin nose, and thin lips (fig 1). The ears were low set with very small lobes. The hands and feet were small with low implanted thumbs, the external genitalia were normal, and a dry skin with livedo was noted. No malformations were found. Clumsiness and a wide based gait were noted at neurological examination. Metabolic screening and fragile X studies were normal, as was a cerebral MRI. Bone age was 3½ years at the age of 5. Psychometric evaluation showed performance at a 2½ year level (DQ = 50).

Cytogenetic and molecular analyses

In order to rule out Prader-Willi syndrome, a methylation assay of the *SNRPN* gene and a FISH study with the *SNRPN* probe were performed. Both analyses showed normal results, as well as standard cytogenetic and telomere analyses using microsatellite polymorphic markers. However, high resolution banding analysis performed on cultured peripheral blood lymphocytes using R and G banding techniques showed a 46,XY, del(6)(q16.1q21) karyotype (fig 2A). Karyotypes of the parents were normal. Comparative genomic hybridisation study confirmed the deletion (fig 2B). Molecular studies using microsatellite polymorphic markers D6S1709, D6S1580, and D6S447 (6q16.3-q21) were fully informative and showed that the 6q deletion was paternal in origin. In addition, molecular studies using an intragenic microsatellite polymorphic marker within the *SIM1* gene (primers 5'-GGCTGGCTCCAACCTCGG-3' and reverse 5'-GATCAGCAGGCAGGAGAG-3') were fully informative and showed that the *SIM1* gene was deleted (fig 2C).



Figure 1 Picture of the child at the age of 5 years. Note generalised obesity, slightly dysmorphic features including a square and flat face, large forehead with protruding metopic ridge, small palpebral fissures, mild strabismus, thin nose, low set ears and thin lips, and small hands and feet.




Figure 2 (A) High resolution banding of chromosome 6, showing the non-deleted chromosome on the left and the deleted chromosome on the right. (B) Comparative genomic hybridisation studies confirming the 6q deletion. (C) Molecular analysis using intragenic polymorphic microsatellite marker of the *SIM1* gene showing the deletion of paternal origin (F: father, M: mother, C: child).

Table 1 Clinical features observed in Prader-Willi syndrome and five cases of interstitial deletion of chromosome 6q associated with Prader-Willi-like phenotype

	Prader-Willi syndrome	Patients with interstitial 6q deletion and a Prader-Willi-like phenotype
Hypotonia	+	4/4
Feeding problems in infancy	+	3/3
Obesity	+	5/5
Facial dysmorphism	Mild	Moderate
Hypogonadism	+	4/5
Developmental delay	Mild	Severe
Hyperphagia	+	2/5
Short extremities	+	5/5
Cardiac defects	-	2/5
CNS abnormalities	-	3/5

DISCUSSION

Here we report on the fifth case of chromosome 6q deletion in association with a Prader-Willi-like phenotype. A review of five patients with an interstitial chromosome 6q deletion and a Prader-Willi-like phenotype showed that they all shared the following features with Prader-Willi syndrome patients: obesity, hypotonia, short extremities, and developmental delay.²⁻⁵ However, there are also distinct differences. For example, excessive appetite was seen in only two patients and cardiac (bicuspid aortic valve, aortic stenosis, right branch block) as well as neurological abnormalities (polygyria, leucomalacia, seizures, hearing loss, Arnold-Chiari malformation) were found in 2/5 and 3/5 patients, respectively. These differences are summarised in table 1.

Interestingly, four out of five patients described have a 6q16.2 deletion (fig 3). This suggests that the 6q16.2 subband could be regarded as a region of interest for obesity related genes. In addition, a balanced translocation between chromosomes 1p22.1 and 6q16.2 has been reported in a patient with profound obesity.⁶ The authors cloned and sequenced both translocation breakpoints. While the translocation did not appear to affect any transcription unit on 1p22.1, it disrupted the *SIM1* gene on 6q16.2. They hypothesised that haploinsufficiency of *SIM1*, possibly acting upstream or downstream of the melanocortin 4 receptor in the paraventricular nuclei of the hypothalamus, could be responsible for obesity in their subject. Indeed, it has been shown in the mouse that the *Sim1* gene is expressed in the central nervous system where it has

an important role in the development of the supraoptic and the paraventricular nuclei of the hypothalamus.⁷ In addition, there is good evidence from anatomical and pharmacological studies that these nuclei are involved in the regulation of body weight as these neurones express the melanocortin-4 receptor⁸ and appear to be a target of alpha melanocyte stimulating hormone, which inhibits food intake.⁹ Interestingly, the *SIM1* gene is also deleted in our patient. Therefore, this observation provides further support for the hypothesis that haploinsufficiency of the *SIM1* gene might be responsible for the obesity observed in our patient.

There is also some evidence in published reports that other chromosome 6q loci might carry obesity related genes, namely: (1) a patient reported by Stein *et al*⁴ with a Prader-Willi-like phenotype and a deletion distal to the subband 6q16.2 as well as patients with distal deletion of chromosome 6q and obesity alone¹⁰; (2) the observation of patients with 6q duplication and obesity^{11 12}; and (3) the discovery of a major locus for fasting insulin concentrations and insulin resistance with strong pleiotropic effects on obesity related phenotypes on distal chromosome 6q¹³ (fig 3).

Finally, there is good evidence that some chromosome 6q genes are submitted to genomic imprinting (<http://www.geneimprint.com>). For example, transient neonatal diabetes mellitus is associated with paternal uniparental disomy for chromosome 6 or paternal duplication of chromosome 6q24.¹⁴ In our report, the deletion was of paternal origin, as was the duplication of chromosome 6q24.3-q27 in the patient reported by Smith *et al*.¹¹ However, the relationship between imprinting and obesity in chromosome 6q deletion remains unknown. Further molecular studies of similar cases might help to resolve the possible causal relationship between imprinting and obesity.

In conclusion, this observation suggests that in patients with a Prader-Willi phenotype and a normal cytogenetic/molecular study of the 15q11-q12 region, deletion of the 6q16.2 region and the *SIM1* gene in particular should be looked for.

ACKNOWLEDGEMENTS

We thank M Pick and C Ozilou for their excellent technical work and Y Deris for his photographic assistance.

.....

Authors' affiliations

L Faivre, V Cormier-Daire, J M Lapierre, L Colleaux, S Jacquemont, D Geneviève, A Munnich, C Turleau, S Romana, M Prieur, M C De Blois, M Vekemans, Département de Génétique, Hôpital Necker-Enfants Malades, Paris, France
 P Saunier, Service de Pédiatrie, CH de Fontainebleau, Fontainebleau, France




Figure 3 Ideogram of chromosome 6 showing the extent of the deletions in children with Prader-Willi-like phenotype as well as other chromosome anomalies of chromosome 6 and obesity.

Correspondence to: Dr M Vekemans, Département de Génétique, Hôpital Necker-Enfants Malades, 149 rue de Sèvres, 75015 Paris, France; vekemans@necker.fr

REFERENCES

- Hopkin RJ, Schorry E, Bofinger M, Milatovich A, Stern HJ, Jayne C, Saal HM. New insights into the phenotypes of 6q deletion. *Am J Med Genet* 1997;**70**:377-86.
- Turleau C, Demay G, Gabanis MO, Lenoir G, de Grouchy J. 6q1 monosomy: a distinctive syndrome. *Clin Genet* 1988;**34**:38-42.
- Villa A, Urioste M, Bofarull JM, Martinez-Frias ML. A de novo interstitial deletion 6q16.2q21 on chromosome 6. *Am J Med Genet* 1995;**55**:379-83.
- Stein CK, Stred SE, Thomson LL, Smith FC, Hoo JJ. Interstitial 6q deletion and Prader-Willi-like phenotype. *Clin Genet* 1996;**49**:306-10.
- Gilhuis HJ, van Ravenswaaij CM, Hamel BJ, Gabreels FJ. Interstitial 6q deletion with a Prader-Willi-like phenotype: a new case and review of the literature. *Eur J Pediatr Neurol* 2000;**4**:39-43.
- Holder JL Jr, Butte NF, Zinn AR. Profound obesity associated with a balanced translocation that disrupts the SIM1 gene. *Hum Mol Genet* 2000;**9**:101-8.
- Michaud J, Rosenquist T, May N, Fan CM. Development of neuroendocrine lineages requires the bHLH-PAS transcription factor SIM1. *Genes Dev* 1998;**12**:3264-75.
- Mountjoy KG, Mortrud MT, Low MJ, Simerly RB, Cone RD. Localization of the melanocortin-4 receptor (MC4-R) in neuroendocrine and autonomic control circuits in the brain. *Mol Endocrinol* 1994;**8**:1298-308.
- Kalra SP, Dube MG, Pu S, Xu B, Horvath TL, Kalra PS. Interacting appetite-regulating pathways in the hypothalamic regulation of body weight. *Endocr Rev* 1999;**20**:68-100.
- Fryns JP, Bettens W, Van den Berghe H. Distal deletion of the long arm of chromosome 6: a specific phenotype? *Am J Med Genet* 1986;**24**:175-8.
- Smith A, Jauch A, Slater H, Rodson L, Sandanam T. Syndromal obesity due to paternal duplication 6(q24.3-q27). *Am J Med Genet* 1999;**84**:125-31.
- Roland B, Lowry RB, Cox DM, Ferreira P, Lin CC. Familial complex chromosomal rearrangement resulting in duplication/deletion of 6q14 to 6q16. *Clin Genet* 1993;**43**:117-21.
- Duggirala R, Blangero J, Almasy L, Catano G, Bamshad M, Bock A, Duggirala R, Williams K, Mummidi S, Clark RA, Ahuja SS, Dolan MJ, Bologna R, Sen L, Ahuja SK. Major locus for fasting insulin concentrations and insulin resistance on chromosome 6q with strong pleiotropic effects on obesity-related phenotype in nondiabetic Mexican Americans. *Am J Hum Genet* 2001;**68**:1149-64.
- Gardner RJ, Mackay DJ, Mungall AJ, Polychronakos C, Siebert R, Shield JP, Temple IK, Robinson DO. An imprinted locus associated with transient neonatal diabetes mellitus. *Hum Mol Genet* 2000;**9**:589-96.

LETTER TO JMG

A novel locus for late onset amyotrophic lateral sclerosis/motor neurone disease variant at 20q13

A L Nishimura, M Mitne-Neto, H C A Silva, J R M Oliveira, M Vainzof, M Zatz

J Med Genet 2004;41:315–320. doi: 10.1136/jmg.2003.013029

Motor neurone disease includes a heterogeneous group of disorders with motor neurone involvement, such as amyotrophic lateral sclerosis, progressive muscular atrophy, progressive bulbar palsy, and primary lateral sclerosis. Amyotrophic lateral sclerosis is the most common adult onset form of motor neurone disease and involves the lower and upper motor neurones. It is characterised by progressive muscle weakness and atrophy, with fasciculations associated with hyperreflexia and spasticity. One of the proposed mechanisms for amyotrophic lateral sclerosis is degeneration of the motor neurone because of abnormal levels of toxic products that accumulate in the cell. Death usually occurs by respiratory failure about 2–3 years after the first symptoms.¹ About 10% of cases are familial amyotrophic lateral sclerosis, and several loci have been associated with this disease. To date, the only two genes identified have been the zinc–copper superoxide dismutase 1 (*SOD1*) gene, which is located on chromosome 21 (ALS1, MIM105400), and the *Alsin* gene, which is located at 2q33 (ALS2, MIM 205100).^{2–4} Autosomal dominant forms of amyotrophic lateral sclerosis also have been linked to 18q21 (ALS3, MIM 606640), 9q34 (ALS4, MIM 602433), and 15q15.1–q21.1 (ALS5, MIM 602099) and amyotrophic lateral sclerosis–frontotemporal dementia (MIM 105550) to 9q21–22.^{5–9} Moreover, mutations in *Dynein* are associated with motor neurone degeneration and defects in retrograde transport. This gene acts in the cellular division, trafficking, and transport of several proteins, such as *SOD1*.^{10–11}

More recently, two new loci have been associated with amyotrophic lateral sclerosis. One of them, reported by three independent groups, is on chromosome 16; the other is at 20p.^{12–14}

We report a Caucasian Brazilian family with 26 members distributed in three generations affected by a late onset autosomal dominant motor neurone disease. Clinical and neurological examination of 11 living members was compatible with the diagnosis of Caucasian amyotrophic lateral sclerosis and motor neurone disease and long survival.

METHODS

Patients

Figure 1 shows the family pedigree. The probands—three affected sisters (IV-12, IV-13, and IV-14)—were referred to the Human Genome Research Center at the Department of Biology, University of São Paulo, with a diagnosis of motor neurone disease. Extended pedigree analysis showed 26 members were affected (10 men and 16 women); 15 of these already were deceased. Family members reported that patients III-26, IV-10, IV-12, IV-31, and IV-38 died of respiratory failure, although no postmortem confirmation of amyotrophic lateral sclerosis was done. The mean age of death was 49.8 (SD 8.1) years, and the mean age at onset was 38 (SD 6) years.

The diagnosis of amyotrophic lateral sclerosis was based on El Escorial revised criteria.¹⁵ All studies were performed after

Key points

- Motor neurone disease includes a heterogeneous group of disorders with motor neurone involvement, such as amyotrophic lateral sclerosis, progressive muscular atrophy, progressive bulbar palsy, and primary lateral sclerosis.
- Amyotrophic lateral sclerosis is the most common adult onset form of motor neurone disease and involves lower and upper motor neurones. It is characterised by progressive muscle weakness and atrophy, with fasciculations associated with hyperreflexia and spasticity.
- One proposed mechanism for amyotrophic lateral sclerosis is degeneration of the motor neurone because of abnormal levels of toxic products that accumulate in cells. Death usually occurs by respiratory failure about 2–3 years after the first symptoms.
- About 10% of cases are familial amyotrophic lateral sclerosis, and several loci have been associated with this condition. To date, the only two genes identified have been the zinc–copper superoxide dismutase 1 (*SOD1*) gene, which is located on chromosome 21 (ALS1, MIM105400), and the *Alsin* gene, which is located at 2q33 (ALS2, MIM 205100).
- Autosomal dominant forms of amyotrophic lateral sclerosis also have been linked to 18q21 (ALS3, MIM 606640), 9q34 (ALS4, MIM 602433), and 15q15.1–q21.1 (ALS5, MIM 602099), and amyotrophic lateral sclerosis and frontotemporal dementia (MIM 105550) have been linked to 9q21–22.
- Mutations in *Dynein* are associated with motor neurone degeneration and defects in retrograde transport. This gene acts in the cellular division, trafficking, and transport of several proteins, such as *SOD1*.
- We report a large Brazilian Caucasian family with clinical and neurological signs compatible with the diagnosis of amyotrophic lateral sclerosis with slow progression.
- The disease seems to affect both sexes equally, with no evidence of clinical anticipation. Clinical onset occurs between age 31 and 45 years, and the cause of death is respiratory failure. Overall, 12 family members were examined personally. All patients had lower motor neurone symptoms, and five also had bulbar involvement. Electromyography, as well as muscle biopsies, showed a neurogenic pattern.
- We mapped a novel locus for autosomal dominant late onset amyotrophic lateral sclerosis/motor neurone disease (ALS/MND) variant at 20q13.33. The identification of a new gene for ALS/MND will contribute to our understanding of this intriguing disorder.




Figure 1 Family pedigree showing a candidate locus on 20q13.33. ■ man affected by disease; □ unaffected man; ● affected woman; ○ unaffected woman; / deceased; * haplotype inferred. Red text indicates haplotype shared by all patients; blue text indicates haplotype of normal siblings. Recombination events were seen in two patients (IV-1 and IV-34), which allowed placement of the candidate region between D20S430 and D20S173.

informed consent. Demographic and clinical data were taken and followed by a complete clinical and neurological examination. Serum levels of creatine kinase, magnetic resonance imaging, nerve conduction study (motor and sensory), and electromyography were performed in the standard way. Muscle biopsies were taken from the biceps of three affected patients (IV-7, IV-14, and IV-37).

We obtained DNA from the peripheral blood of 25 family members (11 affected and 14 unaffected) according to standard procedures. As the disorder has late onset, we included in the linkage analysis only affected patients, older unaffected siblings, and unrelated unaffected spouses. We performed haplotype reconstruction, on the basis of haplotype analysis of members from generation IV, for three patients (III-12, III-20, and III-22), as well as for member III-19.

Linkage analysis

We excluded all previously known loci for amyotrophic lateral sclerosis. We performed scan analysis with distant markers about 10 cM in length from the ABI Prism Linkage Mapping Set kit (version 2; Applied Biosystems, Foster City, CA, USA). Additional polymorphic markers were included in this analysis to refine the region. The PCR products were analysed in MegaBace 1000 DNA Sequencers (Amersham Biosciences, Little Chalfont, UK). The order of the markers was based on different genetic maps of the Marshfield Medical Research Foundation database and the National Center for Biotechnology Information (NCBI) database. All information about sequence tagged sites was obtained from UniSTS of NCBI and the genome database website. As the allele frequency varies according to the population, we analysed at least 30 chromosomes from normal Brazilian controls. For linkage analysis, we used the MLink program of the Fastlink package (Columbia University, New York, NY, USA) for two point analysis, and we assumed autosomal dominant inheritance with penetrance 1 until age 45 years (as the onset in all affected patients occurs before age 45 years), equal male and female recombination rates, and a

gene frequency of 0.0001 for parametric analysis. Multipoint analysis was performed with SimWalk (version 2.86).¹⁶⁻¹⁸

RESULTS

Clinical features

Table 1 shows the affected family members' clinical and neurological features, as well as the results of complementary examinations.

All patients had lower motor neurone symptoms, with signs in the four limbs, and one also had upper neurone signs. Five patients (IV-23, IV-32, IV-33, IV-34, and IV-37) also had bulbar involvement. Postural tremor, one of the first symptoms to appear, was seen in eight patients, but it was stable during disease progression (IV-1, IV-16, IV-17, IV-23, IV-32, IV-33, IV-34, and IV-37).

The symptom of painful cramps was prominent, long standing, and easily obtained, and had a disabling pattern. As lower motor neurone findings confirmed by electrophysiological studies were the most conspicuous and uniform symptom in all patients, such patients first were classified as having motor neurone disease. The presence of pyramidal signs in one patient (IV-1), who could be classified as having clinically probable, laboratory supported amyotrophic lateral sclerosis, subsequently directed the investigation to the amyotrophic lateral sclerosis/motor neurone disease (AML/MND) group.

Serum creatinine kinase was normal or slightly elevated.

Electromyography in eight patients showed a neurogenic pattern compatible with motor neurone disease. Needle electromyography showed abnormal spontaneous activity such as fasciculations, fibrillations, and positive waves in the tongue and muscles of the upper and lower limbs. On effort, polyphasic and giant motor unit action potentials with reduced recruitment were seen. Sensory and motor nerve conduction studies were normal.

Analyses of muscle biopsies showed a neurogenic pattern, with groups of large and small angulated fibres and fibre type grouping.

Table 1 Clinical and neurological features of patients affected by amyotrophic lateral sclerosis and motor neurone disease

Patient	Sex	First symptoms	Age (years)			Test result		Needle electromyography	Cause of death	Age at death (years)
			At first symptom	At onset of weakness	At ascertainment	Serum creatinine kinase (SU)	Findings on neurological examination			
IV-1	Man	Cramps	31	40	42	20.0 and 23.0*	Lower motor neurone signs in lower limbs, pyramidal tract signs in upper limbs. Postural tremor	Motor neurone disease	NA	NA
IV-12	Woman	Fasciculation	40	40	49	7.0	Lower motor neurone signs in the four limbs	Motor neurone disease	Respiratory failure	49
IV-13	Woman	Fasciculation	43	43	49	19.3 and 28.0*	Lower motor neurone signs in the four limbs	Motor neurone disease	NA	NA
IV-14	Woman	Fasciculation	45	45	47	58.0 and 61.0*	Lower motor neurone signs in four limbs	Motor neurone disease	NA	NA
IV-16	Woman	Fasciculation	41	NA	44	ND	Lower motor neurone signs in four limbs† Postural tremor	ND	NA	NA
IV-17	Woman	Cramps and tremor	30	42	43	ND	Lower motor neurone signs in four limbs† Postural tremor	ND	NA	NA
IV-20	Woman	Weakness	42	42	NA	ND	Lower motor neurone signs in the four limbs†	Motor neurone disease	NA	NA
IV-23	Woman	Cramps	25	38	55	18.5	Lower motor neurone signs in the four limbs, cervical muscles, and tongue; dysarthria; and dysphagia† Postural tremor	Motor neurone disease	NA	NA
IV-32	Man	Tremor	41	46	56	25.3	Lower motor neurone signs in the four limbs and thoracic muscles and dysphagia† Postural tremor	ND	NA	NA
IV-33	Man	Tremor	44	No weakness	49	20.3	Lower motor neurone signs in the four limbs, cervico-thoracic muscles, and tongue† Postural tremor	Motor neurone disease	NA	NA
IV-34	Woman	Fatigue	37	40	45	14.0	Lower motor neurone signs in the four limbs and cervical muscles, dysarthria, and dysphagia† Postural tremor	Motor neurone disease	NA	NA
IV-37	Woman	Cramps	37	38	49	18.0	Sensory loss in left lower limb Lower motor neurone signs in the four limbs and dysphagia† Postural tremor	ND	NA	NA

NA, not applicable (patient alive); ND, not done.

* Normal values <12 SU for women and <20 SU for men.

† Signs of lower motor neurone problems: weakness, atrophy, hypotonia, hypoactive or absent reflexes, and fasciculations.

Table 2 Results of multipoint and two point LOD score analyses for informative markers at 20q13. It is considered informative if LOD >3.0.

Marker	Marker position (cM)*	LOD score								
		Multipoint	Two point at $\theta =$							
			0	0.001	0.01	0.05	0.1	0.2	0.3	0.4
D20s1148	55.60	-6.601	-∞	0.668	2.560	3.535	3.585	2.992	2.003	0.784
D20S430	56.78	1.501	-∞	2.245	3.179	3.564	3.437	2.784	1.887	0.847
D20S164	57.69	7.450	5.704	5.694	5.604	5.196	4.664	3.524	2.268	0.923
D20S94	58.13	7.417	0.263	0.263	0.261	0.246	0.217	0.148	0.083	0.033
D20S171	58.41	7.373	6.025	6.015	5.923	5.503	4.961	3.805	2.537	1.149
D20S173	59.51	3.339	2.316	3.573	4.454	4.711	4.437	3.499	2.299	0.930
D20S93	59.55	3.339	2.665	3.921	4.802	5.055	4.775	3.821	2.590	1.139

*Inferred based on data from MapView-NCBI and Ensembl website.




Fig 2 Multipoint LOD score analysis for 20q13 markers based on the Simwalk program. The most significant polymorphic markers are shown.

Linkage data

We detected a significant linkage at region 20q13.3 with the flanking markers D20S178, D20S196, D20S120, D20S100, D20S102, D20S171, and D20S173. Recombination events in two patients (IV-1 and IV-34), as well as the analysis of 12 additional polymorphic markers (D20S857, D20S1083, D20S839, D20S833, D20S606, D20S183, D20S840, D20S211, D20S1148, D20S430, D20S164, and D20S93), allowed us to locate the candidate gene in a region of about 2.7 Mb between the markers D20S430 and D20S173. According to NCBI MapView, this region contains 17 known genes, one pseudogene, and six predicted genes. Table 2 shows the logarithm of odds (LOD) scores. Two point LOD score analysis showed LOD scores >3 for several markers, with a maximum value of 6.02 at $\theta = 0.0$ for the marker D20S171, while multipoint analysis showed a maximum LOD score of 7.45 for the marker D20S164 (fig 2).

We are currently screening functional candidate genes in the region. Until now, the tubulin beta-1 (*TUBB-1*), cathepsin Z (*CTSZ*), and ATP synthase-epsilon subunit (*ATP5E*) genes were screened through single strand conformation polymorphism followed by direct sequencing of all amplicons, but no pathogenic mutation was identified.

DISCUSSION

We have identified a novel locus for an autosomal dominant late onset ALS/MND at 20q13.33 in a large Brazilian Caucasian family. Recently, a new amyotrophic lateral sclerosis locus was also mapped on chromosome 20.¹⁴ This locus is located in the short arm of chromosome 20 (at 20p13), however; while ours is in the long arm.

The disease seems to affect both sexes equally, with no evidence of clinical anticipation. Clinical onset in affected patients occurs earlier than in familial amyotrophic lateral sclerosis or classic amyotrophic lateral sclerosis (mean age 38 (SD 6) years *v* 46 years and 56 years, respectively). In a previous study of Brazilian patients with sporadic amyotrophic lateral sclerosis, the mean age of onset was 52 years, although the first symptoms presented before age 40 years in 18.1% of patients.¹⁹ The progression of the disease in patients from the present family, however, was slower than in classic amyotrophic lateral sclerosis. In addition, other signs, such as postural tremor and disabling cramps, are atypical, although postural tremor because of increased physiological tremor may be seen in patients with weakness.^{20, 21} On the other hand, the presence of atypical signs has prompted us to search for a differential diagnosis from other causes of motor neurone disease associated with postural tremor, such as extrapyramidal degeneration, distal hereditary motor neuropathy, adult onset spinal muscular atrophy, or even an occasional association of motor neurone disease and essential tremor.

Sporadic cases of motor neurone disease with tremor or choreiform movements, or both, have been described.²² These patients, however, did not show the classical findings of amyotrophic lateral sclerosis-dementia-parkinsonism, which are typical of Guam complex. In these cases, autopsy studies showed degeneration in the lower motor neurones and neostriato-pallido-nigral system or in the upper and lower motor neurones, pallido-luysio-nigral system, and brainstem tegmentum.²³

The diagnosis of distal hereditary motor neuropathy was excluded because none of our patients had pes cavus or nerve conduction abnormalities.

Many years ago, we reported a Brazilian family with seven members affected by late onset autosomal dominant spinal muscular atrophy and symptoms that overlapped with amyotrophic lateral sclerosis.²⁴ In this family, however, all patients showed rapid progression, with death occurring 2–3 years after onset.

We propose classifying the present family as having late onset amyotrophic lateral sclerosis/motor neurone disease with atypical signs. Recently, a family with atypical autosomal dominant amyotrophic lateral sclerosis with normal life expectancy, absence of bulbar involvement, and symmetrical distal distribution of atrophy and weakness was mapped at 9q34.⁶ The authors suggested that this form and distal hereditary motor neuropathy with pyramidal signs could be the same disorder. We agree with those authors' comment that it is not easy to classify these neuromuscular conditions on the basis of clinical and neurological signs.

On the other hand, genetic findings and clinical heterogeneity with atypical findings have been common observations, and other families with this intriguing phenotype may exist. Two large Brazilian genealogies with 80 affected members with late onset (mean age 48.8 years) spinal muscular atrophy (MIM 182980) were reported several years ago.²⁵ Patients from these families showed slow loss of muscle strength, with progressive proximal atrophy, hypoactive or absent deep tendon reflexes, and fasciculations. It will be of interest to verify if they have the disorder we describe.

Intrafamilial clinical variability in patients who carry the same pathogenic mutation has been reported, however, for many gene related diseases, such as autosomal recessive limb girdle muscular dystrophies.²⁶ Furthermore, recent studies showed that modifier genes also could alter clinical phenotype. Faster progression of disease has been reported in patients with a mutation in the ciliary neurotrophic factor (*CNTF*) gene associated with the *SOD1* gene compared than in those who lack the *CNTF* mutation.²⁷ The authors concluded that *CNTF* is a modifier gene that could modulate the progression of the disease, but this finding was not confirmed by other investigators.²⁸

On the other hand, the vascular endothelial growth factor (*VEGF*) gene is considered a risk factor for sporadic amyotrophic lateral sclerosis. Patients homozygous for -2578A/-1154A/-634G or -2578A/-1154G/-634G in the *VEGF* promoter/leader sequence were shown to be at 1.8 times greater risk of amyotrophic lateral sclerosis.²⁹

Conclusion

We mapped a novel gene for an ALS/MND variant at 20q13.33 and are currently screening functional candidate genes in the region for mutations. According to the NCBI Map Viewer, the flanking markers D20S430–D20S173 span a region of about 2.7 Mb that contains 17 known genes, six predicted genes, and one pseudogene.

The *TUBB-1* gene was considered to be a strong candidate gene, because it is the major component of microtubules and, recently, alterations of the axonal transport and microtubule

Electronic database information

Database accession numbers from UniGene (NCBI)

- *TUBB1*: Hs.303023
- *CTSZ*: Hs.252549
- *ATP5E*: Hs.177530

Databases

- MapView and UniSTS of National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>)
- Center for Medical Genetics, Marshfield Medical Research Foundation database (<http://www.marshfieldclinic.org/research/genetics/>)
- Ensembl Genome database (www.ensembl.org)

network have been shown to be potential causes of motor neurodegeneration in amyotrophic lateral sclerosis model mice.³⁰ No mutation was found, however, in this gene or in the *CTSZ* or *ATP5E* genes, although the possibility of an alternative splicing or other post-transcriptional mechanism has not been ruled out yet.

Identification of the causative gene will be very important for enhancing our understanding of the underlying pathological mechanisms in this heterogeneous group of disorders and hopefully will contribute to the opening of new avenues for future therapies.

ACKNOWLEDGEMENTS

We are extremely grateful for the participation of the family members in this study. We also thank Dr Maria Rita Passos-Bueno, Dr Julia, Dr Eliete R Bortolini, Dr Luiz Felipe Rocha Vasconcelos, Dr Marli, Constância Urbani, Antonia Maria Cerqueira, Cynthia Esteves, Manuela Tonini, Kikue Abe, Flavia de Paula, Marta Canovas, Roberto Rivelino, Maria Cristina, Eloisa Moreira, and Karina Lezirivitz for their invaluable help. This work was supported by FAPESP-Cepid, PRONEX, and CNPq.

Authors' affiliations

A L Nishimura, M Mitne-Neto, H C A Silva, J R M Oliveira, M Vainzof, M Zatz, Human Genome Research Center, Department of Biology, Biosciences Institute, University of São Paulo, Brazil

Correspondence to: Dr M Zatz, Centro de Estudos do Genoma Humano, Instituto de Biociências, Departamento de Biologia, Universidade de São Paulo, Rua do Matão 277, Cidade Universitária, São Paulo, Brazil; mayatzatz@usp.br

Conflicts of interest: None declared.

Received 6 August 2003

Accepted for publication 26 October 2003

REFERENCES

- Hand CK, Rouleau GA. Familial amyotrophic lateral sclerosis. *Muscle Nerve* 2002;**25**:135–59.
- Rosen DR, Siddique T, Patterson D, Figlewicz DA, Sapp P, Hentati A, Donaldson D, Goto J, O'Regan JP, Deng HX, Rahmani Z, Krizus A, McKenna-Yasek D, Cayabyab A, Gaston SM, Berger R, Tanzi RE, Halperin JJ, Herzfeldt B, Van den Bergh R, Hung WY, Bird T, Deng G, Mulder DM, Smyth C, Laing NG, Soriano E, Pericak-Vance MA, Haines J, Rouleau GA, Gusella JS, Horvitz HR. Brown RH Mutations in Cu/Zn superoxide dismutase gene are associated with familial amyotrophic lateral sclerosis. *Nature* 1993;**362**:59–62.
- Hadano S, Hand CK, Osuga H, Yanagisawa Y, Otomo A, Devon RS, Miyamoto N, Showguchi-Miyata J, Okada Y, Singaraja R, Figlewicz DA, Kwiatkowski T, Hosler BA, Sagie T, Skaug J, Nasir J, Brown RH Jr, Scherer SW, Rouleau GA, Hayden MR, Ikeda JE. A gene encoding a putative GTPase regulator is mutated in familial amyotrophic lateral sclerosis 2. *Nature Genet* 2001;**29**:166–73.
- Yang Y, Hentati A, Deng HX, Dabbagh O, Sasaki T, Hirano M, Hung WY, Ouahchi K, Yan J, Azim AC, Cole N, Gascon G, Yagmour A, Ben-Hamida M, Pericak-Vance M, Hentati F, Siddique T. The gene encoding alsin, a protein with three guanine-nucleotide exchange factor domains, is mutated in a form of recessive amyotrophic lateral sclerosis. *Nat Genet* 2001;**29**:160–5.
- Hand CK, Khoris J, Salachas F, Gros-Louis F, Lopes AA, Mayeux-Portas V, Brewer CG, Brown RH Jr, Meininger V, Camu W, Rouleau GA. A novel locus for familial amyotrophic lateral sclerosis, on chromosome 18q. *Am J Hum Genet* 2002;**70**:251–6.
- De Jonghe P, Auer-Grumbach M, Irobi J, Wagner K, Plecko B, Kennerson M, Zhu D, De Vriendt E, Van Gerwen V, Nicholson G, Hartung HP, Timmerman V. Autosomal dominant juvenile amyotrophic lateral sclerosis and distal hereditary motor neuropathy with pyramidal tract signs: synonyms for the same disorder? *Brain* 2002;**125**:1320–5.
- Chance PF, Rabin BA, Ryan SG, Ding Y, Scavina M, Crain B, Griffin JW, Cornblath DR. Linkage of the gene for an autosomal dominant form of juvenile amyotrophic lateral sclerosis to chromosome 9q34. *Am J Hum Genet* 1998;**62**:633–40.
- Hentati A, Ouahchi K, Pericak-Vance MA, Nijhawan D, Ahmad A, Yang Y, Rimmler J, Hung W, Schlotter B, Ahmed A, Ben Hamida M, Hentati F, Siddique T. Linkage of a commoner form of recessive amyotrophic lateral sclerosis to chromosome 15q15–q22 markers. *Neurogenetics* 1998;**2**:55–60.
- Hosler BA, Siddique T, Sapp PC, Sailor W, Huang MC, Hossain A, Daube JR, Nance M, Fan C, Kaplan J, Hung WY, McKenna-Yasek D, Haines JL, Pericak-Vance MA, Horvitz HR, Brown RH Jr. Linkage of familial amyotrophic lateral sclerosis with frontotemporal dementia to chromosome 9q21–q22. *JAMA* 2000;**284**:1664–9.
- Hafezparast M, Klocke R, Ruhrberg C, Marquardt A, Ahmad-Annur A, Bowen S, Lalli G, Witherden AS, Hummerich H, Nicholson S, Morgan PJ, Oozageer R, Priestley JV, Averill S, King VR, Ball S, Peters J, Toda T, Yamamoto A, Hiraoka Y, Augustin M, Korhau D, Watterl S, Wabnitz P, Dickneite K, Lampel S, Boehme F, Peraus G, Popp A, Rudelius M, Schlegel J, Fuchs H, de Angelis MH, Schiavo G, Shima DT, Russ AP, Stumm G, Martin JE, Fisher EM. Mutations in dynein link motor neuron degeneration to defects in retrograde transport. *Science* 2003;**300**:808–12.
- Toyoshima I, Sugawara M, Kato K, Wada C, Hirota K, Hasegawa K, Kowa H, Sheetz MP, Masamune O. Kinesin and cytoplasmic dynein in spinal spheroids with motor neuron disease. *J Neurol Sci* 1998;**159**:38–44.
- Ruddy DM, Parton MJ, Al-Chalabi A, Lewis CM, Vance C, Smith BN, Leigh PN, Powell JF, Siddique T, Postumus Meyjes E, Baas F, De Jong V, Shaw CE. Two families with familial amyotrophic lateral sclerosis are linked to a novel locus on chromosome 16q. *Am J Hum Genet* 2003;**73**:390–6.
- Abalkhail H, Mitchell J, Habgood J, Orrell R, De Bellerocche J. A new familial amyotrophic lateral sclerosis locus on chromosome 16q12.1–16q12.2. *Am J Hum Genet* 2003;**73**:383–9.
- Sapp PC, Hosler BA, McKenna-Yasek D, Chin W, Gann A, Genise H, Gorenstein J, Huang M, Sailer W, Scheffler M, Valesky M, Haines JL, Pericak-Vance M, Siddique T, Horvitz HR, Brown RH Jr. Identification of two novel loci for dominantly inherited familial amyotrophic lateral sclerosis. *Am J Hum Genet* 2003;**73**:397–403.
- Brooks Br, Miller RG, Swash M, Munsat TL. For the World Federation of Neurology Research Group on Motor Neuron Diseases. El Escorial revisited: revised criteria for the diagnosis of amyotrophic lateral sclerosis. *ALS* 2000;**2**(suppl 2):S293–8.
- Lathrop GM, Lalouel JM, Julier C, Ott J. Strategies for multilocus linkage analysis in humans. *Proc Natl Acad Sci USA* 1984;**81**:3443–6.
- Terwilliger JD, Ott J. *Handbook of human genetic linkage*. Baltimore and London: Johns Hopkins Press, 1994.
- Sobel E, Lange K. Descent graphs in pedigree analysis: applications to haplotyping, location scores, and marker-sharing statistics. *Am J Hum Genet* 1996;**58**:1323–37.
- Dietrich-Neto F, Callegaro D, Dias-Tosta E, Silva HA, Ferraz ME, Lima JM, Oliveira AS. Amyotrophic lateral sclerosis in Brazil: 1998 national survey. *Arg Neuropsiquiatr* 2000;**58**:607–15.
- Said G, Bathien N, Cesaro P. Peripheral neuropathies and tremor. *Neurology* 1982;**32**:480.
- Mahabe H, Sakamoto K. Evaluation of postural tremor of finger for neuromuscular diseases and its application to the classification. *Electromyogr Clin Neurophysiol* 2002;**42**:205–18.
- Kaneto T, Inoue K, Shimoda K, Doi M, Kitagawa T, Kumada H, Miyata H, Ohama E. An autopsy case of progressive spinal muscular atrophy showing tremor and choreiform movements. *Rinsho Shinkeigak* 2000;**40**:801–6.
- Kato S, Oda M, Murahashi M, Suda N. Motor neuron disease with involvement of the pallidum-luysio-nigral system and mesencephalic tegmentum. *Clin Neuropathol* 1995;**14**:241–4.
- Zatz M, Penha-Serrano C, Frota-Pessoa O, Klein D. A malignant form of neurogenic muscular atrophy in adults, with dominant inheritance. *J Genet Hum* 1971;**19**:337–54.
- Richieri-Costa A, Rogatko A, Levisky R, Finkel N, Frota-Pessoa O. Autosomal dominant late adult spinal muscular atrophy type Finkel. *Am J Med Genet* 1981;**9**:119–28.
- Zatz M, Vainzof M, Passos-Bueno MR. Limb-girdle muscular dystrophy: one gene with different phenotypes; one phenotype with different genes. *Curr Opin Neurology* 2000;**13**:511–7.
- Giess R, Holtmann B, Braga M, Grimm T, Muller-Myhsok B, Toyka KV, Sendner M. Early onset of severe familial amyotrophic lateral sclerosis with a SOD-1 mutation: potential impact of CNTF as a candidate modifier gene. *Am J Hum Genet* 2002;**70**:1277–86.
- Al-Chalabi A, Scheffler MD, Smith BN, Parton MJ, Cudkowicz ME, Andersen PM, Hayden DL, Hansen VK, Turner MR, Shaw CE, Leigh PN, Brown RH Jr. Ciliary neurotrophic factor genotype does not influence

- clinical phenotype in amyotrophic lateral sclerosis. *Ann Neurol* 2003;**54**:130-4.
- 29 **Lambrechts D**, Storkebaum E, Morimoto M, Del-Favero J, Desmet F, Marklund SL, Wyns S, Thijs V, Andersson J, van Marion I, Al-Chalabi A, Bornes S, Musson R, Hansen V, Beckman L, Adolfsson R, Pall HS, Prats H, Vermeire S, Rutgeerts P, Katayama S, Awata T, Leigh N, Lang-Lazdunski L, Dewerchin M, Shaw C, Moons L, Vlietinck R, Morrison KE, Robberecht W, Van Broeckhoven C, Collen D, Andersen PM, Carmeliet P. VEGF is a modifier of amyotrophic lateral sclerosis in mice and humans and protects motoneurons against ischemic death. *Nat Genet* 2003;**34**:383-94.
- 30 **Farah CA**, Nguyen MD, Julien JP, Leclerc N. Altered levels and distribution of microtubule-associated proteins before disease onset in a mouse model of amyotrophic lateral sclerosis. *J Neurochem* 2003;**84**:77-86.

CORRECTION

Faivre L, Cormier-Daire V, Lapierre JM, *et al*. Deletion of the *SIM1* gene (6q16.2) in a patient with a Prader-Willi-like phenotype (*J Med Genet* 2002;**39**:594-96). An erratum has been detected in the reverse primer sequence used for amplification of the intragenic microsatellite polymorphic marker of the *SIM1* gene. The sequence should read 5'-CTCTCCTGCCTGCTGATC-3' instead of 5'-GATCAGCAGGCAGGAGAG-3'.