ONLINE MUTATION REPORT

Two novel aspartoacylase gene (*ASPA*) missense mutations specific to Norwegian and Swedish patients with Canavan disease

T R Olsen, L Tranebjærg, E A Kvittingen, L Hagenfeldt, C Møller, Ø Nilssen

J Med Genet 2002;39:e55 (<http://www.jmedgenet.com/cgi/content/full/39/9/e55>)

Canavan disease (CD) (OMIM 271900) is an autosomal recessive leucodystrophy characterised by swelling and spongy degeneration of the white matter of the brain. The biochemical marker for the disorder is increased level of N-acetylaspartic acid (NAA) in cerebrospinal fluid and urine owing to aspartoacylase (*ASPA*) deficiency.^{1,2} The disease is caused by mutations in the gene encoding the enzyme aspartoacylase (EC 3.5.1.15) at 17p13-ter.^{3,4}

Clinical features are macrocephaly, head lag, and hypotonia from the age of 3-6 months followed by rapidly progressive severe mental retardation. Life expectancy is usually into the teens and development of optic atrophy is typical. Only symptomatic treatment is possible. The prevalence is highest among Ashkenazi Jews with a carrier frequency of 1 in 38 and two mutations, Y231X and E285A, constitute 98% of the CD alleles in this population.^{3,5}

Canavan disease is rare in non-Jewish populations and one mutation, A305E (914C→A), is found in about 40%⁶ to 60%⁵ of the disease alleles reported so far. Mutations are scattered all over the six exons of the aspartoacylase gene. In total, 38 different mutations have been reported of which 35 are listed in The Human Gene Mutation Data Base, Cardiff, UK.⁷⁻¹⁰

Prenatal diagnosis by enzyme assay is complicated by technical difficulties.¹¹⁻¹⁵ Measurement of N-acetylaspartate concentration in amniotic fluid by stable isotope dilution is considered a reliable approach. However, for carrier detection and prenatal diagnosis, mutation detection should be performed. Our aim has been to study the mutational spectrum of the aspartoacylase gene in Norwegian and Swedish patients of non-Jewish origin. This knowledge is essential for patient management and for the design of rational molecular genetic diagnosis.

MATERIALS, METHODS, AND SUBJECTS

Eight Norwegian patients from six families and one Swedish patient (table 1) were identified by clinical characterisation and enzyme investigations. Clinical information was provided by the paediatricians who had followed the patients from the time of diagnosis. All the patients were boys and all parents and sibs were healthy.

The pre- and perinatal periods were unremarkable in all cases except in patient N2 where symptoms started after one week and patient N4 where hypotonia was noted from birth (table 1). Through their mother, patients N2/N3 (family 2) were related to patient N1 (family 1) and through their father they were related to patient N4 (family 3). A fourth family (family 4) could not be linked to the other three families but their ancestors originally came from the same small geographical area.

DNA studies**DNA purification**

Genomic DNA was isolated from whole blood by use of an automated DNA extractor (341 Nucleic Acid Purification Sys-

tem DNA extractor, Applied Biosystems), or by salting out methods.

cDNA synthesis

RNA was extracted from cultured skin fibroblast cells by TRI-ZOL[®]LS reagents according to the manufacturer's specifications (GIBCO, BRL). Total RNA was reverse transcribed using 100 pmol oligo dT/ μ g RNA and 200 U Super Script[™] II reverse transcriptase (GIBCO, BRL) at 50°C for 60 minutes.

PCR

Intronic PCR primers flanking each exon were used for PCR amplification and direct sequencing (table 2). All PCR reactions were carried out on a DNA Thermal Cycler/480 (Perkin-Elmer). PCR was performed in a 50 μ l volume containing 400 ng genomic DNA, 10 pmol of each primer, 1 mmol/l dNTPs, 10 $\times$ *Taq* polymerase buffer, 25 mmol/l MgCl₂, and 5U *Taq* polymerase.

DNA sequencing

The PCR products were sequenced with the appropriate primer by the use of PCR product Presequencing Kit (Amersham Life Science) and ABI PRISM BigDye[™] Termination Cycle Sequencing Kit (Perkin-Elmer). The sequences were analysed on an ABI 377 automated sequencer unit.

Key points

- Canavan disease (CD) (OMIM 271900) is an autosomal recessive neurodegenerative disorder characterised by spongy degeneration of the white matter of the brain. The disease is caused by deficiency of the enzyme aspartoacylase, encoded by the *ASPA* gene.
- The mutations E285A and Y231X comprise 98% of all mutations in CD patients in the Ashkenazi Jewish population. In non-Jewish CD patients a total of 38 different mutations have been reported of which one, A305E, is particularly prevalent as it is shared by 40-60% of the patients. The majority of patients investigated originate from southern Europe, United Kingdom, Israel, and Japan. No information exists on mutations in CD patients from Norway and Sweden.
- We report the molecular investigation of eight Norwegian and one Swedish patients deriving from seven families. Mutation analysis of these nine patients, of whom four were related, showed two novel missense mutations, 340G→T (D114Y) and 746A→T (D249V), as well as a previously reported insertion, 245insA. Survival into adulthood was found in four patients, not associated with a particular mutation.

Table 1 Clinical summary and mutations

Family	Patient	Age of onset	Alive at	Age at death	Diagnosis by	Mutation	aa change
1	N1	3 months	25 years		Brain biopsy	340G→T	D114Y
2	N2*	1 week		7 months	Brain biopsy	340G→T	D114Y
	N3	4 months		8 months	Brain biopsy	340G→T	D114Y
3	N4	Birth		6.5 months	U-NAA	340G→T	D114Y
4	N5*	3–4 months		3.5 months	Brain biopsy	340G→T	D114Y
	N6*	8–10 weeks		6.5 years	Brain biopsy	340G→T	D114Y
5	N7	3 months	21 years		U-NAA	746A→T	D249V
6	N8	3 months		17.4 years	U-NAA	245insA	Truncation
7	S9	3 months	15 years		U-NAA	746A→T	D249V

N1: previously reported by Kvittingen *et al.*¹⁶ S9: previously reported by Hagenfeldt *et al.*¹ U-NAA: urine N-acetylaspartic acid.

*Identification of the mutations was based on the genotypes of their parents (N5 and N6) or affected brother (N2).

(Perkin-Elmer) and mutations were detected by manual inspection of the sequence electropherograms.

Mutation detection by PCR and restriction enzyme analyses

In order to facilitate detection of mutations D114Y (340G→T) and D249V (746A→T), assays were developed based on PCR on genomic DNA followed by restriction enzyme digestion and agarose gel electrophoresis. PCR components were as described above. A DNA fragment embracing D114Y (340G→T) was amplified with primers pCANmF and pCANmR (table 2) at the following cycling conditions: denaturation at 95°C for five minutes followed by 36 cycles with denaturation at 94°C for 30 seconds and annealing/extension at 64°C for 90 seconds. A DNA fragment containing mutation 746A→T (D249V) was amplified with primers Caex6F and CAR 3 (table 1) under the following conditions: denaturation at 95°C for five minutes followed by 30 cycles with denaturation at 94°C for one minute, annealing at 55°C for 30 seconds, and extension at 72°C for one minute. The use of restriction enzyme *Sna*BI allowed us to discriminate between the D114 and 114Y alleles, whereas the use of *Mbo*I allowed discrimination between the D249 and 249V alleles. Restriction enzyme digestion was performed according to the manufacturer's specifications.

RESULTS

Case reports

The clinical findings were very similar for all the nine cases and typical for CD: hypotonia developing into spastic tetraplegia, macrocephaly, mental retardation, head lag, and visual impairment. As indicated in table 1, brain biopsy or raised urinary concentrations of NAA led to the diagnosis of Canavan disease. Notably, three of the patients are still alive at the ages of 25, 21, and 15 (N1, N7, and S9, respectively).

Patient N1 showed delayed development at 3 months.¹⁶ Later, he also developed epilepsy, asthma, and diabetes mellitus.

Patient N2 showed failure to thrive, irritability, and spasticity at the age of 1 week. The diagnosis was not established at the time of death, at the age of 7 months. His younger brother, born 6 years later (patient N3), died at the age of 8 months, severely mentally retarded, blind, and spastic. A brain biopsy showed spongy degeneration and Canavan disease was diagnosed and deduced also for his brother (N2).

Patient N4 showed hypotonia, failure to thrive, and irritability immediately after birth. At three months, he had hyperreflexia, spastic lower legs, and made no visual contact. Urinary NAA levels were markedly raised.

The parents of patients N5 and N6 (brothers) originate from the same small geographical area as the three previously mentioned Norwegian families (N1–N4). Cerebral CT scan

Table 2 List of primers

Primer name	Sequence	Derived from GenBank accession number
<i>cDNA primers</i>		
CAF1	5'CTGTACTTTGCCCTTTTGGTA	S67156.1
CAF3	5'CTGATTGAGCATCCTCCCTCA	S67156.1
CAR3	5'ACACCGTGTAAGATGTAAGCT	S67156.1
<i>Genomic DNA primers</i>		
CAex1F	5'CTTCTGAATTGCAGAAATCAGA	I70466
CAex1R	5'CACACCTACCACTTTACACAA	
CAex2F	5'CTCAGGCACAGATGTTGTCA	I70467
CAex2R	5'CAAGTCCTTTGCTGACTTATAA	
CAex3F	5'GGGTTTTTACCCAAGAAAGACGT	I70468
CAex3R	5'CTCTGAGTTTCAGCTAGGACAT	
CAex4F	5'ATGTGACTATCTCCTTCTGT	I70469
CAex4R	5'TCTGACCCAGGTTCCAATTGTGA	
CAex5F	5'CCAGAGATGTTTTAGTTGCCA	I70470
CAex5R	5'TGCTGTATGAGCTATAAACTTCT	
CAex6F	5'GAGTCTGACATAAATTTTAGAGGA	I70471
pCANmF	5'TTATAACAGCAAAAAAATGTACG TAG ATTGCCATATGAAGTG	
pCANmR	5'CCCATGTTAGAGGTGGTGTGTG TAC GT	

To allow the discrimination between alleles D114 and 114Y (340G→T) the primers pCANmF and pCANmR were modified in the following way. For pCANmF the nine nucleotides at the 5' end are derived from the 3' splice acceptor of intron 1.⁴ In order to create an internal *Sna*BI control restriction site nucleotides C21, A22, and A24 (in bold, numbered from the 5' end) were replaced with A, C, and T (underlined), respectively. For pCANmR, nucleotides A25 and G27 were replaced with T and C (underlined), respectively. These replacements allowed *Sna*BI digestion of the mutant 114Y allele while leaving the D114 allele resistant.

Figure 1 (A) Detection of the D114Y mutation by *Sna*BI mediated analyses of a normal control (lane 2), undigested control (lane 3), affected 114Y/114Y homozygote (lane 4), and heterozygote (lane 5). Genomic DNA was amplified and digested with *Sna*BI. Normal, undigested PCR product is 142 bp whereas normal *Sna*BI digested PCR product is 120 bp (+22 bp control fragment). Lack of the 120 bp band shows homozygosity for the mutant 114Y allele. The mutant allele is shown by the presence of the 93 bp. Lanes 1 and 7 are 100 bp ladder. (B) Detection of the D249V mutation. Analyses of a normal control (lane 2), undigested control (lane 3), affected 249V/249V homozygote (lane 4), and heterozygote (lane 5). Genomic DNA was amplified and digested with *Mbo*I. Normal, undigested PCR product is 298 bp whereas the normal digested PCR product gives rise to bands of 167 bp, 69 bp, 32 bp, and 30 bp. Lack of the 69 bp and 30 bp bands indicates homozygosity for the mutant 249V allele. The mutant allele is seen as the presence of 167 bp, 99 bp, and 32 bp. Lanes 1 and 7 are 100 bp ladder.

showed hydrocephalus in both. It was never possible to establish visual contact with patient N5 who showed pronounced head lag at 4.5 months. The steady increase of head circumference in patient N6, from 8-10 weeks, was associated with pronounced irritability and spasticity. Patient N7 showed delayed development from the age of 3 months. The clinical diagnosis was cerebral palsy until Canavan disease was confirmed biochemically at the age of 4.5 years.

The Norwegian patient N8 was diagnosed as having Canavan disease at the age of 4.5 months. He had a typical clinical presentation.

The clinical findings in the Swedish patient, S9, have been reported previously.¹ He presented delay in mental development and motor milestones from the age of 3 months, a large head circumference, and leucodystrophy on cranial CT scan at the age of 6 months. At the age of 15 years, he presents impaired vision, lack of head control, and severe mental retardation (table 1).

Mutation analyses

The study included nine Canavan patients from seven families, of whom three were interrelated. DNA samples from six patients were available for mutation analysis. In three cases, identification of the mutations was based on the genotypes of their parents (N5 and N6) or affected brother (N2). Brief clinical descriptions and genotypes of the Norwegian and Swedish CD patients are listed in table 1.

Two novel missense mutations were identified or deduced in the six probands. In addition, a 1 bp insertion, 245insA,⁶ was identified in one patient.

Four probands were homozygous for nucleotide substitution 340G→T in exon 2, resulting in the replacement of Asp with Tyr at amino acid position 114 (D114Y) (results not shown). Homozygosity for nucleotide substitution 746A→T was identified in one Norwegian as well as in the only Swedish patient included in this study (results not shown). This particular point mutation results in the substitution of Asp with Val at amino acid position 249, D249V, of the 313 residue ASPA polypeptide. The nucleotide substitution 746A→T resides in exon 6 in position +2 next to the intron 5/exon 6 border. Thus, the possibility existed that this mutation might cause splicing deficiency. However, RT-PCR combined with cDNA sequencing detected only normal splicing (results not

shown). Hence, we conclude that 746A→T causes an amino acid replacement only.

PCR based diagnostic tests were developed for both 340G→T (D114Y) and 746A→T (D249V) (fig 1). Neither D114Y nor D249V were found in a panel of 100 Norwegian control alleles (data not shown).

The Norwegian patient N7 was found to be homozygous for the mutation 245insA. This particular mutation in exon 1 causes a shift in the translational reading frame and, hence, a truncated gene product.⁶

DISCUSSION

The majority of non-Jewish Canavan patients investigated originate from southern Europe, United Kingdom, Israel, and Japan. In this work we present brief clinical data and the identification of two novel mutations in Norwegian and Swedish patients with Canavan disease.

The mutation D114Y (340G→T) was found in three related families and in a fourth family originating from the same small geographical region in Norway. It is located in exon 2, resulting in the replacement of the charged, acidic side chain of aspartic acid with the hydrophobic, bulky, aromatic side chain of tyrosine. Hence, the severe consequences of such an alteration can easily be envisaged. Moreover, another amino acid substitution at this particular residue has previously been reported: aspartic acid to glutamic acid (D114E), resulting from 342C→A in a patient of Turkish origin.¹⁷ Such a replacement of Asp with Glu, which conserves the charge and polarity of the side chain but extends the length by only one single methyl group, is apparently sufficient to abolish ASPA activity which was only 0.35% of residual activity.¹⁷ This further shows the importance of residue 114 for the structure/function of the ASPA enzyme.

The mutation D249V (746A→T) was found in one Norwegian and the Swedish patient. The replacement of Asp with Val at amino acid position 249 represents a substitution of a charged, polar side group with an aliphatic, hydrophobic, non-polar side group. The difference in the chemical properties of Asp and Val is consistent with the notion that D249V is abolishing ASPA activity and, hence, is disease causing. Since neither D114Y nor D249V have been reported elsewhere, it is likely that these mutations are specific to Canavan patients from the Scandinavian peninsula.

The mutation, 245insA, a frameshift mutation resulting in a truncated, non-functional gene product, was found in a family from western Norway. This mutation was previously identified in an Italian patient.⁶ It is located in a run of seven adenines, a motif previously documented to be prone to misincorporation of an extra base by the DNA polymerase as a result of slipped mispairing at the replication fork.¹⁸

Both D114 and D249 are conserved in the *ASPA* genes of cow (93% identity), mouse (86% identity), and, strikingly, also in *Cyanobacteria* (46% identity). In the *ASPA* gene of rat (86% identity), however, D249 is conserved whereas D114 is replaced with valine. The D114Y, D249V, and 245insA were the only mutations found in these patients. The fact that they were absent in 100 normal control alleles strongly indicates the pathogenicity of these mutations.

Clinical data showed survival into adulthood in four out of six Norwegian/Swedish patients (table 1). However, since both 114Y/114Y and 249V/249V homozygotes show variable survival time, no genotype-phenotype correlation was evident. Lack of genotype-phenotype correlation has also been previously reported for other CD mutations.¹⁹ It remains to be elucidated how environmental and epigenetic factors may influence the clinical course of Canavan disease. Few data exist concerning residual aspartoacylase activity in CD patients. Such information is crucial to clarify a possible link between genotype, enzyme activity, and clinical phenotype.¹⁷

Our findings add to the spectrum of mutations in Canavan disease and are essential to the genetic diagnosis and counselling of CD families of Norwegian and Swedish origin. The location and nature of the mutations may contribute to further understanding of the function and structure of the *ASPA* gene product.

ACKNOWLEDGEMENTS

The present work was financially supported by The Norwegian Research Council (NFR project No 114575/320), Samordningsrådet for forskningsfondene (SOR), and The Norwegian Research Foundation for Mental Deficiency-Norwegian Red Cross. We are grateful to the participating families and to Drs Dag Veimo, Nordland Central Hospital, Norway, Sigurd Børsting, Innherred Hospital, Levanger, Norway, and Per Erik Waaler, Haukeland University Hospital, Bergen, Norway, for the referral of patients and valuable clinical contributions. We are also grateful to Helle Klenow and Karina Pedersen, University Hospital of Northern Norway, for their technical assistance.

Authors' affiliations

T R Olsen, L Tranebjærg*, Ø Nilssen, Department of Medical Genetics, University Hospital of Northern Norway, N-9038 Tromsø, Norway
E I Kvittingen, Department of Clinical Biochemistry, University Hospital, Rikshospitalet, Oslo, Norway
L Hagenfeldt, Centre for Inherited Metabolic Diseases, Huddinge University Hospital, S-14186 Stockholm, Sweden
C Møller, Department of Paediatrics, Haukeland University Hospital,

Bergen, Norway

Correspondence to: Dr Ø Nilssen, Department of Medical Genetics, University Hospital of Northern Norway, N-9038 Tromsø, Norway; oivindn@fagmed.uit.no

*Present address: Department of Audiology, Bispebjerg Hospital, H:S, and Institute of Medicine, Biochemistry and Genetics, IMBG, University of Copenhagen, Denmark

REFERENCES

- 1 **Hagenfeldt L**, Bollgren I, Venizelos N. N-acetylaspartic aciduria due to aspartoacylase deficiency B, a new aetiology of childhood leukodystrophy. *J Inher Metab Dis* 1987;**10**:135-41.
- 2 **Matalon R**, Michals K, Sebesta D, Deanching M, Gashkoff P, Casanova J. Aspartoacylase deficiency and N-acetylaspartic aciduria in patients with Canavan disease. *Am J Med Genet* 1988;**29**:463-71.
- 3 **Kaul R**, Gao GP, Balamurugan K, Matalon R. Cloning of the human aspartoacylase cDNA and a common missense mutation in Canavan disease. *Nat Genet* 1993;**5**:118-23.
- 4 **Kaul R**, Balamurugan K, Gao PG, Matalon R. Canavan disease: genomic organization and localization of human *ASPA* to 17p13-ter and conservation of the *ASPA* gene during evolution. *Genomics* 1994;**21**:364-70.
- 5 **Kaul R**, Gao GP, Aloya M, Balamurugan K, Petrosky A, Michals K, Matalon R. Canavan disease: mutations among Jewish and non-Jewish patients. *Am J Hum Genet* 1994;**55**:34-41.
- 6 **Elpeleg ON**, Shaah A. The spectrum of mutations of the aspartoacylase gene in Canavan disease in non-Jewish patients. *J Inher Metab Dis* 1999;**22**:531-4.
- 7 **The Human Gene Mutation Data Base Cardiff**, UK. <http://www.uwcm.ac.uk/uwcm/mg/hgmd0.html>
- 8 **Rady PL**, Vargas T, Tyring SK, Matalon R, Langenbeck U. Novel missense mutation (Y231C) in a Turkish patient with Canavan disease. *Am J Med Genet* 1999;**87**:273-5.
- 9 **Rady PL**, Penzien JM, Vargas T, Tyring SK, Matalon R. Novel splice site mutation of aspartoacylase gene in a Turkish patient with Canavan disease. *Eur J Pediatr Neurol* 2000;**4**:27-30.
- 10 **Tahmaz FE**, Sam S, Hoganson GE, Quan F. A partial deletion of the aspartoacylase gene is the cause of Canavan disease in a family from Mexico. *J Med Genet* 2001;**38**:e9.
- 11 **Jakobs C**, Brink Ten, HJ, Langelaar SA, Zee T, Stellaard F, Macek M, Srsnova K, Srsen S, Kleijer WJ. Stable isotope dilution analysis of N-acetylaspartic acid in CSF, blood, urine and amniotic fluid: accurate postnatal diagnosis and the potential for prenatal diagnosis of Canavan disease. *J Inher Metab Dis* 1991;**4**:653-60.
- 12 **Bennett MJ**, Gibson KM, Sherwood WG, Divry P, Rolland MO, Elpeleg ON, Rinaldo P, Jakobs C. Reliable prenatal diagnosis of Canavan disease aspartoacylase deficiency: comparison of enzymatic and metabolite analysis. *J Inher Metab Dis* 1993;**16**:831-6.
- 13 **Matalon R**, Michals K, Kaul R. Canavan disease: from spongy degeneration to molecular analysis. *J Pediatr* 1995;**127**:511-17.
- 14 **Matalon R**, Michaels-Matalon K. Prenatal diagnosis of CD. *Prenat Diagn* 1999;**19**:669-70.
- 15 **Besley GTN**, Elpeleg ON, Shaag A, Manning NJ, Jakobs C, Walter JW. Prenatal diagnosis of the Canavan disease; problems and dilemmas. *J Inher Metab Dis* 1999;**21**:48.
- 16 **Kvittingen EA**, Guldal G, Børsting S, Skälpe IO, Stokke O, Jellum E. N-acetylaspartic aciduria in a child with a progressive cerebral atrophy. *Clin Chim Acta* 1986;**158**:217-27.
- 17 **Kaul R**, Gao GP, Balamurugan K, Petrosky A, Michals K, Matalon R. Identification and expression of eight novel mutations among non-Jewish patients with Canavan disease. *Am J Hum Genet* 1996;**59**:95-102.
- 18 **Cooper DN**, Krawczak M. Mechanisms of insertional mutagenesis in human genes causing genetic disease. *Hum Genet* 1991;**87**:409-15.
- 19 **Matalon R**, Michaels-Matalon K. Recent advances in Canavan disease. *Adv Pediatr* 1999;**46**:493-506.