

## ORIGINAL ARTICLE

## Clinical aspects, prenatal diagnosis, and pathogenesis of trisomy 16 mosaicism

P J Yong, I J Barrett, D K Kalousek, W P Robinson

*J Med Genet* 2003;40:175–182

See end of article for authors' affiliations

Correspondence to:  
Dr W P Robinson,  
Department of Medical  
Genetics, University of  
British Columbia, British  
Columbia Research Institute  
for Children's and  
Women's Health,  
950 West 28th Avenue,  
Vancouver, British  
Columbia, Canada  
V5Z 4H4; wprobins@  
interchange.ubc.ca

Revised version received  
10 November 2002  
Accepted for publication  
11 November 2002

**Introduction:** Analysis of data from cases of trisomy mosaicism can provide insight for genetic counselling after prenatal diagnosis and for the elucidation of the pathogenesis of trisomy during pregnancy.

**Methods:** Statistical analysis was carried out on data from 162 cases of pregnancies with prenatal diagnosis of trisomy 16 mosaicism.

**Results:** The majority of cases resulted in live birth (66%) with an average gestational age of 35.7 weeks and average birth weight of  $-1.93$  standard deviations from the population mean. Among the live births 45% had at least one malformation, the most common being VSD, ASD, and hypospadias. The level of trisomy on direct CVS (cytotrophoblast) was associated with more severe intrauterine growth restriction (IUGR) and higher risk of malformation, while the level of trisomy on cultured CVS (chorionic villous stroma) was associated only with more severe IUGR. Similarly, the presence of trisomy on amniocentesis (amniotic fluid) was associated with both IUGR and malformation, while the presence of trisomy in the amniotic mesenchyme was associated only with IUGR. Surprisingly, the degree of trisomy in placental tissues appeared to be independent of the degree of trisomy in amniotic fluid and amniotic mesenchyme. The sex of the fetus was not associated with any outcome variables, although there was an excess of females (sex ratio = 0.45) that may be explained by selection against male trisomy 16 embryos before the time of CVS ( $-9-12$  weeks).

**Conclusion:** The levels of trisomy in different fetal-placental tissues are significant predictors of some measures of outcome in mosaic trisomy 16 pregnancies.

Mosaicism for trisomy at prenatal diagnosis presents a major dilemma for genetic counselling. Case reports and series have shown outcomes ranging from late spontaneous abortion to seemingly normal live birth. Consolidation and statistical analysis of data from these reports can improve counselling by identifying key correlates with outcome. Such analysis can also provide insight into the pathogenic mechanisms involved in trisomy mosaicism during pregnancy.

Trisomy 16 is of particular importance as it is thought to be the most frequent chromosome abnormality at conception.<sup>1</sup> Among all clinically recognised pregnancies, it has an incidence of  $\sim 1.5\%$ .<sup>2</sup> Although most trisomy 16 embryos are spontaneously aborted or are noted to have arrested development between 8–15 weeks of gestation, some embryos survive and are candidates for prenatal diagnosis.<sup>3</sup> Approximately 34 per 100 000 chorionic villus sampling (CVS) analyses detect trisomy 16,<sup>1</sup> while a recent estimate for amniocentesis has not been reported.<sup>3</sup> These surviving embryos are virtually always mosaic (that is, contain both euploid and trisomic cell lines), with only one apparently non-mosaic case diagnosed at CVS<sup>4</sup> and two apparently non-mosaic cases that were diagnosed at necropsy after intrauterine death in the second and third trimesters.<sup>5,6</sup> Since the trisomic cells must be completely or at least predominantly confined to the placenta for a mosaic trisomy 16 conceptus to survive, the term “confined placental mosaicism” (CPM) can also be used to describe trisomy 16 mosaicism. CPM of trisomy 16 may or may not exhibit low level trisomy in the fetus depending on whether the trisomy is predominantly or completely confined to the placenta, respectively.

Almost all mosaic trisomy 16 pregnancies originate from a trisomy 16 zygote as a consequence of a maternal meiosis I non-disjunction.<sup>7</sup> Thus, trisomy 16 mosaicism is normally caused by “trisomy rescue”, whereby loss of a chromosome 16

in one of the cells of the early trisomic embryo results in a euploid cell line. The eventual distribution of trisomy 16 cells in the placenta and fetus depend on the embryonic stage when trisomy rescue occurs, natural selection against trisomic cells, and stochastic processes during development.

When a trisomy 16 conceptus is rescued, one of the two maternal chromosomes or the paternal chromosome can be lost. If the former occurs, the result is biparental disomy 16 (BPD(16)) or a chromosome 16 inherited from each parent; if the latter occurs, the result is maternal uniparental disomy 16 (UPD(16)mat) or both chromosomes inherited from the mother.<sup>8,9</sup> Uniparental disomy (UPD) may have a distinct phenotypic effect if imprinted genes (that is, genes whose expression depends on whether they are inherited from the mother or father) exist on chromosome 16. We have previously shown that UPD(16)mat is likely to have a subtle phenotypic effect based on statistical analysis of 83 cases and therefore should be considered one of the pathogenic mechanisms of trisomy 16 during pregnancy.<sup>10</sup>

Besides UPD, other factors potentially contributing to the pathogenesis of trisomy 16 mosaicism are (1) the degree of trisomy in various tissues of the placenta and fetal membranes; (2) the degree and distribution of trisomy in tissues of the fetus; and (3) the sex of the fetus. In this paper, statistical analysis was performed on data from a large series ( $n=162$ ) of published and unpublished mosaic trisomy 16 cases with the purpose of (1) summarising the clinical outcome of mosaic trisomy 16 pregnancies; and (2) evaluating the predictive value of various factors for measures of pregnancy outcome. The identification of important predictive factors will aid genetic counselling after prenatal diagnosis and the elucidation of which factors other than UPD are involved in the pathogenesis of trisomy 16 mosaicism during pregnancy.

## METHODS

The study sample ( $n=162$  cases) consists of mosaic trisomy 16 pregnancies diagnosed prenatally by CVS or amniocentesis (with or without molecular (PCR) testing for the UPD status of chromosome 16). Rare cases with paternal origin of the trisomy ( $n=2$ ), partial trisomy ( $n=3$ ), and concomitant aneuploidy ( $n=1$ ) were excluded in order not to confound the analysis (Electronic Appendix 1). Some cases are from a continuing study of trisomy mosaicism at the University of British Columbia (UBC) ( $n=58$ ). This study consists of cases referred from other centres ( $n=51$ ) and cases initially ascertained at the Children's and Women's Health Centre of British Columbia ( $n=7$ ). Some data from most of these cases have been published previously, and there is overlap with cases published by other research groups (Electronic Appendix 1). The Vancouver study was approved by the ethics committee of the University of British Columbia. An additional 103 cases are from other published reports to date (Electronic Appendix 1). Using the review of mosaic trisomy 16 cases by Benn<sup>3</sup> as a starting point, data were verified from the original sources and care was taken to eliminate duplicated cases. In general, cases were ascertained via mothers undergoing prenatal testing for advanced maternal age, abnormal triple screen, and anomalies or growth restriction noted on ultrasound.

Data were collected on the following variables: (1) pregnancy outcome (live birth, intrauterine death, or termination of pregnancy); (2) gestational age at pregnancy outcome; (3) malformation detected in the fetus/neonate/infant ("malformation" used as general term independent of aetiology, including possible disruptions and deformations); (4) fetal/neonatal weight at pregnancy outcome; (5) percent trisomy on amniocentesis (assessing amniotic fluid, thought to be representative of various fetal tissues); (6) percent trisomy on (semi-) direct CVS (assessing the cytotrophoblast of the placenta); (7) percent trisomy on cultured CVS (assessing the chorionic villous stroma of the placenta); (8) percent trisomy in the chorionic membrane (part of the fetal membranes) or chorionic plate of the placenta (together referred to as "chorion" for simplicity) at postnatal or necropsy examination; (9) percent trisomy in the chorionic villous stroma at postnatal or necropsy examination; (10) percent trisomy in the trophoblast at postnatal or necropsy examination (since postnatal or necropsy examination usually involved FISH without previous culture, both cytotrophoblast and syncytiotrophoblast nuclei were included; thus the general term "trophoblast" is used); (11) percent trisomy in the amniotic mesenchyme at postnatal or necropsy examination of the amnion (part of the fetal membranes) ("amniotic mesenchyme" refers to the layer of the amnion that is studied when the amnion is cultured before analysis<sup>13</sup>); and (12) confirmation of trisomy in various fetal tissues. Since FISH was used to confirm the results from conventional cytogenetics only very rarely ( $n=4$ ) for CVS and amniocentesis, only data from conventional cytogenetics were used for variables (5), (6), and (7). As FISH was more common during postnatal or necropsy investigation of the placenta, FISH was given precedence over conventional cytogenetics for variables (8), (9), and (10) because FISH samples a greater number of cells.<sup>7</sup> Only data from conventional cytogenetics was considered for variable (11). For variable (12), data from conventional cytogenetics, FISH, or molecular methods (PCR) were taken into account; if results were contradictory (that is, one was positive for trisomy, the other negative), then the fetal tissue was coded as positive. Molecular detection of trisomy in fetal tissues was only considered for cases from the UBC study where detailed information on chromosome 16 markers was available, and cases from other published reports where there was an explicit statement that PCR showed or excluded trisomy.

It should be emphasised that the study sample may be biased towards cases with poorer outcomes, since such cases


are more likely to be ascertained (for example, because of anomalies observed on ultrasound), referred for research purposes, and/or submitted for publication. Therefore, purely descriptive statistics should not be considered estimates of the actual values in the general population, but are intended as descriptions of the study sample specifically. Statistical associations between variables are more likely to be unbiased, but any potential biases should be considered when interpreting results. For example, postnatal or necropsy examination of the amniotic mesenchyme and fetal tissues may have been more thorough in pregnancies with abnormal outcome or with trisomy detected on amniocentesis. Furthermore, it should be noted that there was variation in the quantity and quality of data available among the cases (for example, in the descriptions of malformations).

Associations between variables were tested for statistically (for example,  $t$  test, Fisher exact test, Yates chi-square test) using SPSS 10.0 and the VassarStats Web Site for Statistical Computation (<http://faculty.vassar.edu/lowry/VassarStats.html>). All tests were one tailed (except for the Yates chi-square test). Data were grouped or categorised for analysis to maximise power (for instance, the percent trisomy on CVS was grouped into 100% and <100%). As an example, 50% and 60% trisomy on CVS will be detected as a true difference on a non-grouped analysis, but may be biologically equivalent because the level of trisomy on CVS is affected by stochastic processes (such as the area of the placenta that is sampled and culture of the sample before cytogenetics).


## RESULTS

### Clinical outcome of prenatally diagnosed mosaic trisomy 16 pregnancies

Even with an expected bias towards poor outcome, the majority of cases (66%) resulted in live births, of which 93% survived beyond the neonatal period. Eleven percent of pregnancies ended in intrauterine death (IUD), while 22% of the pregnancies were terminated. Results are for 157 cases informative for pregnancy outcome. Fig 1 shows the distribution of gestational ages for the live births (including both those that survived beyond the neonatal period and neonatal deaths); the average gestational age was 35.7 (0.41) (mean (SE)) weeks. The distribution of birth weights (number of SDs



**Figure 1** Histogram of gestational ages for live births, including seven cases that resulted in neonatal death and 70 cases that survived beyond the neonatal period.



**Figure 2** Histogram of birth weights (number of SDs from the gestational age corrected general population mean birth weights<sup>14</sup>) for live births (including both those that survived beyond the neonatal period and neonatal deaths). Results are for 72 informative cases.

from gestational age corrected general population mean birth weights<sup>14</sup>) for the live births is illustrated in fig 2. Virtually all birth weights (90%) were below the general population mean birth weights (that is, 0 standard deviations), with an average birth weight of  $-1.92$  ( $0.16$ ) (mean (SE)) SD from the mean.

Considering all cases regardless of pregnancy outcome, at least one malformation was present in 49% of the 129 cases where malformation status was known. Malformations were classified anatomically into cardiac, pulmonary, genitourinary, gastrointestinal, craniofacial, extra-craniofacial musculo-skeletal, and miscellaneous dermatological. Cardiac malformations were most prevalent (62% of cases positive for malformation), with the other malformation classes present in  $\sim 20$ –45% of positive cases.

Considering individual types of malformations, almost all were present in three or fewer cases in the study sample (of 129 cases informative for malformation status), and were considered sporadic or random. Several malformations were present in eight or more cases (that is, in  $>6\%$  of informative cases, equivalent to at least twice the risk of major structural anomalies in the general population at birth ( $2$ – $3\%$ )<sup>15</sup>), and thus were considered more likely to have a true association with trisomy 16 mosaicism: VSD 17.8% ( $n=23$ ), two vessel cord 10.9% ( $n=14$ ), ASD 7.8% ( $n=10$ ), clinodactyly 7.8% ( $n=10$ ), and pulmonary hypoplasia 7.0% ( $n=9$ ). It should be noted that at least five of the cases of pulmonary hypoplasia were probably secondary to other malformations (for example, diaphragmatic hernia, renal anomalies, premature rup-

ture of membranes, severe scoliosis, and omphalocele with kyphoscoliosis).

Considering only live births, 45% of the 92 cases informative for malformation status had at least one malformation. Some malformations were present in six or more cases (that is, in  $>6\%$  of informative cases) and were considered more likely to have a true association with trisomy 16 mosaicism: VSD 16.3% ( $n=15$ ), ASD 9.8% ( $n=9$ ), and hypospadias 26.9% ( $n=7$ , of 26 informative male cases). All of these malformations were significantly more frequent when compared to the corresponding prevalences among neonates in the general population (Electronic Appendix 2,  $p < 0.0001$ ).

### Trisomy 16 in the placenta

Since most trisomy 16 cases show 100% trisomy in both direct CVS (cytotrophoblast) and cultured CVS (chorionic villous stroma),<sup>16</sup> more meaningful binary “indicator” variables, referred to as dCVS and cCVS respectively, were set up: the presence of euploidy ( $<100\%$  trisomy) was coded as “0”, while the presence of non-mosaic trisomy (100% trisomy) was coded as “1”. Associations between dCVS and cCVS, and four outcome variables of interest (table 1) were then tested for. Non-mosaic trisomy on direct CVS (cytotrophoblast) (dCVS=1) was associated with lower fetal/neonatal weight in terms of the number of SDs from the gestational age corrected general population mean birth weights<sup>14</sup> (BW) ( $t=2.22$ ,  $df=n-2 = 36$ ,  $p = 0.016$ ; table 2) and higher risk of malformation (MALF) (Fisher exact test,  $n=50$ ,  $p=0.015$ ,  $RR=\infty$ ; table 3), but was not associated with the risk of intra-uterine or neonatal death (OUT) (Fisher exact test,  $n=51$ ,  $p=0.589$ ) or the gestational age at live birth or intrauterine death (GEST) ( $t=0.67$ ,  $df=41$ ,  $p=0.255$ ). In contrast, non-mosaic trisomy on cultured CVS (chorionic villous stroma) (cCVS=1) was associated with only lower BW ( $t=1.91$   $df=34$ ,  $p=0.032$ ; table 2), and less significantly so compared to non-mosaic trisomy on direct CVS (cytotrophoblast).

Statistical analysis involving postnatal or necropsy investigation of the different placental tissues (trophoblast, chorionic villous stroma, and chorion) should be viewed more cautiously since the number of informative cases for any given outcome was small ( $n=15$ ). To determine whether there is potentially selection against trisomic cells during development, the percent trisomy in cultured and direct CVS, and at postnatal or necropsy assessment of the placental tissues, were coded so that “0” refers to 0%, “1” refers to 1%–10%, “2” refers to 11%–20%, etc. For those cases with both cultured CVS and postnatal or necropsy examination of the chorionic villous stroma, the chorionic villous stroma had significantly lower levels of trisomy: 6.05 (0.79) (mean (SE))  $v$  9.23 (0.39) (mean (SE)) (paired sample  $t=3.89$ ,  $df=21$ ,  $p=0.0004$ ). For those cases with both direct CVS and postnatal or necropsy examination of the trophoblast, there was a trend towards a lower level of trisomy at postnatal or necropsy examination of the trophoblast: 7.75 (0.90) (mean (SE)) versus 9.42 (0.58) (mean (SE)) (paired sample  $t=1.62$ ,  $df=11$ ,  $p=0.067$ ). It should be noted that the latter may not be significant because the paired sample size ( $n=df+1=12$ ) was smaller than the

**Table 1** Outcome variables of interest

Variable	Value	Explanation
OUT*	1	Intrauterine or neonatal death
	0	Live birth with survival beyond the neonatal period
GEST*		Gestational age at live birth or intrauterine death
BW†		Weight of the fetus/neonate in terms of the number of SDs from gestational age corrected general population mean birth weights <sup>14</sup>
MALF†	1	Presence of malformation in the fetus/neonate/infant
	0	Absence of malformation in the fetus/neonate/infant

\*Excluding cases with termination of pregnancy.

†Including all cases regardless of pregnancy outcome (live birth, intrauterine death, or termination of pregnancy).

**Table 2** Associations with BW\*

Variable	Tissue	Procedure	Value	Level of trisomy	No	Mean BW*	SD	t statistic, df, and p value
dCVS	Cytotrophoblast	Direct CVS	1	100%	34	-2.05	1.19	$t = 2.22, df = 36, p = 0.016$
			0	<100%	4	-0.52	2.20	
cCVS	Chorionic villous stroma	Cultured CVS	1	100%	31	-1.80	1.10	$t = 1.91, df = 34, p = 0.032$
			0	<100%	5	-0.71	1.70	
AF	Amniotic fluid	Amniocentesis	1	>0%	22	-2.82	1.07	$t = 3.55, df = 68, p = 0.0003$
			0	0%	48	-1.67	1.33	
AMN	Amniotic mesenchyme	Postnatal or necropsy sampling of amnion followed by culture	1	>0%	7	-3.13	0.87	$t = 2.66, df = 16, p = 0.009$
			0	0%	11	-1.48	1.48	

\*Weight of the fetus/neonate in terms of the number of SDs from gestational age corrected general population mean birth weights.<sup>14</sup>

**Table 3** Associations with risk of malformation

Variable	Tissue	Procedure	Value	Level of trisomy	MALF		Risk	Statistic, RR and p value
					Absence of malformation	Presence of malformation		
dCVS	Cytotrophoblast	Direct CVS	1	100%	23	19	45%	Fisher exact test, RR = ∞, p = 0.015
			0	<100%	8	0	0%	
AF	Amniotic fluid	Amniocentesis	1	>0%	15	37	71%	Yates chi-square = 13.90, RR = 2.10, p = 0.0002
			0	0%	39	20	34%	

paired sample size for the comparison between cultured CVS and chorionic villous stroma ( $n = df + 1 = 22$ ).

### Trisomy 16 in the amniotic mesenchyme

Since about half of postnatal or necropsy investigations of the amniotic mesenchyme (that is, sampling of the amnion followed by culture before analysis) yielded a euploid karyotype, percent trisomy in the amniotic mesenchyme was coded into an indicator variable AMN: "0" refers to the presence of non-mosaic euploidy (0% trisomy) and "1" refers to the presence of trisomy (>0% trisomy). The presence of trisomy in the amniotic mesenchyme (AMN=1) was only associated with lower BW (Welch's approximate  $t = 2.98, df = 15, p < 0.005$ ; table 2).

### Trisomy 16 in the fetus

For the cases where percent trisomy on amniocentesis (in amniotic fluid) was known, about half also had a non-mosaic euploid karyotype. Therefore, another indicator variable (AF) was set up; the presence of non-mosaic euploidy (0% trisomy) on amniocentesis was coded as "0", while the presence of trisomy (>0% trisomy) was coded as "1". The presence of trisomy on amniocentesis (AF=1) was associated with lower BW ( $t = 3.55, df = 68, p = 0.0003$ ; table 2) and higher risk of

malformation (Yates chi-square = 13.90,  $n = 111, p = 0.0002, RR = 2.10$ ; table 3). Removing cases with AF=0 (that is, considering only cases where the amniotic fluid is positive for trisomy), the level of trisomy in amniotic fluid (coded so that "1" refers to 1%-10%, "2" refers to 11%-20%, etc) was not associated with BW ( $r = 0.13, n = 20, p = 0.297$ ) or malformation ( $t = 0.58, df = 43, p = 0.282$ ). Therefore, the presence of trisomy in amniotic fluid, but not the actual level above 0%, is predictive of outcome. It should be noted, however, that there was considerable variability in outcomes even for a given result on amniocentesis. Table 2 shows that there is a large standard deviation for the BW distribution for cases with trisomy present on amniocentesis (SD 1.07), and for the BW distribution for cases with no trisomy present on amniocentesis (SD 1.33). In addition, table 3 shows that although the risk of malformation increases when trisomy is present on amniocentesis, the risk of malformation is still considerable (34%) even when no trisomy is detected on amniocentesis.

In addition, the presence of trisomy on amniocentesis (AF=1) was also associated with the presence of trisomy in amniotic mesenchyme (AMN=1) (Fisher exact test,  $n = 21, 7.70 p = 0.007, RR = 7.70$ ; table 4). However, there were four discordant cases, with three cases being positive for trisomy in amniotic fluid but negative in the amniotic mesenchyme, and

**Table 4** Associations between trisomy on amniocentesis and trisomy in amniotic mesenchyme and fetal/neonatal/infant tissues

Variable	Tissue	Procedure	Value	Level of trisomy	AF		RR and p value*
					0% trisomy on amniocentesis	>0% trisomy on amniocentesis	
AMN	Amniotic mesenchyme	Postnatal or necropsy sampling of amnion followed by culture	1	>0%	1	7	RR = 7.70, p = 0.007
			0	0%	10	3	
FETUS	Fetal/neonatal/infant tissues	Postnatal or necropsy sampling	1	>0%	3	17	RR = 4.10, p = 0.004
			0	0%	31	30	
BLOOD	Fetal/neonatal/infant blood	Cordocentesis or postnatal or necropsy sampling	1	>0%	1	3	RR = 2.40, p = 0.396
			0	0%	27	32	

\*RR calculated with AF as "exposure" variable from which AMN, FETUS, and BLOOD were ascertained, and p value derived from Fisher exact test.

one case being negative in amniotic fluid but positive in the amniotic mesenchyme. Moreover, a variety of fetal tissues were examined for trisomy at postnatal or necropsy examination, including blood (n=73) and skin (n=45) as well as a number of other tissues (kidney, lung, liver, brain, heart, gonad, spleen, adrenal, thymus, intestine, buccal, cartilage, connective tissue, diaphragm, eye, fascia, muscle, tendon, and rectum) that were examined less often (all n<10 except for kidney (n=16) and lung (n=15)). The presence of trisomy in tissues of the fetus/neonate/infant detected at postnatal or necropsy examination was coded into an indicator variable FETUS: "0" refers to non-mosaic euploidy (0% trisomy) in all tissues examined and "1" refers to the presence of trisomy (>0% trisomy) in at least one tissue. The presence of trisomy in at least one fetal tissue (FETUS=1) was not associated with the presence of trisomy in the amniotic mesenchyme (AMN=1) (Fisher exact test, n=23, p=0.596), but was associated with the presence of trisomy on amniocentesis (AF=1) (Fisher exact test, n=81, p=0.004, RR=4.10; table 4). Again, it should be noted that there were 33 discordant cases, with 30 cases being positive for trisomy in amniotic fluid but negative in fetal/neonatal/infant tissues, and three cases being negative in amniotic fluid but positive in fetal/neonatal/infant tissues.

Since trisomy in fetal blood can be tested for prenatally (cordocentesis), the clinical value of trisomy detected in blood lymphocytes was of interest. As was already established,<sup>3</sup> trisomy 16 was rarely found in blood (trisomy was detected in only 5.5% of cases where the blood was tested for trisomy via cordocentesis, or postnatal or necropsy sampling), and so cordocentesis is not of prenatal diagnostic value in this context. The presence of trisomy in the blood (coded into an indicator variable BLOOD) was not significantly associated with the presence of trisomy on amniocentesis (AF=1) (Fisher exact test, n=63, p=0.396, RR=2.40; table 4), although this result is probably influenced by the small number of cases positive for trisomy in blood.

### Independence of variables

To determine whether the effects of the variables (dCVS, cCVS, AF, and AMN) on outcome are independent, ideally regression modelling should be used. For instance, dCVS and AF may be positively correlated; that is, cases with non-mosaic trisomy on direct CVS (dCVS=1) may be more likely to have the presence of trisomy on amniocentesis (AF=1). However, sample sizes were too small so that significance was lost even in simple regression models between one explanatory variable and one outcome variable when only cases informative for a potentially confounding variable were considered. Therefore, in lieu of regression modelling, pairwise associations between the variables were determined. dCVS (level of trisomy in the cytotrophoblast determined by direct CVS) and cCVS (level of trisomy in the chorionic villous stroma determined by cultured CVS) were associated (that is, were not independent) (Fisher exact test, n=35, p=0.002, RR=5.17). As noted previously, AF (presence of trisomy in the amniotic fluid determined by amniocentesis) and AMN (presence of trisomy in the amniotic mesenchyme determined by postnatal or necropsy sampling followed by culture) were not independent (Fisher exact test, n=21, p=0.007, RR=7.70). In contrast, both dCVS (Fisher exact test, n=42, p=0.557) and cCVS (Fisher exact test, n=43, p=0.523) were not associated with (that is, were independent of) AF. Similarly, both dCVS (Fisher exact test, n=10, p=0.133) and cCVS (Fisher exact test, n=13, p=0.577) were independent of AMN. Although the sample sizes for the latter associations were small (n=10 and n=13), there were no clear trends, which is consistent with the lack of association between dCVS/cCVS and AF. Electronic Appendix 3 summarises which variables were (and were not) independent of each other.

### Sex of the fetus in trisomy 16 mosaicism


The sex ratio of informative cases in the study sample was 0.45 (45 males and 101 females), which is significantly different from the expected ratio of 1.07 calculated from prenatal controls by Huether *et al*<sup>17</sup> (Yates chi-square = 23.63, df=1, p<0.0001). This sex ratio bias towards females confirms a previous observation made by Benn.<sup>3</sup> Since a significant difference was found between the sex ratios found at CVS (1.28) and amniocentesis (1.06) in the prenatal controls (the average of the two, weighted by sample size, being 1.07),<sup>17</sup> comparisons were made separately with the sex ratio for cases with CVS and the sex ratio for cases with amniocentesis in the study sample. The sex ratio at CVS was found to be 0.41 (31 males and 76 females), which was significantly different from the expected 1.28 (Yates chi-square = 30.98, df=1, p<0.0001). The sex ratio at amniocentesis was 0.43 (33 males and 76 females), which was significantly different from the expected 1.06 (Yates chi-square = 18.74, df=1, p<0.0001).

To determine when the bias in sex ratio occurs, the sex of the fetus was coded into an indicator variable SEX, where SEX = 0 and SEX = 1 refer to female and male respectively. Sex of the fetus was not found to be significantly associated with any of the outcome variables, also confirming previous observations.<sup>3</sup> Since the cases in the study sample are of trisomy 16 fetuses that survived beyond the time of amniocentesis (with few exceptions, cases of early terminations or intrauterine deaths), this suggests that there is little or no selection against a particular sex after the time of amniocentesis (on average, ~17 weeks in the study sample). Furthermore, the sex ratios at CVS (0.41) and amniocentesis (0.43) were not significantly different (Yates chi-square = 0, df=1, p=1.00). This suggests that no selection against male fetuses occurs between the times of CVS (~9–12 weeks) and amniocentesis (~17 weeks), assuming there is not a significant number of mosaic trisomy 16 pregnancies diagnosed at CVS that spontaneously abort soon after and thus are not referred or published. In contrast, the sex ratio at CVS was significantly lower than the sex ratio of non-mosaic trisomy 16 spontaneous abortions (1.01)<sup>18</sup> (Yates chi-square = 18.53, df=1, p<0.0001), suggesting that the bias in sex ratio is set in the first trimester before ~9–12 weeks.

## DISCUSSION

### Clinical outcome of prenatally diagnosed mosaic trisomy 16 pregnancies

Although this sample is likely to be biased towards poorer outcome, the majority of prenatally diagnosed mosaic trisomy 16 pregnancies resulted in live births with survival beyond the neonatal period. The distribution of gestational ages for all the live births suggests that trisomy 16 pregnancies may be at higher risk for preterm delivery (fig 1). Further research is needed to determine whether there is truly a higher risk of preterm delivery in an unbiased population, and, if so, whether it is primarily the result of preterm labour or induction or caesarean section secondary to other complications. In addition, virtually all birth weights for the live births were below the gestational age corrected mean birth weights in the general population<sup>14</sup> (fig 2). This indicates that some level of below average growth is a nearly universal phenomenon in trisomy 16 mosaicism, and supports the hypothesis of undetected trisomy mosaicism as an aetiological factor in both severe and mild idiopathic intrauterine growth restriction. Anatomical classes of malformations were present at approximately the same frequency (~20–45%), with cardiac malformations most commonly seen (62%). However, some individual malformations (VSD, ASD, and hypospadias) were particularly frequent among live births (>6% of informative cases) and were significantly more frequent compared to the



**Figure 3** Summary of the pathogenesis of trisomy 16 mosaicism. The cytotrophoblast was assessed via direct CVS. The chorionic villous stroma was assessed via cultured CVS. The amniotic fluid was assessed via amniocentesis. The amniotic mesenchyme was assessed via postnatal or necropsy sampling of the amnion followed by culture. Growth restriction was coded as BW (table 1). Malformation was coded as MALF (table 1). The inclusion of UPD(16)mat is from previously reported results.<sup>10</sup>

general neonatal population (Electronic Appendix 2). These malformations are likely to have a true association with trisomy mosaicism of chromosome 16.

### Trisomy 16 in the placenta

It has previously been shown that the clinical outcome of trisomy predominantly or completely confined to the placenta was strongly associated with the level of trisomy in the term trophoblast but not with the level of trisomy on CVS.<sup>7</sup> However, the data were confounded by the inclusion of a mixture of trisomies involving different chromosomes and different origins. In this study sample, non-mosaic trisomy on direct CVS (cytotrophoblast) was found to be associated with low BW (table 2) and increased risk of malformation (table 3). Cytotrophoblast function is important for implantation and for differentiation to the hormone secreting syncytiotrophoblast and invasive extravillous cytotrophoblast. Placental insufficiency is a cause of intrauterine growth restriction and can be caused by poor remodelling of maternal spiral arteries, which is thought to depend on normal extravillous cytotrophoblast function.<sup>19</sup> Increased apoptosis of syncytiotrophoblast has been observed in pregnancies complicated by growth restriction,<sup>20</sup> while decreased villous trophoblast proliferation has been seen in spontaneous abortions with chromosome abnormalities (mostly single autosomal trisomies).<sup>21</sup> Notably, it has been shown that trisomy 21 placentas have a defect in cytotrophoblast differentiation to syncytiotrophoblast.<sup>22</sup>

Non-mosaic trisomy on cultured CVS (chorionic villous stroma) was also found to be associated with BW (table 2). This may be because of correlation with high levels of trisomy in the cytotrophoblast or malfunction of the chorionic villous stroma. For instance, the chorionic villous stroma is thought to be an inducer of overlying cytotrophoblast proliferation.<sup>23</sup> Increased apoptosis and decreased proliferation of chorionic villous stromal cells have also been observed in spontaneous abortions with chromosome abnormalities (mostly single autosomal trisomies).<sup>21</sup>

### Trisomy 16 in the amniotic mesenchyme

The presence of trisomy in the amniotic mesenchyme was associated with lower BW (table 2). This may be because of a correlation between trisomy in the amniotic mesenchyme and

trisomy in the amniotic fluid (and therefore, fetal tissues) (Electronic Appendix 3). The association between trisomy in the amniotic mesenchyme and outcome may also be the result of functional effects of the trisomy. For example, the amniochorionic membrane is part of the “transmembranous” pathway of fluid exchange between the amniotic fluid and maternal blood that is involved in amniotic fluid regulation.<sup>24</sup>

### Trisomy 16 in the fetus

The presence of trisomy on amniocentesis (that is, in amniotic fluid) was associated with lower BW (table 2) and increased risk of malformation (table 3) confirming the importance of trisomy in fetal tissues for outcome. Interestingly, it is the simple presence of trisomy at amniocentesis, but not the level above 0%, that appears to be important. This finding is similar to the observation that the level of trisomy 18 in lymphocytes does not appear to correlate with outcome in mosaic trisomy 18 subjects.<sup>25, 26</sup> It also suggests that the level of trisomy on amniocentesis is influenced by stochastic processes during cell culture and during development, and should be considered a random “snapshot” of the degree and distribution of trisomy in the fetus.

### Implications for the prenatal diagnosis of trisomy 16 mosaicism

Tables 2 and 3 describe the predictive value of amniocentesis and direct and cultured CVS for outcome. Also, the risk of finding trisomy on amniocentesis is independent of the levels of trisomy on direct and cultured CVS (Electronic Appendix 3). In other words, cases with non-mosaic trisomy on either direct or cultured CVS were not more likely to have trisomy detected at amniocentesis compared to cases with <100% trisomy on either direct or cultured CVS.

It is also important to re-emphasise that most cases of trisomy 16 are initially diagnosed with non-mosaic trisomy on CVS. However, as noted previously, virtually all embryos that survive to the time of prenatal diagnosis are mosaic. Thus, even with non-mosaic (100%) trisomy detected on CVS, the clinical outcomes of the pregnancy will correspond to outcomes seen in the population of mosaic trisomy 16 pregnancies. Moreover, although several variables have been shown to be predictive of outcome, there is variation in

outcome even for a given value of a variable (tables 2 and 3). It should also be noted that ultrasound is one prenatal variable that has not been assessed in this study, but is likely to be important for identifying those cases with the worse prognoses.<sup>11</sup>

### Implications for the pathogenesis of trisomy 16 mosaicism

Fig 3 summarises the results pertinent to the pathogenesis of trisomy 16 mosaicism. Trisomy in the cytotrophoblast (detected by direct CVS), chorionic villous stroma (detected by cultured CVS), amniotic fluid (detected by amniocentesis), and amniotic mesenchyme (detected by postnatal or necropsy sampling followed by culture) appear to have an effect on risk of intrauterine growth restriction, while only trisomy in the cytotrophoblast and amniotic fluid appear to have an effect on risk of malformation. Since the levels of trisomy in the placental tissues may be independent of the levels of trisomy in the amniotic fluid and amniotic mesenchyme (Electronic Appendix 3), it is possible that the effect of placental trisomy on outcome seen in this study sample may be independent (at least in part) of the effect of trisomy present in the fetus. Thus, it is theoretically possible that prenatal therapeutic interventions to improve placental function may improve outcome in trisomic pregnancies.

### Sex of the fetus in trisomy 16 mosaicism

The sex ratio in the study sample (0.45) was biased towards females, as initially noted by Benn,<sup>3</sup> and was significantly different from the expected sex ratio calculated from prenatal controls.<sup>17</sup> This is similar to a report of a significant excess of females in prenatally diagnosed trisomy 21 mosaicism (sex ratio = 0.72) compared to an excess of males in prenatally diagnosed non-mosaic trisomy 21.<sup>27</sup> The same phenomenon was found in another report,<sup>17</sup> in addition to a significant excess of females in prenatally diagnosed trisomy 18 mosaicism (sex ratio = 0.52) and a trend towards excess females in prenatally diagnosed trisomy 13 mosaicism (sex ratio = 0.76) compared to prenatal controls.

We found no evidence for selection against male fetuses after the time of CVS, indicating that the bias in sex ratio is set in the first trimester (before ~9-12 weeks). As previously suggested,<sup>3</sup> two possible explanations are (1) increased probability of rescue in female non-mosaic trisomy 16 embryos and/or (2) increased selection against male mosaic trisomy 16 embryos before the time of CVS.

### Future directions

Although this study has focused on intrauterine growth restriction and malformation, the most important outcome, long term postnatal prognosis, has been not been assessed because adequate follow up data are not yet available. The most common malformations in neonates (hypospadias, and ASD/VSD if clinically insignificant) may not have an impact on long term outcome, and of the few cases with long term follow up (S Langlois, unpublished data),<sup>11,12</sup> postnatal development seems to progress quite well in general. We encourage clinical geneticists world wide formally to follow up cases of prenatal diagnosis of trisomy mosaicism at their centres and to publish this information for the improvement of genetic counselling.

### ACKNOWLEDGEMENTS

We thank Dr Deborah McFadden for information regarding fetal malformation and FISH examination of the placenta. This research was supported by the Canadian Institutes of Health Research (CIHR) (grant 15667). PJY is supported by studentships from the CIHR and the Killam Trust.

### Authors' affiliations

**P J Yong**, MD/PhD and Experimental Medicine Programs, University of British Columbia and the British Columbia Research Institute for Children's and Women's Health, Vancouver, British Columbia, Canada  
**I J Barrett**, **W P Robinson**, Department of Medical Genetics, University of British Columbia and the British Columbia Research Institute for Children's and Women's Health, Vancouver, British Columbia, Canada.  
**D K Kalousek**, Department of Pathology and Laboratory Medicine, University of British Columbia and the British Columbia Research Institute for Children's and Women's Health, Vancouver, British Columbia, Canada



Supplementary appendices can be found on our website, [www.jmedgenet.com](http://www.jmedgenet.com)

### REFERENCES

- Wolstenholme J.** An audit of trisomy 16 in man. *Prenat Diagn* 1995;**15**:109-21.
- Hassold TJ, Jacobs PA.** Trisomy in man. *Ann Rev Genet* 1984;**18**:69-97.
- Benn P.** Trisomy 16 and trisomy 16 mosaicism: a review. *Am J Med Genet* 1998;**79**:121-33.
- Association of Clinical Cytogenetics Working Party on Chorionic Villi in Prenatal Diagnosis.** Cytogenetic analysis of chorionic villi for prenatal diagnosis: an ACC collaborative study of UK data. *Prenat Diagn* 1994;**14**:363-79.
- Cusick W, Bork M, Fabri B, Benn P, Rodis JF, Butting L Jr.** Trisomy 16 fetus surviving into the second trimester. *Prenat Diagn* 1995;**15**:1078-81.
- Yancey MK, Hardin EL, Pacheco C, Kuslich CD, Donlon TA.** Non-mosaic trisomy 16 in a third-trimester fetus. *Obstet Gynecol* 1996;**87**:856-60.
- Robinson WP, Barrett IJ, Bernard L, Telenius A, Bernasconi F, Wilson RD, Best RG, Howard-Peebles PN, Langlois S, Kalousek DK.** Meiotic origin of trisomy in confined placental mosaicism is correlated with presence of fetal uniparental disomy, high levels of trisomy in trophoblast, and increased risk of fetal intrauterine growth restriction. *Am J Hum Genet* 1997;**60**:917-27.
- Engel E.** A new genetic concept: uniparental disomy and its potential effect, isodisomy. *Am J Med Genet* 1980;**6**:137-43.
- Spence JE, Perciaccante RG, Greig GM, Willard HF, Ledbetter DH, Hejtmancik JF, Pollack MS, O'Brien WE, Beaudet AL.** Uniparental disomy as a mechanism for human genetic disease. *Am J Hum Genet* 1988;**42**:217-26.
- Yong PJ, Marion SA, Barrett IJ, Kalousek DK, Robinson WP.** Evidence for imprinting on chromosome 16: effect of uniparental disomy on the outcome of mosaic trisomy 16 pregnancies. *Am J Med Genet* 2002;**112**:123-32.
- Peñaherrera MS, Barrett IJ, Brown CJ, Langlois S, Yong SL, Lewis S, Bruyere H, Howard-Peebles PN, Kalousek DK, Robinson WP.** An association between skewed X-chromosome inactivation and abnormal outcome in mosaic trisomy 16 confined predominantly to the placenta. *Clin Genet* 2000;**58**:436-46.
- Schneider AS, Bischoff FZ, McCaskill C, Coady ML, Stopfer JE, Shaffer LG.** Comprehensive 4-year follow-up on a case of maternal heterodisomy for chromosome 16. *Am J Med Genet* 1996;**66**:204-8.
- Robinson WP, McFadden DE, Barrett IJ, Kuchinka B, Peñaherrera MS, Bruyere H, Best RG, Pedreira DAL, Langlois S, Kalousek DK.** Origin of amnion and implications for evaluation of the fetal genotype in cases of mosaicism. *Prenat Diagn* 2002;**22**:1076-85.
- Usher R, McLean F.** Intrauterine growth of live-born Caucasian infants at sea-level: standards obtained from measurements in 7 dimensions of infants born between 25 and 44 weeks of gestation. *J Pediatr* 1969;**74**:901-10.
- Sadler TW.** *Langman's medical embryology*. Philadelphia: Lippincott Williams & Wilkins, 2000.
- Wolstenholme J.** Confined placental mosaicism for trisomies 2, 3, 6, 8, 9, 16, and 22: their incidence, likely origins, and mechanisms for cell lineage compartmentalization. *Prenat Diagn* 1996;**16**:511-24.
- Hueith CA, Martin RLM, Stoppelman SM, D'Souza S, Bishop JK, Torfs CP, Lorey F, May KM, Hanna JS, Baird PA, Kelly JC.** Sex ratios in fetuses and liveborn infants with autosomal aneuploidy. *Am J Med Genet* 1996;**63**:492-500.
- Hassold T, Quillen SD, Yamane JA.** Sex ratio in spontaneous abortions. *Ann Hum Genet* 1983;**47**, Pt 1:39-47.
- Cartwright JE, Kenny LC, Dash PR, Crocker IP, Aplin JD, Baker PN, Whitley GS.** Trophoblast invasion of spiral arteries: a novel in vitro model. *Placenta* 2002;**23**:232-5.
- Ishihara N, Matsuo H, Murakoshi H, Laog-Fernandez LB, Samoto T, Maruo T.** Increased apoptosis in the syncytiotrophoblast in human term placentas complicated by either preeclampsia or intrauterine growth retardation. *Am J Obstet Gynecol* 2002;**186**:158-66.

- 21 **Qumsiyeh MB**, Kim KR, Ahmed MN, Bradford W. Cytogenetics and mechanisms of spontaneous abortions: increased apoptosis and decreased cell proliferation in chromosomally abnormal villi. *Cytogenet Cell Genet* 2000;**88**:230-5.
- 22 **Frendo JL**, Vidaud M, Guibourdenche J, Luton D, Muller F, Bellet D, Giovagrandi Y, Tarrade A, Porquet D, Blot P. Defect of villous cytotrophoblast differentiation into syncytiotrophoblast in Down's syndrome. *J Clin Endocrinol Metab* 2001;**85**:3700-7.
- 23 **Moore KL**, Persaud TVN. *Before we are born: essentials of embryology and birth defects*. Philadelphia: Saunders, 1998.
- 24 **Brace RA**, Resnik R. Dynamics and disorders of amniotic fluid. In: Creasy RK, Resnik R, eds. *Maternal-fetal medicine*. Philadelphia: Saunders, 1999:632-43.
- 25 **Graham DA**, Jewitt MM, Fitzgerald PH. Trisomy 18 mosaicism with complete peripheral lymphocyte trisomy and normal intelligence. *Clin Genet* 1992;**41**:36-8.
- 26 **Lim AS**, Su LC. Mosaic trisomy 18 male with normal intelligence who fathered a normal baby girl. *Am J Med Genet* 1998;**76**:365-6.
- 27 **Hook EB**, Cross PK, Mutton DE. Female predominance (low sex ratio) in 47,+21 mosaics. *Am J Med Genet* 1999;**84**:316-19.

Find out what's in the latest issue  
the moment it's published

Email Alerts

Sign up to receive the table of contents by email every month. You can select from three alerts: Table of Contents (full), TOC Awareness (notice only); *Journal of Medical Genetics* related announcements.

[www.jmedgenet.com](http://www.jmedgenet.com)