

Subject index

- 11p**, Is the locus for Costello syndrome on 11p?, **469**
- 11p15**, Paternal uniparental disomy in monozygotic twins discordant for hemihypertrophy, **223**
- 1555A>G mutation**, Heteroplasmy for the 1555A>G mutation in the mitochondrial 12S rRNA gene in six Spanish families with non-syndromic hearing loss, **632**
- 1q21-q23**, DFNA49, a novel locus for autosomal dominant non-syndromic hearing loss, maps proximal to DFNA7/DFNM1 region on chromosome 1q21-q23, **832**
- 22 translocation**, Neurofibromatosis 2 in a patient with a *de novo* balanced reciprocal translocation 46,X,t(X;22)(p11.2;q11.2), **682**
- 22q13**, Molecular characterisation of the 22q13 deletion syndrome supports the role of haploinsufficiency of *SHANK3/PROSAP2* in the major neurological symptoms, **575**
- 2p12**, Mapping of a new autosomal dominant non-syndromic hearing loss locus (DFNA43) to chromosome 2p12, **278**
- 5,10-methylenetetrahydrofolate reductase**, Maternal 677CT/1298AC genotype of the *MTHFR* gene as a risk factor for cleft lip, **e64**
- 677C>T**, Geographical and ethnic variation of the 677C>T allele of 5,10 methylenetetrahydrofolate reductase (*MTHFR*): findings from over 7000 newborns from 16 areas world wide, **619**
- 7q36**, No association between the *EN2* gene and autistic disorder, **e4**
- 9q34**, Identification of recurrent regions of chromosome loss and gain in vestibular schwannomas using comparative genomic hybridisation, **802**
- 9q34 deletion**, Cryptic terminal deletion of chromosome 9q34: a novel cause of syndromic obesity in childhood?, **300**
- α -galactosidase deficiency**, Fabry disease: molecular studies in Italian patients and X inactivation analysis in manifesting carriers, **e103**
- ABCA1**, A common variant in the *ABCA1* gene is associated with a lower risk for premature coronary heart disease in familial hypercholesterolaemia, **163**
- ABCB4**, *ABCB4* gene sequence variation in women with intrahepatic cholestasis of pregnancy, **e70**
- ABL gene**, Deletion in the *ABL* gene resulting from a meiotic recombination of a maternal (3;22;9)(q22;q12;q34.1) translocation, **e6**
- acanthosis nigricans**, Studies on the pathogenesis of Costello syndrome, **e37**
- ACO2**, Mutation and LOH analysis of *ACO2* in colorectal cancer: no evidence of biallelic genetic inactivation, **e73**
- acrocapitofemoral dysplasia**, Acrocapitofemoral dysplasia: an autosomal recessive skeletal dysplasia with cone shaped epiphyses in the hands and hips, **201**
- acro-renal-ocular syndrome**, Mutations at the *SALL4* locus on chromosome 20 result in a range of clinically overlapping phenotypes, including Okihiro syndrome, Holt-Oram syndrome, acro-renal-ocular syndrome, and patients previously reported to represent thalidomide embryopathy, **473**
- ACTG1**, A mutation in the gamma actin 1 (*ACTG1*) gene causes autosomal dominant hearing loss (DFNA20/26), **879**
- actin**, A mutation in the gamma actin 1 (*ACTG1*) gene causes autosomal dominant hearing loss (DFNA20/26), **879**
- ADHD**, A novel pericentric inversion of chromosome 3 cosegregates with a developmental-behavioural phenotype, **e15**
Disruption of a novel member of a sodium/hydrogen exchanger family and *DOCK3* is associated with an attention deficit hyperactivity disorder-like phenotype, **733**
- adolescents**, A genetic screening programme for Tay-Sachs disease and cystic fibrosis for Australian Jewish high school students, **e45**
- adrenal hypoplasia**, A complex deletion-inversion-deletion event results in a chimeric *IL1RAPL1*-dystrophin transcript and a contiguous gene deletion syndrome, **127**
- AEC syndrome**, Rapp-Hodgkin and AEC syndromes due to a new frameshift mutation in the TP63 gene, **e133**
- AGTR2 gene**, Rare polymorphic variants of the *AGTR2* gene in boys with non-specific mental retardation, **357**
- AIB1**, Somatic instability of the DNA sequences encoding the polymorphic polyglutamine tract of the *AIB1* gene, **885**
- AIB1 and breast cancer**, Somatic instability of the DNA sequences encoding the polymorphic polyglutamine tract of the *AIB1* gene, **885**
- Aicardi-Goutières syndrome**, Cree encephalitis is allelic with Aicardi-Goutières syndrome: implications for the pathogenesis of disorders of interferon alpha metabolism, **183**
- Alagille syndrome**, Consequences of *JAG1* mutations, **891**
- aldosterone synthase**, Gender specific association of aldosterone synthase gene polymorphism with renal survival in patients with IgA nephropathy, **372**
- ALK1**, Hereditary haemorrhagic telangiectasia: a questionnaire based study to delineate the different phenotypes caused by endoglin and *ALK1* mutations, **585**
- ALK-1**, Molecular and functional analysis identifies *ALK-1* as the predominant cause of pulmonary hypertension related to hereditary haemorrhagic telangiectasia, **865**
- alkaline phosphatase**, Molecular study of three cases of odontohypophosphatasia resulting from heterozygosity for mutations in the tissue non-specific alkaline phosphatase gene, **605**
- allele specific replication timing**, The imprinted region on human chromosome 7q32 extends to the carboxypeptidase A gene cluster: an imprinted candidate for Silver-Russell syndrome, **249**
- allelic imbalance**, *CHEK2* 1100delC and colorectal cancer, **e110**
- alternative splicing**, Transcript analysis of the cystic fibrosis splicing mutation 1525-1G>A shows use of multiple alternative splicing sites and suggests a putative role of exonic splicing enhancers, **e88**
- Alzheimer's disease**, Association of 3'-UTR polymorphisms of the oxidised LDL receptor 1 (*OLR1*) gene with Alzheimer's disease, **424**
- amelogenesis imperfecta**, Novel *ENAM* mutation responsible for autosomal recessive amelogenesis imperfecta and localised enamel defects, **900**
- amyloid neuropathy**, Genetic study of transthyretin amyloid neuropathies: carrier risks among French and Portuguese families, **e120**
- analphoid marker chromosome**, Analphoid *de novo* marker chromosome inv dup(3)(q28qter) with neocentromere in a dysmorphic and developmentally retarded girl, **e27**
- aneuploidy**, Rapid, high throughput prenatal detection of aneuploidy using a novel quantitative method (MLPA), **907**
- Angelman syndrome**, Angelman syndrome: a review of the clinical and genetic aspects, **87**
- angio-oedema**, The effect of sequence variations within the coding region of the C1 inhibitor gene on disease expression and protein function in families with hereditary angio-oedema, **e114**
- angiotensin-converting enzyme**, Renoprotective efficacy of renin-angiotensin inhibitors in IgA nephropathy is influenced by *ACE* A2350G polymorphism, **e130**
- angiotensin-converting enzyme inhibitor**, Renoprotective efficacy of renin-angiotensin inhibitors in IgA nephropathy is influenced by *ACE* A2350G polymorphism, **e130**
- animal model of human disease**, Canine tricuspid valve malformation, a model of human Ebstein anomaly, maps to dog chromosome 9, **320**
- aplastic anaemia**, Short-limbed dwarfism with bowing, combined immune deficiency, and late onset aplastic anaemia caused by novel mutations in the *RMPR* gene, **761**
- APOE**, *APOE* and TGF- β 1 genes are associated with obesity phenotypes, **918**
- apolipoprotein E**, Does apolipoprotein E polymorphism influence susceptibility to malaria?, **348**
The role of *APOE* in the phenotypic expression of Leber hereditary optic neuropathy, **e41**
- apolipoprotein E gene**, Small babies receive the cardiovascular protective apolipoprotein ϵ 2 allele less frequently than expected, **626**
- apolipoprotein AV gene**, Ser19 \rightarrow Trp polymorphism within the apolipoprotein AV gene in hypertriglyceridaemic people, **e105**

- apoptosis**, Promoter polymorphisms of the *CD14* gene in Italian patients with coeliac disease, **e108**
- arCRD**, Evidence of *RPGRI1* gene mutations associated with recessive cone-rod dystrophy, **616**
- Arg276X**, High incidence of the R276X *SALL1* mutation in sporadic but not familial Townes-Brocks syndrome and report of the first familial case, **e127**
- arteriovenous**, Visceral manifestations in hereditary haemorrhagic telangiectasia type 2, **494**
- Ashkenazi Jews**, A twofold increase in *BRCA* mutation related prostate cancer among Ashkenazi Israelis is not associated with distinctive histopathology, **787**
Genetic homogeneity and phenotypic variability among Ashkenazi Jews with Usher syndrome type III, **767**
The *RNASEL* 471delAAAG allele and prostate cancer in Ashkenazi Jewish men, **e22**
- asphyxiating thoracic dystrophy**, A locus for asphyxiating thoracic dystrophy, *ATD*, maps to chromosome 15q13, **431**
- assessment**, Evaluation of breast cancer risk assessment packages in the family history evaluation and screening programme, **807**
- association**, The metabotropic glutamate receptor 8 gene at 7q31: partial duplication and possible association with autism, **e42**
- association study**, A Y chromosomal influence on prostate cancer risk: the multi-ethnic cohort study, **815**
- atlastin**, Science in motion: common molecular pathological themes emerge in the hereditary spastic paraplegias, **81**
- atopy**, A common IL-1 complex haplotype is associated with an increased risk of atopy, **e66**
- atrial fibrillation**, Expanding the phenotype of *LMNA* mutations in dilated cardiomyopathy and functional consequences of these mutations, **560**
- atrial/ventricular septal defects**, Novel *TBX5* mutations and molecular mechanism for Holt-Oram syndrome, **e29**
- attitudes**, Genetic information but not termination: pregnant women's attitudes and willingness to pay for carrier screening for deafness genes, **e80**
- atypical Kearns Sayre syndrome**, Compensatory amplification of mtDNA in a patient with anovel deletion/duplication and high mutant load, **e125**
- auditory neuropathy**, Non-syndromic recessive auditory neuropathy is the result of mutations in the otoferlin (*OTOF*) gene, **45**
- autism**, Association of *INPP1*, *PIK3CG*, and *TSC2* gene variants with autistic disorder: implications for phosphatidylinositol signalling in autism, **e119**
No association between the *EN2* gene and autistic disorder, **e4**
The neurobeachin gene is disrupted by a translocation in a patient with idiopathic autism, **352**
- autistic behaviour**, Myhre syndrome: new reports, review, and differential diagnosis, **546**
- autoimmunity**, Promoter polymorphisms of the *CD14* gene in Italian patients with coeliac disease, **e108**
- autosomal dominant**, Evidence of autosomal dominant Leber congenital amaurosis (LCA) underlain by a *CRX* heterozygous null allele, **e90**
- autosomal recessive**, Mapping of a new locus for autosomal recessive non-syndromic mental retardation in the chromosomal region 19p13.12-p13.2: further genetic heterogeneity, **729**
Microcephaly-cardiomyopathy syndrome: expansion of the phenotype, **e78**
- autosomal recessive inheritance**, Clinical, radiological, and chondro-osseous findings in opsismodysplasia: survey of a series of 12 unreported cases, **195**
- autozygosity mapping**, A locus for asphyxiating thoracic dystrophy, *ATD*, maps to chromosome 15q13, **431**
A novel locus for autosomal recessive primary microcephaly (MCPH6) maps to 13q12.2, **540**
- azoospermia factor b**, The human Y chromosome's azoospermia factor b (AZFb) region: sequence, structure, and deletion analysis in infertile men, **18**
- Bannayan-Riley-Ruvalcaba syndrome**, *PTEN* hamartoma tumour syndrome: variability of an entity, **e111**
- Bardet-Biedl syndrome**, Further support for digenic inheritance in Bardet-Biedl syndrome, **e104**
- Barrett's oesophagus**, Demographic and phenotypic features of 70 families segregating Barrett's oesophagus and oesophageal adenocarcinoma, **651**
- basal ganglia**, Karak syndrome: a novel degenerative disorder of the basal ganglia and cerebellum, **543**
Neuroferritinopathy in a French family with late onset dominant dystonia, **e69**
- BDA1**, Identification of a mutation in the Indian Hedgehog (*IHH*) gene causing brachydactyly type A1 and evidence for a third locus, **42**
- BDA-1**, Mild brachydactyly type A1 maps to chromosome 2q35-q36 and is caused by a novel *IHH* mutation in a three generation family, **132**
- Beckwith-Wiedemann syndrome**, Beckwith-Wiedemann syndrome and assisted reproduction technology (ART), **62**
Paternal uniparental disomy in monozygotic twins discordant for hemihypertrophy, **223**
Silencing of *CDKN1C* (*p57^{KIP2}*) is associated with hypomethylation at KvDMR1 in Beckwith-Wiedemann syndrome, **797**
- benign telangiectasia**, Autosomal dominant hereditary benign telangiectasia maps to the *CMC1* locus for capillary malformation on chromosome 5q14, **849**
- BHD**, Alterations of the Birt-Hogg-Dubé gene (*BHD*) in sporadic colorectal tumours, **511**
Analysis of the Birt-Hogg-Dubé (*BHD*) tumour suppressor gene in sporadic renal cell carcinoma and colorectal cancer, **820**
- biparental hydatidiform mole**, Genetic refinement and physical mapping of a biparental complete hydatidiform mole locus on chromosome 19q13.4., **e95**
- bipolar affective disorder**, Prader-Willi syndrome: the psychopathological phenotype in uniparental disomy, **e112**
- birth defects**, Complete sequencing shows a role for *MSX1* in non-syndromic cleft lip and palate, **399**
- Birt-Hogg-Dubé syndrome**, Mutations of the Birt-Hogg-Dubé (*BHD*) gene in sporadic colorectal carcinomas and colorectal carcinoma cell lines with microsatellite instability, **364**
- bisulphite**, Modulation of methylation in the *FMR1* promoter region after long term treatment with L-carnitine and acetyl-L-carnitine, **1**
- bladder carcinoma**, Studies on the pathogenesis of Costello syndrome, **e37**
- BMPR-II**, Molecular and functional analysis identifies ALK-1 as the predominant cause of pulmonary hypertension related to hereditary haemorrhagic telangiectasia, **865**
- bone marrow transplantation**, Short-limbed dwarfism with bowing, combined immune deficiency, and late onset aplastic anaemia caused by novel mutations in the *RMPR* gene, **761**
- Börjeson-Forsmann-Lehmann syndrome**, Novel *PHF6* mutation p.D333del causes Börjeson-Forsman-Lehmann syndrome, **e50**
- BOS2**, A gene locus for branchio-otic syndrome maps to chromosome 14q21.3-q24.3, **515**
- BOS3**, A gene locus for branchio-otic syndrome maps to chromosome 14q21.3-q24.3, **515**
- brachydactyly**, Myhre syndrome: new reports, review, and differential diagnosis, **546**
- brachydactyly type A1**, Identification of a mutation in the Indian Hedgehog (*IHH*) gene causing brachydactyly type A1 and evidence for a third locus, **42**
Mild brachydactyly type A1 maps to chromosome 2q35-q36 and is caused by a novel *IHH* mutation in a three generation family, **132**
- brain**, Association of 3'-UTR polymorphisms of the oxidised LDL receptor 1 (*OLR1*) gene with Alzheimer's disease, **424**
- brain development**, Mutations in the oligophrenin-1 gene (*OPHN1*) cause X linked congenital cerebellar hypoplasia, **441**
- branchio-otic syndrome**, A gene locus for branchio-otic syndrome maps to chromosome 14q21.3-q24.3, **515**
- branchio-oto-renal syndrome (BOR)**, A gene locus for branchio-otic syndrome maps to chromosome 14q21.3-q24.3, **515**
- Brazil**, High frequency of novel germline mutations in the *VHL* gene in the heterogeneous population of Brazil, **e31**
- BRCA**, A twofold increase in *BRCA* mutation related prostate cancer among Ashkenazi Israelis is not associated with distinctive histopathology, **787**
- BRCA1**, Application of haplotype pair analysis for the identification of hemizygous loci, **346**
BRCA1 and *BRCA2* mutations have no major role in predisposition to prostate cancer in Finland, **e98**
BRCA1 c.2845insA is a recurring mutation with a founder effect in Singapore Malay women with early onset breast/ovarian cancer, **e117**

- Germline *TP53* mutations in breast cancer families with multiple primary cancers: is *TP53* a modifier of *BRCA1*?, **e34**
- Individual and family characteristics associated with protein truncating *BRCA1* and *BRCA2* mutations in an Ontario population based series from the Cooperative Family Registry for Breast Cancer Studies, **e91**
- Neuronal migration defect in a *BRCA1* gene carrier: possible focal nullisomy?, **e24**
- Non-random transmission of mutant alleles to female offspring of *BRCA1* carriers in Poland, **719**
- Pre-test prediction models of *BRCA1* or *BRCA2* mutation in breast/ovarian families attending familial cancer clinics, **503**
- Sensitivity of *BRCA1/2* mutation testing in 466 breast/ovarian cancer families, **e107**
- BRCA1/BRCA2**, Rapid, inexpensive scanning for all possible *BRCA1* and *BRCA2* gene sequence variants in a single assay: implications for genetic testing, **e33**
- BRCA2**, Individual and family characteristics associated with protein truncating *BRCA1* and *BRCA2* mutations in an Ontario population based series from the Cooperative Family Registry for Breast Cancer Studies, **e91**
- BRCA1* and *BRCA2* mutations have no major role in predisposition to prostate cancer in Finland, **e98**
- Pre-test prediction models of *BRCA1* or *BRCA2* mutation in breast/ovarian families attending familial cancer clinics, **503**
- Sensitivity of *BRCA1/2* mutation testing in 466 breast/ovarian cancer families, **e107**
- The *BRCA2* variant 8204G>A is a splicing mutation and results in an in frame deletion of the gene, **e23**
- breast and ovarian cancer susceptibility**, Mutation screening of *Mre11* complex genes: indication of *RAD50* involvement in breast and ovarian cancer susceptibility, **e131**
- breast cancer**, Absence of constitutional *H2AX* gene mutations in 101 hereditary breast cancer families, **e51**
- Evaluation of breast cancer risk assessment packages in the family history evaluation and screening programme, **807**
- Germline *TP53* mutations in breast cancer families with multiple primary cancers: is *TP53* a modifier of *BRCA1*?, **e34**
- Identification of a splice acceptor site mutation in *p16^{INK4A}/p14^{ARF}* within a breast cancer, melanoma, neurofibroma prone kindred, **e102**
- Individual and family characteristics associated with protein truncating *BRCA1* and *BRCA2* mutations in an Ontario population based series from the Cooperative Family Registry for Breast Cancer Studies, **e91**
- Neuronal migration defect in a *BRCA1* gene carrier: possible focal nullisomy?, **e24**
- Prevalence of family histories of breast cancer in the general population and the incidence of related seeking of health care, **e83**
- Sensitivity of *BRCA1/2* mutation testing in 466 breast/ovarian cancer families, **e107**
- breast neoplasms**, Little value from including cousins in individual risk assessment of hereditary breast cancer: a simulation study, **e25**
- breast/ovarian cancer**, *BRCA1* c.2845insA is a recurring mutation with a founder effect in Singapore Malay women with early onset breast/ovarian cancer, **e117**
- bronchiectasis**, High frequency of T9 and *CFTR* mutations in children with idiopathic bronchiectasis, **530**
- C1 inhibitor**, The effect of sequence variations within the coding region of the C1 inhibitor gene on disease expression and protein function in families with hereditary angio-oedema, **e114**
- C282Y mutation**, Haemochromatosis (*HFE*) gene C282Y mutation and the risk of coronary artery disease and myocardial infarction: a study in 1279 patients undergoing coronary angiography, **e58**
- C3**, Haemolytic uraemic syndrome and mutations of the factor H gene: a registry-based study of German speaking countries, **676**
- C7orf10**, Genomic characterisation of *C7orf10* in Silver-Russell syndrome patients, **e44**
- CAG repeats in AIB1**, Somatic instability of the DNA sequences encoding the polymorphic polyglutamine tract of the *AIB1* gene, **885**
- cancer**, Analysis of the Birt-Hogg-Dubé (*BHD*) tumour suppressor gene in sporadic renal cell carcinoma and colorectal cancer, **820**
- cancer genetics**, Chromosome 2 (2p16) abnormalities in Carney complex tumours, **268**
- RNASEL* mutations in hereditary prostate cancer, **e21**
- cancer predisposition**, Cancer risk in 348 French *MSH2* or *MLH1* gene carriers, **208**
- cancer screening**, Cancer surveillance is often inadequate in people at high risk for colorectal cancer, **e54**
- capillary malformation**, Autosomal dominant hereditary benign telangiectasia maps to the *CMC1* locus for capillary malformation on chromosome 5q14, **849**
- carbonic anhydrase II**, A phenocopy of *CAII* deficiency: a novel genetic explanation for inherited infantile osteopetrosis with distal renal tubular acidosis, **115**
- carbonic anhydrases**, Expression of cell surface transmembrane carbonic anhydrase genes *CA9* and *CA12* in the human eye: overexpression of *CA12* (*CAXII*) in glaucoma, **257**
- carboxypeptidase A (CPA)**, The imprinted region on human chromosome 7q32 extends to the carboxypeptidase A gene cluster: an imprinted candidate for Silver-Russell syndrome, **249**
- CARD15 SNPs**, Linkage of ulcerative colitis to the pericentromeric region of chromosome 16 in Italian inflammatory bowel disease families is independent of the presence of common *CARD15* mutations, **837**
- cardiomyopathy**, Expanding the phenotype of *LMNA* mutations in dilated cardiomyopathy and functional consequences of these mutations, **560**
- carrier detection**, An alternative to FISH: detecting deletion and duplication carriers within 24 hours, **e113**
- carrier females**, Clinical presentation in female patients with Fabry disease, **e38**
- carrier frequency**, A Tyr368His *RPE65* founder mutation is associated with variable expression and progression of early onset retinal dystrophy in 10 families of a genetically isolated population, **709**
- carrier risk**, Carrier risk status changes resulting from mutation testing in hereditary non-polyposis colorectal cancer and hereditary breast-ovarian cancer, **591**
- carrier screening**, Genetic information but not termination: pregnant women's attitudes and willingness to pay for carrier screening for deafness genes, **e80**
- cartilage oligomeric matrix protein**, A recurrent R718W mutation in *COMP* results in multiple epiphyseal dysplasia with mild myopathy: clinical and pathogenetic overlap with collagen IX mutations, **942**
- cartilage-hair hypoplasia**, Short-limbed dwarfism with bowing, combined immune deficiency, and late onset aplastic anaemia caused by novel mutations in the *RMPR* gene, **761**
- cat eye syndrome**, Partial trisomy of chromosome 22 resulting from an interstitial duplication of 22q11.2 in a child with typical cat eye syndrome, **e62**
- cataracts**, Genotype-phenotype correlations for cataracts in neurofibromatosis 2, **758**
- CD14**, Association of the *CD14* gene -159C polymorphism with progression of IgA nephropathy, **104**
- Promoter polymorphisms of the *CD14* gene in Italian patients with coeliac disease, **e108**
- CDH1**, A novel germline mutation in the *MET* extracellular domain in a Korean patient with the diffuse type of familial gastric cancer, **e97**
- CDSN**, Haplotype analysis of distantly related populations implicates corneodesmosin in psoriasis susceptibility, **447**
- central nervous system disease**, Hereditary xanthinuria type II associated with mental delay, autism, cortical renal cysts, nephrocalcinosis, osteopenia, and hair and teeth defects, **e121**
- centromere specific multicolour FISH (cenM FISH)**, Partial hexasomy 15pter→15q13 including *SNRPN* and *D15S10*: first molecular cytogenetically proven case report, **e28**
- cerebellar ataxia**, Hypogonadotrophic hypogonadism, short stature, cerebellar ataxia, rod-cone retinal dystrophy, and hypersegmented neutrophils: a novel disorder or a new variant of Boucher-Neuhauser syndrome?, **e2**
- cerebellar atrophy**, Progressive cerebellocerebral atrophy: a new syndrome with microcephaly, mental retardation, and spastic quadriplegia, **e96**
- cerebellum**, Karak syndrome: a novel degenerative disorder of the basal ganglia and cerebellum, **543**
- Mutations in the oligophrenin-1 gene (*OPHN1*) cause X linked congenital cerebellar hypoplasia, **441**
- cerebral cortex**, A novel locus for autosomal recessive primary microcephaly (*MCPH6*) maps to 13q12.2, **540**
- Transcript analysis of the cystic fibrosis splicing mutation 1525-1G>A shows use of multiple alternative splicing sites and suggests a putative role of exonic splicing enhancers, **e88**
- CFTR**, High frequency of T9 and *CFTR* mutations in children with idiopathic bronchiectasis, **530**

- CGH**, Identification of recurrent regions of chromosome loss and gain in vestibular schwannomas using comparative genomic hybridisation, **802**
- Charcot-Marie-Tooth disease**, Phenotypic and genetic exploration of severe demyelinating and secondary axonal neuropathies resulting from *GDAP1* nonsense and splicing mutations, **e87**
- CHEK2**, *CHEK2* 1100delC and colorectal cancer, **e110**
- child**, Melatonin supplementation for severe and intractable sleep disturbance in young people with genetically determined developmental disabilities: short review and commentary, **793**
- cholestasis**, A second heterozygous *MDR3* nonsense mutation associated with intrahepatic cholestasis of pregnancy, **e32**
ABCB4 gene sequence variation in women with intrahepatic cholestasis of pregnancy, **e70**
- chondrodysplasia punctata**, Greenberg dysplasia (HEM) and lethal X linked dominant Conradi-Hünermann chondrodysplasia punctata (CDPX2): presentation of two cases with overlapping phenotype, **e129**
- chromosome**, Telomeres: a diagnosis at the end of the chromosomes, **385**
- chromosome 14**, Comprehensive microsatellite marker analysis contradicts previous report of segmental maternal heterodisomy of chromosome 14, **e26**
Exclusion of maternal uniparental disomy of chromosome 14 in patients referred for Prader-Willi syndrome using a multiplex methylation polymerase chain reaction assay, **e46**
- chromosome 14q**, Molecular characterisation of a 3.5 Mb interstitial 14q deletion in a child with several phenotypic anomalies, **e47**
- chromosome 15**, Angelman syndrome: a review of the clinical and genetic aspects, **87**
Chromosome 15 maternal uniparental disomy and psychosis in Prader-Willi syndrome, **72**
Exclusion of maternal uniparental disomy of chromosome 14 in patients referred for Prader-Willi syndrome using a multiplex methylation polymerase chain reaction assay, **e46**
Identification of a locus for type I punctate palmoplantar keratoderma on chromosome 15q22-q24, **872**
- chromosome 15(q11-13)**, Death in adults with Prader-Willi syndrome may be correlated with maternal uniparental disomy, **e63**
- chromosome 17q25.3**, Homozygosity mapping of a Desbuquois dysplasia locus to chromosome 17q25.3, **282**
- chromosome 19q13.4**, Genetic refinement and physical mapping of a biparental complete hydatidiform mole locus on chromosome 19q13.4., **e95**
- chromosome 22**, Linkage and linkage disequilibrium searched for between non-syndromic cleft palate and four candidate loci, **464**
Neurofibromatosis 2 in a patient with a *de novo* balanced reciprocal translocation 46,X,t(X;22)(p11.2;q11.2), **682**
- chromosome 22q11.2**, Partial trisomy of chromosome 22 resulting from an interstitial duplication of 22q11.2 in a child with typical cat eye syndrome, **e62**
- chromosome breakpoint**, Array painting: a method for the rapid analysis of aberrant chromosomes using DNA microarrays, **664**
- chromosome deletion**, Unusual cognitive and behavioural profile in a Williams syndrome patient with atypical 7q11.23 deletion, **526**
- chromosome imbalance**, Distal trisomy 6p and 20q owing to the concurrent transposition of distal 6p and 20q to the 22q telomere: a genomic polymorphism?, **e94**
- chromosome instability**, Fanconi anaemia, **1**
- chromosome inversion**, A novel pericentric inversion of chromosome 3 cosegregates with a developmental-behavioural phenotype, **e15**
Disruption of a novel member of a sodium/hydrogen exchanger family and *DOCK3* is associated with an attention deficit hyperactivity disorder-like phenotype, **733**
- chromosome rearrangement**, Array painting: a method for the rapid analysis of aberrant chromosomes using DNA microarrays, **664**
- chronic progressive external ophthalmoplegia**, Characterisation of repeat and palindrome elements in patients harbouring single deletions of mitochondrial DNA, **e86**
- ciliary abnormalities**, *RPGR* mutation associated with retinitis pigmentosa, impaired hearing, and sinorespiratory infections, **609**
- circadian rhythm**, β_1 -adrenergic antagonists and melatonin reset the clock and restore sleep in a circadian disorder, Smith-Magenis syndrome, **74**
- cleft lip**, Maternal 677CT/1298AC genotype of the *MTHFR* gene as a risk factor for cleft lip, **e64**
- cleft lip and palate**, Complete sequencing shows a role for *MSX1* in non-syndromic cleft lip and palate, **399**
- cleft palate**, Linkage and linkage disequilibrium searched for between non-syndromic cleft palate and four candidate loci, **464**
Maternal 677CT/1298AC genotype of the *MTHFR* gene as a risk factor for cleft lip, **e64**
- clinical management**, Fanconi anaemia, **1**
- clinical variability**, Mild brachydactyly type A1 maps to chromosome 2q35-q36 and is caused by a novel *IHH* mutation in a three generation family, **132**
- club foot**, Recessive multiple epiphyseal dysplasia (rMED): phenotype delineation in eighteen homozygotes for *DTDST* mutation R279W, **65**
- CMC1**, Autosomal dominant hereditary benign telangiectasia maps to the *CMC1* locus for capillary malformation on chromosome 5q14, **849**
- CMT**, Diagnosis of gene dosage alterations at the *PMP22* gene using MAPH, **e123**
- CMT4A**, Phenotypic and genetic exploration of severe demyelinating and secondary axonal neuropathies resulting from *GDAP1* nonsense and splicing mutations, **e87**
- COCH**, Subcellular localisation, secretion, and post-translational processing of normal cochlin, and of mutants causing the sensorineural deafness and vestibular disorder, DFNA9, **479**
- cochlin**, Subcellular localisation, secretion, and post-translational processing of normal cochlin, and of mutants causing the sensorineural deafness and vestibular disorder, DFNA9, **479**
- COD1**, A new genetic locus for X linked progressive cone-rod dystrophy, **418**
- COD2**, A new genetic locus for X linked progressive cone-rod dystrophy, **418**
- coeliac disease**, Promoter polymorphisms of the *CD14* gene in Italian patients with coeliac disease, **e108**
- cognitive functioning**, Predictive testing for cognitive functioning in female carriers of the fragile X syndrome using hair root analysis, **377**
- cognitive profile**, Unusual cognitive and behavioural profile in a Williams syndrome patient with atypical 7q11.23 deletion, **526**
- Cohen syndrome**, Diagnostic criteria, clinical characteristics, and natural history of Cohen syndrome, **233**
- COL10A1**, Genetic changes in the RNA components of RNase MRP and RNase P in Schmid metaphyseal chondrodysplasia, **741**
- collagen IX**, A recurrent R718W mutation in *COMP* results in multiple epiphyseal dysplasia with mild myopathy: clinical and pathogenetic overlap with collagen IX mutations, **942**
- colonoscopy**, Experience of discharge from colonoscopy of mutation negative HNPCC family members, **e55**
- colorectal**, Analysis of the Birt-Hogg-Dubé (*BHD*) tumour suppressor gene in sporadic renal cell carcinoma and colorectal cancer, **820**
- colorectal cancer**, *CHEK2* 1100delC and colorectal cancer, **e110**
Mutation and LOH analysis of *ACO2* in colorectal cancer: no evidence of biallelic genetic inactivation, **e73**
Mutations of the Birt-Hogg-Dubé (*BHD*) gene in sporadic colorectal carcinomas and colorectal carcinoma cell lines with microsatellite instability, **364**
- colorectal tumours**, Alterations of the Birt-Hogg-Dubé gene (*BHD*) in sporadic colorectal tumours, **511**
- compensatory reversion**, In vivo reversion to normal of inherited mutations in humans, **721**
- complement factor H**, Haemolytic uraemic syndrome and mutations of the factor H gene: a registry-based study of German speaking countries, **676**
- complex chromosomal rearrangements**, Multicolour FISH fine mapping unravels an insertion as a complex chromosomal rearrangement involving six breakpoints and a 5.89 Mb large deletion, **e60**
- complex disease**, Inbreeding and risk of late onset complex disease, **925**
- complex trait**, Confirmation linkage study in support of the X chromosome harbouring a QTL underlying human height variation, **825**
- complex traits**, New approaches to investigating heterogeneity in complex traits, **553**
- complex translocation**, Deletion in the *ABL* gene resulting from a meiotic recombination of a maternal (3;22;9)(q22;q12;q34.1) translocation, **e6**
- conduct disorder**, A novel pericentric inversion of chromosome 3 cosegregates with a developmental-behavioural phenotype, **e15**

- cone shaped epiphyses**, Acrocapitofemoral dysplasia: an autosomal recessive skeletal dysplasia with cone shaped epiphyses in the hands and hips, **201**
- congenital cerulean cataract**, Gamma-D crystallin gene (*CRYGD*) mutation causes autosomal dominant congenital cerulean cataracts, **262**
- congenital heart defects**, Canine tricuspid valve malformation, a model of human Ebstein anomaly, maps to dog chromosome 9, **320**
Correlation between *PTPN11* gene mutations and congenital heart defects in Noonan and LEOPARD syndromes, **704**
High incidence of the R276X *SALL1* mutation in sporadic but not familial Townes-Brocks syndrome and report of the first familial case, **e127**
- congenital heart disease**, Is growth hormone treatment beneficial or harmful in Costello syndrome?, **e74**
- congenital melanocytic naevi**, Mutational analysis of *N-ras*, *p53*, *CDKN2A (p16)*, *p14*, *CDK4*, and *MC1R* genes in human dysplastic melanocytic naevi, **e14**
- congenital myasthenic syndromes**, Possible founder effect of rapsyn N88K mutation and identification of novel rapsyn mutations in congenital myasthenic syndromes, **e81**
- congenital nuclear cataract**, A novel *GJA8* mutation in an Iranian family with progressive autosomal dominant congenital nuclear cataract, **e124**
- connective tissue**, Homozygosity mapping of a gene for arterial tortuosity syndrome to chromosome 20q13, **747**
- connexin26**, Contribution of connexin26 (*GJB2*) mutations and founder effect to non-syndromic hearing loss in India, **e68**
Uniparental disomy of chromosome 13q causing homozygosity for the 35delG mutation in the gene encoding connexin26 (*GJB2*) results in prelingual hearing impairment in two unrelated Spanish patients, **636**
- consanguinity**, Inbreeding and risk of late onset complex disease, **925**
- copper disorders**, A copper treatable Menkes disease mutation associated with defective trafficking of a functional Menkes copper ATPase, **290**
- copy number changes**, An alternative to FISH: detecting deletion and duplication carriers within 24 hours, **e113**
- corneodesmosin**, Haplotype analysis of distantly related populations implicates corneodesmosin in psoriasis susceptibility, **447**
- coronary artery disease**, Association of genetic variants in the HDL receptor, SR-B1, with abnormal lipids in women with coronary artery disease, **453**
Haemochromatosis (*HFE*) gene C282Y mutation and the risk of coronary artery disease and myocardial infarction: a study in 1279 patients undergoing coronary angiography, **e58**
- coronary heart disease**, A common variant in the *ABCA1* gene is associated with a lower risk for premature coronary heart disease in familial hypercholesterolaemia, **163**
- Costello syndrome**, Is growth hormone treatment beneficial or harmful in Costello syndrome?, **e74**
Is the locus for Costello syndrome on 11p?, **469**
Studies on the pathogenesis of Costello syndrome, **e37**
- Cowden syndrome**, *PTEN* hamartoma tumour syndrome: variability of an entity, **e111**
- Cowden's disease**, De novo germline *PTEN* mutation in a man with Lhermitte-Duclos disease which arose on the paternal chromosome and was transmitted to his child with polydactyly and Wormian bones, **e92**
- CpG island**, Differential targets of CpG island hypermethylation in primary and metastatic head and neck squamous cell carcinoma (HNSCC), **25**
- CRB1**, Study of the involvement of the *RGR*, *CRPB1*, and *CRB1* genes in the pathogenesis of autosomal recessive retinitis pigmentosa, **e89**
- Cree encephalitis**, Cree encephalitis is allelic with Aicardi-Goutières syndrome: implications for the pathogenesis of disorders of interferon alpha metabolism, **183**
- CRX**, Evidence of autosomal dominant Leber congenital amaurosis (LCA) underlain by a *CRX* heterozygous null allele, **e90**
- CYP11B2**, Gender specific association of aldosterone synthase gene polymorphism with renal survival in patients with IgA nephropathy, **372**
- CYP1B1**, *CYP1B1* gene analysis in primary congenital glaucoma in Indonesian and European patients, **e9**
- CYP21A2**, Mapping of a de novo unequal crossover causing a deletion of the steroid 21-hydroxylase (*CYP21A2*) gene and a non-functional hybrid tenascin-X (*TNXB*) gene, **e53**
- cystic fibrosis**, A genetic screening programme for Tay-Sachs disease and cystic fibrosis for Australian Jewish high school students, **e45**
Transcript analysis of the cystic fibrosis splicing mutation 1525-1G>A shows use of multiple alternative splicing sites and suggests a putative role of exonic splicing enhancers, **e88**
- cytochrome P4501B1**, *CYP1B1* gene analysis in primary congenital glaucoma in Indonesian and European patients, **e9**
- de novo mutation**, De novo germline *PTEN* mutation in a man with Lhermitte-Duclos disease which arose on the paternal chromosome and was transmitted to his child with polydactyly and Wormian bones, **e92**
- deafness**, A mutation in the gamma actin 1 (*ACTG1*) gene causes autosomal dominant hearing loss (DFNA20/26), **879**
Contribution of connexin26 (*GJB2*) mutations and founder effect to non-syndromic hearing loss in India, **e68**
Distinctive audiometric profile associated with DFNB21 alleles of *TECTA*, **360**
Genetic homogeneity and phenotypic variability among Ashkenazi Jews with Usher syndrome type III, **767**
Myhre syndrome: new reports, review, and differential diagnosis, **546**
Origins and frequencies of *SLC26A4 (PDS)* mutations in east and south Asians: global implications for the epidemiology of deafness, **242**
Pathogenic mutations but not polymorphisms in congenital and childhood onset autosomal recessive deafness disrupt the proteolytic activity of TMPRSS3, **629**
- deafness and vestibular disorder**, Subcellular localisation, secretion, and post-translational processing of normal cochlin, and of mutants causing the sensorineural deafness and vestibular disorder, DFNA9, **479**
- deafness genes**, Genetic information but not termination: pregnant women's attitudes and willingness to pay for carrier screening for deafness genes, **e80**
- del(7)(q34q35)**, Multicolour FISH fine mapping unravels an insertion as a complex chromosomal rearrangement involving six breakpoints and a 5.89 Mb large deletion, **e60**
- deletion**, Death in adults with Prader-Willi syndrome may be correlated with maternal uniparental disomy, **e63**
Deletion in the *ABL* gene resulting from a meiotic recombination of a maternal (3;22;9)(q22;q12;q34.1) translocation, **e6**
Molecular characterisation of a 3.5 Mb interstitial 14q deletion in a child with several phenotypic anomalies, **e47**
Molecular characterisation of the 22q13 deletion syndrome supports the role of haploinsufficiency of *SHANK3/PROSAP2* in the major neurological symptoms, **575**
Mosaicism del(8p)/inv dup(8p) in a dysmorphic female infant: a mosaic formed by a meiotic error at the 8p OR gene and an independent terminal deletion event, **e93**
Multiple exon skipping and RNA circularisation contribute to the severe phenotypic expression of exon 5 dystrophin deletion, **e100**
Novel *PHF6* mutation p.D333del causes Börjeson-Forsman-Lehmann syndrome, **e50**
Prader-Willi syndrome: the psychopathological phenotype in uniparental disomy, **e112**
The severe form of type I hyperproliferative anaemia results from homozygous inactivation of the *PRODH* gene, **e7**
Using case study comparisons to explore genotype-phenotype correlations in Williams-Beuren syndrome, **136**
- deletion 22q13**, Telomeric 22q13 deletions resulting from rings, simple deletions, and translocations: cytogenetic, molecular, and clinical analyses of 32 new observations, **690**
- denaturing high performance liquid chromatography**, Association of single nucleotide polymorphisms in the oxidised LDL receptor 1 (*OLR1*) gene in patients with acute myocardial infarction, **933**
Comparison of fluorescent SSCP and denaturing HPLC analysis with direct sequencing for mutation screening in hypertrophic cardiomyopathy, **e59**
- Desbuquois dysplasia**, Homozygosity mapping of a Desbuquois dysplasia locus to chromosome 17q25.3, **282**
- development**, Gene expression signatures identify novel regulatory pathways during murine lung development: implications for lung tumorigenesis, **408**

- developmental delay**, Developmental delay and the methyl binding genes, **e13**
- DFNA20**, A mutation in the gamma actin 1 (*ACTG1*) gene causes autosomal dominant hearing loss (DFNA20/26), **879**
- DFNA26**, A mutation in the gamma actin 1 (*ACTG1*) gene causes autosomal dominant hearing loss (DFNA20/26), **879**
- DFNA43**, Mapping of a new autosomal dominant non-syndromic hearing loss locus (DFNA43) to chromosome 2p12, **278**
- DFNA49**, DFNA49, a novel locus for autosomal dominant non-syndromic hearing loss, maps proximal to DFNA7/DFNM1 region on chromosome 1q21-q23, **832**
- DFNA9**, Subcellular localisation, secretion, and post-translational processing of normal cochlin, and of mutants causing the sensorineural deafness and vestibular disorder, DFNA9, **479**
- DFNB1**, Uniparental disomy of chromosome 13q causing homozygosity for the 35delG mutation in the gene encoding connexin26 (*GJB2*) results in prelingual hearing impairment in two unrelated Spanish patients, **636**
- DFNB4**, Origins and frequencies of *SLC26A4* (*PDS*) mutations in east and south Asians: global implications for the epidemiology of deafness, **242**
- DFNB8/DFNB10**, Pathogenic mutations but not polymorphisms in congenital and childhood onset autosomal recessive deafness disrupt the proteolytic activity of *TMPRSS3*, **629**
- DHPLC**, A novel germline mutation in the *MET* extracellular domain in a Korean patient with the diffuse type of familial gastric cancer, **e97**
- diabetes mellitus**, Wolcott-Rallison syndrome: pathogenic insights into neonatal diabetes from new mutation and expression studies of *EIF2AK3*, **685**
- diagnostics**, Telomeres: a diagnosis at the end of the chromosomes, **385**
- digenic inheritance**, Further support for digenic inheritance in Bardet-Biedl syndrome, **e104**
- disability**, Melatonin supplementation for severe and intractable sleep disturbance in young people with genetically determined developmental disabilities: short review and commentary, **793**
- discordance**, Appraisal of genetic and epigenetic congruity of a monozygotic twin pair discordant for schizophrenia, **e16**
- DLL3**, Novel mutations in *DLL3*, a somitogenesis gene encoding a ligand for the Notch signalling pathway, cause a consistent pattern of abnormal vertebral segmentation in spondylocostal dysostosis, **333**
- DNA damage**, Absence of constitutional *H2AX* gene mutations in 101 hereditary breast cancer families, **e51**
- DNA diagnosis**, The use of genotype-phenotype correlations in mutation analysis for the long QT syndrome, **141**
- DNA double-strand break repair**, Mutation screening of *Mre11* complex genes: indication of *RAD50* involvement in breast and ovarian cancer susceptibility, **e131**
- DNA methylation**, Differential targets of CpG island hypermethylation in primary and metastatic head and neck squamous cell carcinoma (HNSCC), **25**
- DNA microarrays**, Array painting: a method for the rapid analysis of aberrant chromosomes using DNA microarrays, **664**
- DNA repair**, Fanconi anaemia, **1**
- dominant transmission**, Molecular study of three cases of odontohypophosphatasia resulting from heterozygosity for mutations in the tissue non-specific alkaline phosphatase gene, **605**
- DTDST**, Recessive multiple epiphyseal dysplasia (rMED): phenotype delineation in eighteen homozygotes for *DTDST* mutation R279W, **65**
- Duchenne muscular dystrophy**, An alternative to FISH: detecting deletion and duplication carriers within 24 hours, **e113**
- duplication**, Analphoid de novo marker chromosome inv dup(3)(q28qter) with neocentromere in a dysmorphic and developmentally retarded girl, **e27**
The metabotropic glutamate receptor 8 gene at 7q31: partial duplication and possible association with autism, **e42**
- dysmorphology**, Rapp-Hodgkin and AEC syndromes due to a new frameshift mutation in the *TP63* gene, **e133**
- dysplastic naevi mutation**, Mutational analysis of *N-ras*, *p53*, *CDKN2A* (*p16*), *p14*, *CDK4*, and *MC1R* genes in human dysplastic melanocytic naevi, **e14**
- dystonia**, Neuroferritinopathy in a French family with late onset dominant dystonia, **e69**
- dystrophin**, A complex deletion-inversion-deletion event results in a chimeric *IL1RAPL1*-dystrophin transcript and a contiguous gene deletion syndrome, **127**
Multiple exon skipping and RNA circularisation contribute to the severe phenotypic expression of exon 5 dystrophin deletion, **e100**
- dystrophin-glycoprotein complex**, Genotype-phenotype correlations in 35 Brazilian families with sarcoglycanopathies including the description of three novel mutations, **e12**
- dystrophy**, The genetics of inherited macular dystrophies, **641**
- DYX3**, A genome scan for developmental dyslexia confirms linkage to chromosome 2p11 and suggests a new locus on 7q32, **340**
- early diagnosis**, Cancer risk in 348 French *MSH2* or *MLH1* gene carriers, **208**
- early onset retinal dystrophy**, A Tyr368His *RPE65* founder mutation is associated with variable expression and progression of early onset retinal dystrophy in 10 families of a genetically isolated population, **709**
- early-onset**, *BRCA1* c.2845insA is a recurring mutation with a founder effect in Singapore Malay women with early onset breast/ovarian cancer, **e117**
- ectodermal dysplasia**, Hereditary xanthinuria type II associated with mental delay, autism, cortical renal cysts, nephrocalcinosis, osteopenia, and hair and teeth defects, **e121**
- EEC syndrome**, *P63* mutations are not a major cause of non-syndromic split hand/foot malformation, **55**
- Ehlers-Danlos syndrome**, Mapping of a de novo unequal crossover causing a deletion of the steroid 21-hydroxylase (*CYP21A2*) gene and a non-functional hybrid tenascin-X (*TNXB*) gene, **e53**
- Emery-Dreifuss muscular dystrophy**, Mutation analysis of the lamin A/C gene (*LMNA*) among patients with different cardiomyopathy phenotypes, **e132**
- EN2**, No association between the *EN2* gene and autistic disorder, **e4**
- enamel pitting**, Novel *ENAM* mutation responsible for autosomal recessive amelogenesis imperfecta and localised enamel defects, **900**
- enamelin**, Novel *ENAM* mutation responsible for autosomal recessive amelogenesis imperfecta and localised enamel defects, **900**
- endoglin**, Hereditary haemorrhagic telangiectasia: a questionnaire based study to delineate the different phenotypes caused by endoglin and *ALK1* mutations, **585**
- epigenetics**, Silencing of *CDKN1C* (*p57^{KIP2}*) is associated with hypomethylation at KvDMR1 in Beckwith-Wiedemann syndrome, **797**
- epilepsy**, Genetic mapping of a new Lafora progressive myoclonus epilepsy locus (*EPM2B*) on 6p22, **671**
- essential hypertension**, Evidence for linkage and association of the markers near the *LPL* gene with hypertension in Chinese families, **e57**
- expression signatures**, Gene expression signatures identify novel regulatory pathways during murine lung development: implications for lung tumorigenesis, **408**
- eye humour**, Expression of cell surface transmembrane carbonic anhydrase genes *CA9* and *CA12* in the human eye: overexpression of *CA12* (*CAXII*) in glaucoma, **257**
- Fabry disease**, Clinical presentation in female patients with Fabry disease, **e38**
Fabry disease: molecular studies in Italian patients and X inactivation analysis in manifesting carriers, **e103**
- FACL4**, A third MRX family (MRX68) is the result of mutation in the long chain fatty acid-CoA ligase 4 (*FACL4*) gene: proposal of a rapid enzymatic assay for screening mentally retarded patients, **11**
- false negative FISH result**, Partial deletion of the critical 1.5 Mb interval in Williams-Beuren syndrome, **e99**
- familial**, Familial vestibulocerebellar disorder maps to chromosome 13q31-q33: a new nystagmus locus, **37**
- familial adenomatous polyposis**, Cancer surveillance is often inadequate in people at high risk for colorectal cancer, **e54**
- familial breast cancer**, Women's preferences and consultants' communication of risk in consultations about familial breast cancer: impact on patient outcomes, **e56**
- familial hypercholesterolaemia**, A common variant in the *ABCA1* gene is associated with a lower risk for premature coronary heart disease in familial hypercholesterolaemia, **163**
Quality of life in a family based genetic cascade screening programme for familial hypercholesterolaemia: a longitudinal study among participants, **e3**
- familial Sotos syndrome**, Familial Sotos syndrome is caused by a novel 1 bp deletion of the *NSD1* gene, **51**

- familial tumours**, Hereditary pheochromocytomas and paragangliomas: a study of five susceptibility genes, **e75**
- family**, Evidence for linkage and association of the markers near the *LDL* gene with hypertension in Chinese families, **e57**
- family history**, Prevalence of family histories of breast cancer in the general population and the incidence of related seeking of health care, **e83**
- family studies**, The effect of sequence variations within the coding region of the C1 inhibitor gene on disease expression and protein function in families with hereditary angio-oedema, **e114**
- Fanconi anaemia**, Fanconi anaemia, **1**
- fetal hypokinesia**, Early onset asymmetrical intrauterine growth retardation with fetal hypokinesia and variable expression of acral and genitourinary malformations: a new lethal MCA syndrome, **e1**
- FGFR3**, Studies on the pathogenesis of Costello syndrome, **e37**
- fibrillin-1 gene**, In frame fibrillin-1 gene deletion in autosomal dominant Weill-Marchesani syndrome, **34**
- fibrocystic**, Molecular pathology and genetics of congenital hepatorenal fibrocystic syndromes, **311**
- FISH**, Distal trisomy 6p and 20q owing to the concurrent transposition of distal 6p and 20q to the 22q telomere: a genomic polymorphism?, **e94**
- SHOX* mutations detected by FISH and direct sequencing in patients with short stature, **e11**
- FMRP expression**, FMRP expression studies in blood and hair roots in a fragile X family with methylation mosaics, **535**
- FMRP test**, Predictive testing for cognitive functioning in female carriers of the fragile X syndrome using hair root analysis, **377**
- focal nullisomy**, Neuronal migration defect in a *BRCA1* gene carrier: possible focal nullisomy?, **e24**
- founder**, *BRCA1* c.2845insA is a recurring mutation with a founder effect in Singapore Malay women with early onset breast/ovarian cancer, **e117**
- founder effect**, Contribution of connexin26 (*GJB2*) mutations and founder effect to non-syndromic hearing loss in India, **e68**
- founder mutations**, The *RNASEL* 471delAAAG allele and prostate cancer in Ashkenazi Jewish men, **e22**
- FOXP2**, A genome scan for developmental dyslexia confirms linkage to chromosome 2p11 and suggests a new locus on 7q32, **340**
- fragile X syndrome**, FMRP expression studies in blood and hair roots in a fragile X family with methylation mosaics, **535**
- Modulation of methylation in the *FMR1* promoter region after long term treatment with L-carnitine and acetyl-L-carnitine, **1**
- Predictive testing for cognitive functioning in female carriers of the fragile X syndrome using hair root analysis, **377**
- Fukutin**, A homozygous nonsense mutation in the *Fukutin* gene causes a Walker-Warburg syndrome phenotype, **845**
- fumarase**, No fumarate hydratase (*FH*) mutations in hereditary prostate cancer, **e19**
- functional analysis**, Heterogeneity in residual function of MeCP2 carrying missense mutations in the methyl CpG binding domain, **487**
- G13513A**, The mitochondrial DNA G13513A MELAS mutation in the NADH dehydrogenase 5 gene is a frequent cause of Leigh-like syndrome with isolated complex I deficiency, **188**
- γ -crystallin**, Gamma-D crystallin gene (*CRYGD*) mutation causes autosomal dominant congenital cerulean cataracts, **267**
- gamete selection**, Non-random transmission of mutant alleles to female offspring of *BRCA1* carriers in Poland, **719**
- gangliosides**, Phenotypic and genetic exploration of severe demyelinating and secondary axonal neuropathies resulting from *GDAP1* nonsense and splicing mutations, **e87**
- gastric cancer**, A novel germline mutation in the *MET* extracellular domain in a Korean patient with the diffuse type of familial gastric cancer, **e97**
- GDAP1**, Phenotypic and genetic exploration of severe demyelinating and secondary axonal neuropathies resulting from *GDAP1* nonsense and splicing mutations, **e87**
- gender**, Gender specific association of aldosterone synthase gene polymorphism with renal survival in patients with IgA nephropathy, **372**
- gene**, Familial vestibulocerebellar disorder maps to chromosome 13q31-q33: a new nystagmus locus, **37**
- Genetic mapping of a new Lafora progressive myoclonus epilepsy locus (*EPM2B*) on 6p22, **671**
- gene conversion**, In vivo reversion to normal of inherited mutations in humans, **721**
- gene discovery**, Disruption of a novel member of a sodium/hydrogen exchanger family and *DOCK3* is associated with an attention deficit hyperactivity disorder-like phenotype, **733**
- gene dosage**, Diagnosis of gene dosage alterations at the *PMP22* gene using MAPH, **e123**
- gene expression**, Microarray analysis of gene/transcript expression in Prader-Willi syndrome: deletion versus UPD, **568**
- Wolcott-Rallison syndrome: pathogenic insights into neonatal diabetes from new mutation and expression studies of *EIF2AK3*, **685**
- gene mutation**, Novel sarcoglycan gene mutations in a large cohort of Italian patients, **e67**
- gene promoter**, Single nucleotide polymorphic alleles in the 5' region of the *RET* proto-oncogene define a risk haplotype in Hirschsprung's disease, **714**
- gene rearrangement**, Application of haplotype pair analysis for the identification of hemizygous loci, **346**
- gene silencing**, Silencing of *CDKN1C* (*p57^{KIP2}*) is associated with hypomethylation at KvDMR1 in Beckwith-Wiedemann syndrome, **797**
- gene-infectious agent interactions**, Does apolipoprotein E polymorphism influence susceptibility to malaria?, **348**
- general population**, Prevalence of family histories of breast cancer in the general population and the incidence of related seeking of health care, **e83**
- genes**, Association of genetic variants in the HDL receptor, SR-B1, with abnormal lipids in women with coronary artery disease, **453**
- genetic counselling**, Little value from including cousins in individual risk assessment of hereditary breast cancer: a simulation study, **e25**
- Mutation analysis in the *MECP2* gene and genetic counselling for Rett syndrome, **380**
- Non-random transmission of mutant alleles to female offspring of *BRCA1* carriers in Poland, **719**
- genetic disease**, A copper treatable Menkes disease mutation associated with defective trafficking of a functional Menkes copper ATPase, **290**
- genetic heterogeneity**, Gamma-D crystallin gene (*CRYGD*) mutation causes autosomal dominant congenital cerulean cataracts, **262**
- Identification of a mutation in the Indian Hedgehog (*IHH*) gene causing brachydactyly type A1 and evidence for a third locus, **42**
- In frame fibrillin-1 gene deletion in autosomal dominant Weill-Marchesani syndrome, **34**
- Mutational and gross deletion study of the *MENT1* gene and correlation with clinical features in Spanish patients, **e72**
- Novel *ENAM* mutation responsible for autosomal recessive amelogenesis imperfecta and localised enamel defects, **900**
- genetic isolate**, A Tyr368His *RPE65* founder mutation is associated with variable expression and progression of early onset retinal dystrophy in 10 families of a genetically isolated population, **709**
- Inbreeding and risk of late onset complex disease, **925**
- genetic linkage**, A genome scan for developmental dyslexia confirms linkage to chromosome 2p11 and suggests a new locus on 7q32, **340**
- genetic screening**, Quality of life in a family based genetic cascade screening programme for familial hypercholesterolaemia: a longitudinal study among participants, **e3**
- genetic testing**, Cancer surveillance is often inadequate in people at high risk for colorectal cancer, **e54**
- genetic variant**, Effect of the peroxisome proliferator activated receptor- γ gene Pro12Ala variant on body mass index: a meta-analysis, **773**
- genetics**, A second heterozygous *MDR3* nonsense mutation associated with intrahepatic cholestasis of pregnancy, **e32**
- Canine tricuspid valve malformation, a model of human Ebstein anomaly, maps to dog chromosome 9, **320**
- Genetic study of transthyretin amyloid neuropathies: carrier risks among French and Portuguese families, **e120**
- Melatonin supplementation for severe and intractable sleep disturbance in young people with genetically determined developmental disabilities: short review and commentary, **793**
- The genetics of inherited macular dystrophies, **641**
- genitourinary malformations**, Early onset asymmetrical intrauterine growth retardation with fetal hypokinesia and variable expression of acral and genitourinary malformations: a new lethal MCA syndrome, **e1**
- genomic imprinting**, Angelman syndrome: a review of the clinical and genetic aspects, **87**
- Genetic refinement and physical mapping of a biparental complete hydridiform mole locus on chromosome 19q13.4., **e95**

- The imprinted region on human chromosome 7q32 extends to the carboxypeptidase A gene cluster: an imprinted candidate for Silver-Russell syndrome, **249**
- genomic rearrangements**, Genomic rearrangements of *hMSH6* contribute to the genetic predisposition in suspected hereditary non-polyposis colorectal cancer syndrome, **597**
- genomic variants**, Searching for genomic variants in the *MEST11* transcript in Silver-Russell syndrome patients, **e65**
- genotype-phenotype**, Genotype-phenotype correlations for cataracts in neurofibromatosis 2, **758**
- High frequency of novel germline mutations in the *VHL* gene in the heterogeneous population of Brazil, **e31**
- genotype-phenotype correlation**, Distinctive audiometric profile associated with DFNB21 alleles of *TECTA*, **360**
- Genotype-phenotype correlations in 35 Brazilian families with sarcoglycanopathies including the description of three novel mutations, **e12**
- Independent *NF1* mutations in two large families with spinal neurofibromatosis, **122**
- Multicolour FISH fine mapping unravels an insertion as a complex chromosomal rearrangement involving six breakpoints and a 5.89 Mb large deletion, **e60**
- Partial deletion of the critical 1.5 Mb interval in Williams-Beuren syndrome, **e99**
- Spectrum of *NSD1* mutations in Sotos and Weaver syndromes, **436**
- The use of genotype-phenotype correlations in mutation analysis for the long QT syndrome, **141**
- genotype-phenotype relationship**, Evaluation of genotype-phenotype correlations in neurofibromatosis type 1, **e109**
- germ cells**, Y chromosome haplotypes and testicular cancer in the English population, **e20**
- germline mutation**, Cancer risk in 348 French *MSH2* or *MLH1* gene carriers, **208**
- γ -crystallin**, Gamma-D crystallin gene (*CRYGD*) mutation causes autosomal dominant congenital cerulean cataracts, **262**
- GJA8**, A novel *GJA8* mutation in an Iranian family with progressive autosomal dominant congenital nuclear cataract, **e124**
- GJB2**, Contribution of connexin26 (*GJB2*) mutations and founder effect to non-syndromic hearing loss in India, **e68**
- GLA deficiency**, Fabry disease: molecular studies in Italian patients and X inactivation analysis in manifesting carriers, **e103**
- glaucoma**, Expression of cell surface transmembrane carbonic anhydrase genes *CA9* and *CA12* in the human eye: overexpression of *CA12* (*CAXII*) in glaucoma, **257**
- Nail patella syndrome: a review of the phenotype aided by developmental biology, **153**
- Prevalence of optineurin sequence variants in adult primary open angle glaucoma: implications for diagnostic testing, **e101**
- The M98K variant of the OPTINEURIN (*OPTN*) gene modifies initial intraocular pressure in patients with primary open angle glaucoma, **842**
- glomerulonephritis**, Association of the *CD14* gene -159C polymorphism with progression of IgA nephropathy, **104**
- GLRA1**, A novel mutation (R218Q) at the boundary between the N-terminal and the first transmembrane domain of the glycine receptor in a case of sporadic hyperekplexia, **e71**
- Glutathione transferase**, Patient with bilateral periventricular nodular heterotopia and polymicrogyria with apparently balanced reciprocal translocation t(1;6)(p12;p12.2) that interrupts the mannosidase alpha, class 1A, and glutathione S-transferase A2 genes, **e128**
- glycerol kinase**, A complex deletion-inversion-deletion event results in a chimeric *IL1RAPL1*-dystrophin transcript and a contiguous gene deletion syndrome, **127**
- glycine receptor**, A novel mutation (R218Q) at the boundary between the N-terminal and the first transmembrane domain of the glycine receptor in a case of sporadic hyperekplexia, **e71**
- gonadosomic mosaicism**, Mitotic recombination mediated by the *JJAZF1* (*KIAA0160*) gene causing somatic mosaicism and a new type of constitutional *NF1* microdeletion in two children of a mosaic female with only few manifestations, **520**
- Greenberg dysplasia**, Greenberg dysplasia (HEM) and lethal X linked dominant Conradi-Hünermann chondrodysplasia punctata (CDPX2): presentation of two cases with overlapping phenotype, **e129**
- Greenberg/HEM skeletal dysplasia**, Congenital abnormalities reported in Pelger-Huët homozygosity as compared to Greenberg/HEM dysplasia: highly variable expression of allelic phenotypes, **937**
- GRM8**, The metabotropic glutamate receptor 8 gene at 7q31: partial duplication and possible association with autism, **e42**
- growth hormone**, Is growth hormone treatment beneficial or harmful in Costello syndrome?, **e74**
- H2AX**, Absence of constitutional *H2AX* gene mutations in 101 hereditary breast cancer families, **e51**
- haem oxygenase-1**, Relationship between microsatellite polymorphism in the haem oxygenase-1 gene promoter and longevity of the normal Japanese population, **146**
- haemochromatosis gene**, Haemochromatosis (*HFE*) gene C282Y mutation and the risk of coronary artery disease and myocardial infarction: a study in 1279 patients undergoing coronary angiography, **e58**
- Haemolytic uraemic syndrome**, Haemolytic uraemic syndrome and mutations of the factor H gene: a registry-based study of German speaking countries, **676**
- hair root**, FMRP expression studies in blood and hair roots in a fragile X family with methylation mosaics, **535**
- hair roots**, Predictive testing for cognitive functioning in female carriers of the fragile X syndrome using hair root analysis, **377**
- hand-foot-genital syndrome**, A novel duplication in the *HOXA13* gene in a family with atypical hand-foot-genital syndrome, **e49**
- haploinsufficiency**, Sotos syndrome and haploinsufficiency of *NSD1*: clinical features of intragenic mutations and submicroscopic deletions, **285**
- haplotype**, A common IL-1 complex haplotype is associated with an increased risk of atopy, **e66**
- Application of haplotype pair analysis for the identification of hemizygous loci, **346**
- haplotypes**, Chinese patients with sporadic Hirschsprung's disease are predominantly represented by a single *RET* haplotype, **e122**
- haptoglobin**, Haptoglobin and its association with the HELLP syndrome, **214**
- hearing**, Genetic homogeneity and phenotypic variability among Ashkenazi Jews with Usher syndrome type III, **767**
- hearing impairment**, Heteroplasmy for the 1555A>G mutation in the mitochondrial 12S rRNA gene in six Spanish families with non-syndromic hearing loss, **632**
- Uniparental disomy of chromosome 13q causing homozygosity for the 35delG mutation in the gene encoding connexin26 (*GJB2*) results in prelingual hearing impairment in two unrelated Spanish patients, **636**
- hearing loss**, Clinical and immunohistochemical evidence for an X linked retinitis pigmentosa syndrome with recurrent infections and hearing loss in association with an *RPGR* mutation, **e118**
- Mapping of a new autosomal dominant non-syndromic hearing loss locus (DFNA43) to chromosome 2p12, **278**
- RPGR* mutation associated with retinitis pigmentosa, impaired hearing, and sinorespiratory infections, **609**
- height**, Confirmation linkage study in support of the X chromosome harbouring a QTL underlying human height variation, **825**
- HELLP syndrome**, Haptoglobin and its association with the HELLP syndrome, **214**
- HEM**, Greenberg dysplasia (HEM) and lethal X linked dominant Conradi-Hünermann chondrodysplasia punctata (CDPX2): presentation of two cases with overlapping phenotype, **e129**
- hereditary breast cancer**, The *BRCA2* variant 8204G>A is a splicing mutation and results in an in frame deletion of the gene, **e23**
- hereditary breast-ovarian cancer**, Carrier risk status changes resulting from mutation testing in hereditary non-polyposis colorectal cancer and hereditary breast-ovarian cancer, **591**
- hereditary cancer syndrome**, Demographic and phenotypic features of 70 families segregating Barrett's oesophagus and oesophageal adenocarcinoma, **651**
- hereditary haemorrhagic telangiectasia**, Hereditary haemorrhagic telangiectasia: a questionnaire based study to delineate the different phenotypes caused by endoglin and *ALK1* mutations, **585**
- Molecular and functional analysis identifies *ALK-1* as the predominant cause of pulmonary hypertension related to hereditary haemorrhagic telangiectasia, **865**
- hereditary leiomyomatosis and renal cell cancer (HLRCC)**, No fumarate hydratase (*FH*) mutations in hereditary prostate cancer, **e19**

- hereditary motor and sensory neuropathies**, Phenotypic and genetic exploration of severe demyelinating and secondary axonal neuropathies resulting from *GDAP1* nonsense and splicing mutations, **e87**
- hereditary non-polyposis colorectal cancer**, Cancer surveillance is often inadequate in people at high risk for colorectal cancer, **e54**
Carrier risk status changes resulting from mutation testing in hereditary non-polyposis colorectal cancer and hereditary breast-ovarian cancer, **591**
Experience of discharge from colonoscopy of mutation negative HNPCC family members, **e55**
- hereditary pancreatitis**, *SPINK1* mutations and phenotypic expression in patients with pancreatitis associated with trypsinogen mutations, **e40**
- hereditary prostate cancer (HPRCA)**, No fumarate hydratase (*FH*) mutations in hereditary prostate cancer, **e19**
- hereditary spastic paraplegia**, Meta-analysis of age at onset in spastin-associated hereditary spastic paraplegia provides no evidence for a correlation with mutational class, **e106**
Science in motion: common molecular pathological themes emerge in the hereditary spastic paraplegias, **81**
- hereditary xanthinuria**, Hereditary xanthinuria type II associated with mental delay, autism, cortical renal cysts, nephrocalcinosis, osteopenia, and hair and teeth defects, **e121**
- heterogeneity**, New approaches to investigating heterogeneity in complex traits, **553**
- heterozygous null allele**, Evidence of autosomal dominant Leber congenital amaurosis (LCA) underlain by a *CRX* heterozygous null allele, **e90**
- high resolution breakpoint mapping**, Multicolour FISH fine mapping unravels an insertion as a complex chromosomal rearrangement involving six breakpoints and a 5.89 Mb large deletion, **e60**
- high resolution mapping**, A familial unbalanced subtelomeric translocation resulting in monosomy 6q27→qter, **e48**
- Hirschsprung disease**, Investigation of germline *GFRA4* mutations and evaluation of the involvement of *GFRA1*, *GFRA2*, *GFRA3*, and *GFRA4* sequence variants in *Hirschsprung disease*, **e18**
Mowat-Wilson syndrome, **305**
- Hirschsprung's disease**, Chinese patients with sporadic Hirschsprung's disease are predominantly represented by a single *RET* haplotype, **e122**
Single nucleotide polymorphic alleles in the 5' region of the *RET* proto-oncogene define a risk haplotype in Hirschsprung's disease, **714**
- histone**, Absence of constitutional *H2AX* gene mutations in 101 hereditary breast cancer families, **e51**
- history of neurofibromatosis**, From Aldrovandi's "Homuncio" (1592) to Buffon's girl (1749) and the "Wart Man" of Tilesius (1793): antique illustrations of mosaicism in neurofibromatosis?, **227**
- HLA class I polymorphism**, Significant frequency deviation of the class I polymorphism HLA-A10 in schizophrenic patients, **217**
- HLA immunogenetics**, Significant frequency deviation of the class I polymorphism HLA-A10 in schizophrenic patients, **217**
- hMSH6***, Genomic rearrangements of *hMSH6* contribute to the genetic predisposition in suspected hereditary non-polyposis colorectal cancer syndrome, **597**
- HNPCC**, Frequent genomic disorganisation of *MLH1* in hereditary non-polyposis colorectal cancer (HNPCC) screened by RT-PCR on puromycin treated samples, **e30**
Genomic rearrangements of *hMSH6* contribute to the genetic predisposition in suspected hereditary non-polyposis colorectal cancer syndrome, **597**
- HNPP**, Diagnosis of gene dosage alterations at the *PMP22* gene using MAPH, **e123**
- HNSSC**, Differential targets of CpG island hypermethylation in primary and metastatic head and neck squamous cell carcinoma (HNSSC), **25**
- Holt-Oram syndrome**, Mutations at the *SALL4* locus on chromosome 20 result in a range of clinically overlapping phenotypes, including Okhiro syndrome, Holt-Oram syndrome, acro-renal-ocular syndrome, and patients previously reported to represent thalidomide embryopathy, **473**
Novel *TBX5* mutations and molecular mechanism for Holt-Oram syndrome, **e29**
- homeobox**, Complete sequencing shows a role for *MSX1* in non-syndromic cleft lip and palate, **399**
- homozygosity**, Inbreeding and risk of late onset complex disease, **925**
- homozygosity mapping**, Genetic mapping of a new Lafora progressive myoclonus epilepsy locus (*EPM2B*) on 6p22, **671**
Homozygosity mapping of a Desbuquois dysplasia locus to chromosome 17q25.3, **282**
Homozygosity mapping of a gene for arterial tortuosity syndrome to chromosome 20q13, **747**
Mapping of a new locus for autosomal recessive non-syndromic mental retardation in the chromosomal region 19p13.12-p13.2: further genetic heterogeneity, **729**
- HOXA13***, A novel duplication in the *HOXA13* gene in a family with atypical hand-foot-genital syndrome, **e49**
- HPT-JT**, *HRPT2* mutations are associated with malignancy in sporadic parathyroid tumours, **657**
- HRPT2**, *HRPT2* mutations are associated with malignancy in sporadic parathyroid tumours, **657**
- hyperparathyroidism**, *HRPT2* mutations are associated with malignancy in sporadic parathyroid tumours, **657**
- hyperprolinaemia**, The severe form of type I hyperprolinaemia results from homozygous inactivation of the *PRODH* gene, **e7**
- hypersegmented neutrophils**, Hypogonadotrophic hypogonadism, short stature, cerebellar ataxia, rod-cone retinal dystrophy, and hypersegmented neutrophils: a novel disorder or a new variant of Boucher-Neuhauser syndrome?, **e2**
- hypertrophic cardiomyopathy**, Comparison of fluorescent SSCP and denaturing HPLC analysis with direct sequencing for mutation screening in hypertrophic cardiomyopathy, **e59**
- hypocomplementaemia**, Haemolytic uraemic syndrome and mutations of the factor H gene: a registry-based study of German speaking countries, **676**
- hypodontia**, A unique form of hypodontia seen in Vietnamese patients: clinical and molecular analysis, **e79**
- hypogonadism**, Hypogonadotrophic hypogonadism, short stature, cerebellar ataxia, rod-cone retinal dystrophy, and hypersegmented neutrophils: a novel disorder or a new variant of Boucher-Neuhauser syndrome?, **e2**
- hypophosphatasia**, Molecular study of three cases of odontohypophosphatasia resulting from heterozygosity for mutations in the tissue non-specific alkaline phosphatase gene, **605**
- hypospadias**, A novel duplication in the *HOXA13* gene in a family with atypical hand-foot-genital syndrome, **e49**
- IB**, High frequency of T9 and *CFTR* mutations in children with idiopathic bronchiectasis, **530**
- ICSI**, Beckwith-Wiedemann syndrome and assisted reproduction technology (ART), **62**
- idiopathic congenital central hypoventilation syndrome**, Association of germline mutations and polymorphisms of the *RET* proto-oncogene with idiopathic congenital central hypoventilation syndrome in 33 patients, **e10**
- IgA nephropathy**, Gender specific association of aldosterone synthase gene polymorphism with renal survival in patients with IgA nephropathy, **372**
Renoprotective efficacy of renin-angiotensin inhibitors in IgA nephropathy is influenced by *ACE* A2350G polymorphism, **e130**
- IHH**, Identification of a mutation in the Indian Hedgehog (*IHH*) gene causing brachydactyly type A1 and evidence for a third locus, **42**
- IHH gene**, Mild brachydactyly type A1 maps to chromosome 2q35-q36 and is caused by a novel *IHH* mutation in a three generation family, **132**
- IL-10***, Inflammation, genetics, and longevity: further studies on the protective effects in men of *IL-10* -1082 promoter SNP and its interaction with *TNF- α* -308 promoter SNP, **296**
- immunodeficiency**, Short-limbed dwarfism with bowing, combined immune deficiency, and late onset aplastic anaemia caused by novel mutations in the *RMPR* gene, **761**
- imprinting**, Beckwith-Wiedemann syndrome and assisted reproduction technology (ART), **62**
Death in adults with Prader-Willi syndrome may be correlated with maternal uniparental disomy, **e63**
Microarray analysis of gene/transcript expression in Prader-Willi syndrome: deletion versus UPD, **568**
- impulsivity**, A novel pericentric inversion of chromosome 3 cosegregates with a developmental-behavioural phenotype, **e15**
Disruption of a novel member of a sodium/hydrogen exchanger family and *DOCK3* is associated with an attention deficit hyperactivity disorder-like phenotype, **733**
- inbreeding**, Inbreeding and risk of late onset complex disease, **925**

- inflammation**, Inflammation, genetics, and longevity: further studies on the protective effects in men of *IL-10* -1082 promoter SNP and its interaction with *TNF- α* -308 promoter SNP, **296**
- inflammatory bowel disease**, Linkage of ulcerative colitis to the pericentromeric region of chromosome 16 in Italian inflammatory bowel disease families is independent of the presence of common *CARD15* mutations, **837**
- inheritance**, Non-random transmission of mutant alleles to female offspring of *BRCA1* carriers in Poland, **719**
- inherited**, The genetics of inherited macular dystrophies, **641**
- inherited predisposition**, *RNASEL* mutations in hereditary prostate cancer, **e21**
- INPP1**, Association of INPP1, PIK3CG, and TSC2 gene variants with autistic disorder: implications for phosphatidylinositol signalling in autism, **e119**
- interferon alpha**, Cree encephalitis is allelic with Aicardi-Goutières syndrome: implications for the pathogenesis of disorders of interferon alpha metabolism, **183**
- interleukin-1**, A common IL-1 complex haplotype is associated with an increased risk of atopy, **e66**
- interstitial duplication**, Partial trisomy of chromosome 22 resulting from an interstitial duplication of 22q11.2 in a child with typical cat eye syndrome, **e62**
- intrachromosomal recombination**, Heterogeneity of breakpoints in non-LCR-mediated large constitutional deletions of the 17q11.2 NF1 tumour suppressor region, **e116**
- intrauterine recombination**, In vivo reversion to normal of inherited mutations in humans, **721**
- intrauterine growth restriction**, Clinical aspects, prenatal diagnosis, and pathogenesis of trisomy 16 mosaicism, **175**
Small babies receive the cardiovascular protective apolipoprotein ϵ 2 allele less frequently than expected, **626**
- intrauterine growth retardation**, Early onset asymmetrical intrauterine growth retardation with fetal hypokinesia and variable expression of acral and genitourinary malformations: a new lethal MCA syndrome, **e1**
- inversion**, A complex deletion-inversion-deletion event results in a chimeric *IL1RAPL1*-dystrophin transcript and a contiguous gene deletion syndrome, **127**
- IQ**, FMRP expression studies in blood and hair roots in a fragile X family with methylation mosaics, **535**
- iron accumulation**, Karak syndrome: a novel degenerative disorder of the basal ganglia and cerebellum, **543**
- isolated hemihypertrophy**, Paternal uniparental disomy in monozygotic twins discordant for hemihypertrophy, **223**
- IVF**, Beckwith-Wiedemann syndrome and assisted reproduction technology (ART), **62**
- Jeune syndrome**, A locus for asphyxiating thoracic dystrophy, *ATD*, maps to chromosome 15q13, **431**
- Karak syndrome**, Karak syndrome: a novel degenerative disorder of the basal ganglia and cerebellum, **543**
- Kearns-Sayre syndrome**, Characterisation of repeat and palindrome elements in patients harbouring single deletions of mitochondrial DNA, **e86**
Identical large scale rearrangement of mitochondrial DNA causes Kearns-Sayre syndrome in a mother and her son, **858**
- KIAA0160 gene**, Mitotic recombination mediated by the *JJAZF1* (*KIAA0160*) gene causing somatic mosaicism and a new type of constitutional NF1 microdeletion in two children of a mosaic female with only few manifestations, **520**
- kidney**, Molecular pathology and genetics of congenital hepatorenal fibrocystic syndromes, **311**
- kidney disease**, Hereditary xanthinuria type II associated with mental delay, autism, cortical renal cysts, nephrocalcinosis, osteopenia, and hair and teeth defects, **e121**
- kinesin**, Science in motion: common molecular pathological themes emerge in the hereditary spastic paraplegias, **81**
- Krebs' cycle**, Mutation and LOH analysis of *ACO2* in colorectal cancer: no evidence of biallelic genetic inactivation, **e73**
- kyphomelic dysplasia**, Short-limbed dwarfism with bowing, combined immune deficiency, and late onset aplastic anaemia caused by novel mutations in the *RMPR* gene, **761**
- lamin A/C**, A synonymous codon change in the *LMNA* gene alters mRNA splicing and causes limb girdle muscular dystrophy type 1B, **e115**
Mutation analysis of the lamin A/C gene (*LMNA*) among patients with different cardiomyopathy phenotypes, **e132**
Expanding the phenotype of *LMNA* mutations in dilated cardiomyopathy and functional consequences of these mutations, **560**
Mandibuloacral dysplasia caused by homozygosity for the R527H mutation in lamin A/C, **854**
- lamin B receptor**, Greenberg dysplasia (HEM) and lethal X linked dominant Conradi-Hünemann chondrodysplasia punctata (CDPX2): presentation of two cases with overlapping phenotype, **e129**
- laminopathy**, Mandibuloacral dysplasia caused by homozygosity for the R527H mutation in lamin A/C, **854**
- large genomic disorganisation**, Frequent genomic disorganisation of *MLH1* in hereditary non-polyposis colorectal cancer (HNPCC) screened by RT-PCR on puromycin treated samples, **e30**
- large scale rearrangement**, Identical large scale rearrangement of mitochondrial DNA causes Kearns-Sayre syndrome in a mother and her son, **858**
- Laugier-Hunziker syndrome**, Laugier-Hunziker syndrome: an important differential diagnosis for Peutz-Jeghers syndrome, **e77**
- LCA**, Evidence of *RPGRIP1* gene mutations associated with recessive cone-rod dystrophy, **616**
- L-carnitine**, Modulation of methylation in the *FMRI* promoter region after long term treatment with L-carnitine and acetyl-L-carnitine, **1**
- LCRs at 17q11.2**, Mitotic recombination mediated by the *JJAZF1* (*KIAA0160*) gene causing somatic mosaicism and a new type of constitutional NF1 microdeletion in two children of a mosaic female with only few manifestations, **520**
- Leber congenital amaurosis**, Evidence of autosomal dominant Leber congenital amaurosis (LCA) underlain by a *CRX* heterozygous null allele, **e90**
- Leber hereditary optic neuropathy**, Frequency of rare mitochondrial DNA mutations in patients with suspected Leber's hereditary optic neuropathy, **e85**
The role of *APOE* in the phenotypic expression of Leber hereditary optic neuropathy, **e41**
- Leigh disease**, The mitochondrial DNA G13513A MELAS mutation in the NADH dehydrogenase 5 gene is a frequent cause of Leigh-like syndrome with isolated complex I deficiency, **188**
- LEOPARD syndrome**, Correlation between *PTPN11* gene mutations and congenital heart defects in Noonan and LEOPARD syndromes, **704**
- LGMD**, Novel sarcoglycan gene mutations in a large cohort of Italian patients, **e67**
- lifetime genetic testing**, Rapid, inexpensive scanning for all possible *BRCA1* and *BRCA2* gene sequence variants in a single assay: implications for genetic testing, **e33**
- likelihood ratio test**, Genetic variation analysis of *MLP*, *TFAP2A*, and *CSK* in patients with neural tube defects, **e43**
- limb girdle muscular dystrophy (LGMD1B)**, A synonymous codon change in the *LMNA* gene alters mRNA splicing and causes limb girdle muscular dystrophy type 1B, **e115**
- limb-girdle muscular dystrophy**, Genotype-phenotype correlations in 35 Brazilian families with sarcoglycanopathies including the description of three novel mutations, **e12**
- LMK1**, Using case study comparisons to explore genotype-phenotype correlations in Williams-Beuren syndrome, **136**
- linkage**, Autosomal dominant hereditary benign telangiectasia maps to the *CMC1* locus for capillary malformation on chromosome 5q14, **849**
Confirmation linkage study in support of the X chromosome harbouring a QTL underlying human height variation, **825**
Evidence for linkage and association of the markers near the *LPL* gene with hypertension in Chinese families, **e57**
Genetic mapping of a new Lafora progressive myoclonus epilepsy locus (*EPM2B*) on 6p22, **671**
- linkage analysis**, DFNA49, a novel locus for autosomal dominant non-syndromic hearing loss, maps proximal to DFNA7/DFNM1 region on chromosome 1q21-q23, **832**
Identification of a locus for type I punctate palmoplantar keratoderma on chromosome 15q22-q24, **872**
Linkage of ulcerative colitis to the pericentromeric region of chromosome 16 in Italian inflammatory bowel disease families is independent of the presence of common *CARD15* mutations, **837**
- lipoproteins**, Association of genetic variants in the HDL receptor, SR-B1, with abnormal lipids in women with coronary artery disease, **453**

- Lisch nodules**, Evaluation of genotype-phenotype correlations in neurofibromatosis type 1, **e109**
- lissencephaly**, A homozygous nonsense mutation in the *Fukutin* gene causes a Walker-Warburg syndrome phenotype, **845**
- liver**, Molecular pathology and genetics of congenital hepatorenal fibrocystic syndromes, **311**
- LMNA gene**, A synonymous codon change in the *LMNA* gene alters mRNA splicing and causes limb girdle muscular dystrophy type 1B, **e115**
- LMX1B**, Nail patella syndrome: a review of the phenotype aided by developmental biology, **153**
- LOH**, Alterations of the Birt-Hogg-Dubé gene (*BHD*) in sporadic colorectal tumours, **511**
- long QT syndrome**, The use of genotype-phenotype correlations in mutation analysis for the long QT syndrome, **141**
- longevity**, Inflammation, genetics, and longevity: further studies on the protective effects in men of *IL-10*-1082 promoter SNP and its interaction with *TNF- α* -308 promoter SNP, **296**
Relationship between microsatellite polymorphism in the haem oxygenase-1 gene promoter and longevity of the normal Japanese population, **146**
- longitudinal melanonychia**, Laugier-Hunziker syndrome: an important differential diagnosis for Peutz-Jeghers syndrome, **e77**
- loss of heterozygosity (LOH)**, Molecular study of frequency of mosaicism in neurofibromatosis 2 patients with bilateral vestibular schwannomas, **109**
- LPL gene**, Evidence for linkage and association of the markers near the *LPL* gene with hypertension in Chinese families, **e57**
- LTA**, Effect of *TNF* and *LTA* polymorphisms on biological markers of response to oxidative stimuli in coal miners: a model of gene-environment interaction, **96**
- lung**, Gene expression signatures identify novel regulatory pathways during murine lung development: implications for lung tumorigenesis, **408**
Single nucleotide polymorphism (SNP) analysis of mouse pulmonary adenoma susceptibility loci 1–4 for identification of candidate genes, **e36**
- Lynch syndromes**, Cancer risk in 348 French *MSH2* or *MLH1* gene carriers, **208**
- M98K**, The M98K variant of the OPTINEURIN (*OPTN*) gene modifies initial intraocular pressure in patients with primary open angle glaucoma, **842**
- malaria**, Does apolipoprotein E polymorphism influence susceptibility to malaria?, **348**
- Malay**, *BRCA1* c.2845insA is a recurring mutation with a founder effect in Singapore Malay women with early onset breast/ovarian cancer, **e117**
- male**, No correlation between phenotype and genotype in boys with a truncating *MECP2* mutation, **e5**
Sensitivity of *BRCA1/2* mutation testing in 466 breast/ovarian cancer families, **e107**
- male infertility**, The human Y chromosome's azoospermia factor b (*AZFb*) region: sequence, structure, and deletion analysis in infertile men, **18**
- malformation**, Clinical aspects, prenatal diagnosis, and pathogenesis of trisomy 16 mosaicism, **175**
- malignancy**, Is growth hormone treatment beneficial or harmful in Costello syndrome?, **e74**
- malignant melanoma**, Identification of a splice acceptor site mutation in *p16^{INK4A}/p14^{ARF}* within a breast cancer, melanoma, neurofibroma prone kindred, **e102**
- mandibuloacral dysplasia**, Mandibuloacral dysplasia caused by homozygosity for the R527H mutation in lamin A/C, **854**
- Mannosidase**, Patient with bilateral periventricular nodular heterotopia and polymicrogyria with apparently balanced reciprocal translocation t(1;6)(p12;p12.2) that interrupts the mannosidase alpha, class 1A, and glutathione S-transferase A2 genes, **e128**
- maternal inheritance**, Heteroplasmy for the 1555A>G mutation in the mitochondrial 12S rRNA gene in six Spanish families with non-syndromic hearing loss, **632**
- maternal uniparental disomy**, Chromosome 15 maternal uniparental disomy and psychosis in Prader-Willi syndrome, **72**
- MCA/MR syndrome**, Early onset asymmetrical intrauterine growth retardation with fetal hypokinesia and variable expression of acral and genitourinary malformations: a new lethal MCA syndrome, **e1**
- MCPH6**, A novel locus for autosomal recessive primary microcephaly (*MCPH6*) maps to 13q12.2, **540**
- MDR1**, *ABCB4* gene sequence variation in women with intrahepatic cholestasis of pregnancy, **e70**
- MDR3**, A second heterozygous *MDR3* nonsense mutation associated with intrahepatic cholestasis of pregnancy, **e32**
- MECP2**, Developmental delay and the methyl binding genes, **e13**
Mutation analysis in the *MECP2* gene and genetic counselling for Rett syndrome, **380**
No correlation between phenotype and genotype in boys with a truncating *MECP2* mutation, **e5**
- MECP2 mutation**, Heterogeneity in residual function of MeCP2 carrying missense mutations in the methyl CpG binding domain, **487**
- MECP2 mutations**, Patients with the R133C mutation: is their phenotype different from patients with Rett syndrome with other mutations?, **e52**
- meiotic error**, Mosaicism del(8p)/inv dup(8p) in a dysmorphic female infant: a mosaic formed by a meiotic error at the 8p *OR* gene and an independent terminal deletion event, **e93**
- melanoma**, Mutational analysis of *N-ras*, *p53*, *CDKN2A (p16)*, *p14*, *CDK4*, and *MCT1R* genes in human dysplastic melanocytic naevi, **e14**
- melatonin**, β_1 -adrenergic antagonists and melatonin reset the clock and restore sleep in a circadian disorder, Smith-Magenis syndrome, **74**
Melatonin supplementation for severe and intractable sleep disturbance in young people with genetically determined developmental disabilities: short review and commentary, **793**
- MEN 2**, Investigation of germline *GFRA4* mutations and evaluation of the involvement of *GFRA1*, *GFRA2*, *GFRA3*, and *GFRA4* sequence variants in Hirschsprung disease, **e18**
- MEN1**, Mutational and gross deletion study of the *MEN1* gene and correlation with clinical features in Spanish patients, **e72**
- MEN1 gene**, Mutational and gross deletion study of the *MEN1* gene and correlation with clinical features in Spanish patients, **e72**
- Menkes disease**, A copper treatable Menkes disease mutation associated with defective trafficking of a functional Menkes copper ATPase, **290**
- mental illness**, Disruption of the neuronal PAS3 gene in a family affected with schizophrenia, **325**
- mental retardation**, Cryptic terminal deletion of chromosome 9q34: a novel cause of syndromic obesity in childhood?, **300**
Diagnostic criteria, clinical characteristics, and natural history of Cohen syndrome, **233**
Identification of a 650 kb duplication at the X chromosome breakpoint in a patient with 46,X,t(X;8)(q28;q12) and non-syndromic mental retardation, **169**
Microcephaly-cardiomyopathy syndrome: expansion of the phenotype, **e78**
Mutations in the oligophrenin-1 gene (*OPHN1*) cause X linked congenital cerebellar hypoplasia, **441**
Progressive cerebellocerebral atrophy: a new syndrome with microcephaly, mental retardation, and spastic quadriplegia, **e96**
The severe form of type I hyperprolinaemia results from homozygous inactivation of the *PRODH* gene, **e7**
- MESTIT1**, Searching for genomic variants in the *MESTIT1* transcript in Silver-Russell syndrome patients, **e65**
- MET**, A novel germline mutation in the *MET* extracellular domain in a Korean patient with the diffuse type of familial gastric cancer, **e97**
- metaphyseal chondrodysplasia type Schmid**, Genetic changes in the RNA components of RNase MRP and RNase P in Schmid metaphyseal chondrodysplasia, **741**
- metastatic tumour**, Differential targets of CpG island hypermethylation in primary and metastatic head and neck squamous cell carcinoma (HNSCC), **25**
- meta-analysis**, Effect of the peroxisome proliferator activated receptor- γ gene Pro12Ala variant on body mass index: a meta-analysis, **773**
- methyl binding genes**, Developmental delay and the methyl binding genes, **e13**
- methylation**, Beckwith-Wiedemann syndrome and assisted reproduction technology (ART), **62**
Heterogeneity in residual function of MeCP2 carrying missense mutations in the methyl CpG binding domain, **487**
Modulation of methylation in the *FMR1* promoter region after long term treatment with L-carnitine and acetyl-L-carnitine, **1**
- microarray**, Gene expression signatures identify novel regulatory pathways during murine lung development: implications for lung tumorigenesis, **408**

- Microarray analysis of gene/transcript expression in Prader-Willi syndrome: deletion versus UPD, **568**
- microcephaly**, A novel locus for autosomal recessive primary microcephaly (MCPH6) maps to 13q12.2, **540**
- Molecular characterisation of a 3.5 Mb interstitial 14q deletion in a child with several phenotypic anomalies, **e47**
- Progressive cerebellocerebral atrophy: a new syndrome with microcephaly, mental retardation, and spastic quadriplegia, **e96**
- microcephaly-cardiomyopathy syndrome**, Microcephaly-cardiomyopathy syndrome: expansion of the phenotype, **e78**
- microfibrillopathies**, In frame fibrillin-1 gene deletion in autosomal dominant Weill-Marchesani syndrome, **34**
- microsatellite instability**, Genomic rearrangements of *hMSH6* contribute to the genetic predisposition in suspected hereditary non-polyposis colorectal cancer syndrome, **597**
- Mutations of the Birt-Hogg-Dubé (*BHD*) gene in sporadic colorectal carcinomas and colorectal carcinoma cell lines with microsatellite instability, **364**
- microsatellite polymorphism**, Relationship between microsatellite polymorphism in the haem oxygenase-1 gene promoter and longevity of the normal Japanese population, **146**
- mild phenotype**, Patients with the R133C mutation: is their phenotype different from patients with Rett syndrome with other mutations?, **e52**
- minigene**, Identification of a mutation that perturbs *NF1* gene splicing using genomic DNA samples and a minigene assay, **220**
- mitochondrial 12S rRNA**, Heteroplasmy for the 1555A>G mutation in the mitochondrial 12S rRNA gene in six Spanish families with non-syndromic hearing loss, **632**
- mitochondrial disease**, Identical large scale rearrangement of mitochondrial DNA causes Kearns-Sayre syndrome in a mother and her son, **858**
- mitochondrial DNA**, Frequency of rare mitochondrial DNA mutations in patients with suspected Leber's hereditary optic neuropathy, **e85**
- Identical large scale rearrangement of mitochondrial DNA causes Kearns-Sayre syndrome in a mother and her son, **858**
- The mitochondrial DNA G13513A MELAS mutation in the NADH dehydrogenase 5 gene is a frequent cause of Leigh-like syndrome with isolated complex I deficiency, **188**
- The role of *APOE* in the phenotypic expression of Leber hereditary optic neuropathy, **e41**
- mitochondrial DNA deletion syndrome**, Compensatory amplification of mtDNA in a patient with anovel deletion/duplication and high mutant load, **e125**
- mitochondrial DNA mutation**, A tRNA^{Ala} mutation causing mitochondrial myopathy clinically resembling myotonic dystrophy, **752**
- mitochondrial proliferation**, Compensatory amplification of mtDNA in a patient with anovel deletion/duplication and high mutant load, **e125**
- MLH1**, Frequent genomic disorganisation of *MLH1* in hereditary non-polyposis colorectal cancer (HNPCC) screened by RT-PCR on puromycin treated samples, **e30**
- models**, Evaluation of breast cancer risk assessment packages in the family history evaluation and screening programme, **807**
- molecular cytogenetics**, Deletion in the *ABL* gene resulting from a meiotic recombination of a maternal (3;22;9)(q22;q12;q34.1) translocation, **e6**
- molecular genetics**, Expanding the phenotype of *LMNA* mutations in dilated cardiomyopathy and functional consequences of these mutations, **560**
- monozygotic twins**, Appraisal of genetic and epigenetic congruity of a monozygotic twin pair discordant for schizophrenia, **e16**
- mosaicism**, From Aldrovandi's "Homuncio" (1592) to Buffon's girl (1749) and the "Wart Man" of Tilesius (1793): antique illustrations of mosaicism in neurofibromatosis?, **227**
- Molecular study of frequency of mosaicism in neurofibromatosis 2 patients with bilateral vestibular schwannomas, **109**
- Mosaicism del(8p)/inv dup(8p) in a dysmorphic female infant: a mosaic formed by a meiotic error at the 8p OR gene and an independent terminal deletion event, **e93**
- Somatic mosaicism in neurofibromatosis 2: prevalence and risk of disease transmission to offspring, **459**
- mouse**, Single nucleotide polymorphism (SNP) analysis of mouse pulmonary adenoma susceptibility loci 1-4 for identification of candidate genes, **e36**
- movement disorder**, Neuroferritinopathy in a French family with late onset dominant dystonia, **e69**
- Mowat-Wilson syndrome**, Characterisation of deletions of the *ZFX1B* region and genotype-phenotype analysis in Mowat-Wilson syndrome, **601**
- Mowat-Wilson syndrome, **305**
- MRE11, NBS1, and RAD50 germline mutations**, Mutation screening of Mre11 complex genes: indication of *RAD50* involvement in breast and ovarian cancer susceptibility, **e131**
- MRX**, Identification of a 650 kb duplication at the X chromosome breakpoint in a patient with 46,X,t(X;8)(q28;q12) and non-syndromic mental retardation, **169**
- MRX68**, A third MRX family (MRX68) is the result of mutation in the long chain fatty acid-CoA ligase 4 (*FACL4*) gene: proposal of a rapid enzymatic assay for screening mentally retarded patients, **11**
- MSI**, Alterations of the Birt-Hogg-Dubé gene (*BHD*) in sporadic colorectal tumours, **511**
- MSX1**, A unique form of hypodontia seen in Vietnamese patients: clinical and molecular analysis, **e79**
- Complete sequencing shows a role for *MSX1* in non-syndromic cleft lip and palate, **399**
- Linkage and linkage disequilibrium searched for between non-syndromic cleft palate and four candidate loci, **464**
- mtDNA**, Characterisation of repeat and palindrome elements in patients harbouring single deletions of mitochondrial DNA, **e86**
- mtDNA amplification**, Compensatory amplification of mtDNA in a patient with anovel deletion/duplication and high mutant load, **e125**
- mtDNA deletion**, Identical large scale rearrangement of mitochondrial DNA causes Kearns-Sayre syndrome in a mother and her son, **858**
- MTHFR**, Geographical and ethnic variation of the 677C>T allele of 5,10 methylenetetrahydrofolate reductase (*MTHFR*): findings from over 7000 newborns from 16 areas world wide, **619**
- mucocutaneous pigmentation**, Laugier-Hunziker syndrome: an important differential diagnosis for Peutz-Jeghers syndrome, **e77**
- mucopolysaccharidosis IIID**, Sanfilippo syndrome type D: identification of the first mutation in the N-acetylglucosamine-6-sulphatase gene, **192**
- "multiple" colorectal adenomas, Germline mutations in the TGF- β and Wnt signalling pathways are a rare cause of the "multiple" adenoma phenotype, **e35**
- multiple endocrine neoplasia**, Chromosome 2 (2p16) abnormalities in Carney complex tumours, **268**
- multiple epiphyseal dysplasia**, A recurrent R718W mutation in *COMP* results in multiple epiphyseal dysplasia with mild myopathy: clinical and pathogenetic overlap with collagen IX mutations, **942**
- Recessive multiple epiphyseal dysplasia (rMED): phenotype delineation in eighteen homozygotes for *DTDST* mutation R279W, **65**
- multiple sclerosis**, New approaches to investigating heterogeneity in complex traits, **553**
- multiplex amplifiable probe hybridisation**, An alternative to FISH: detecting deletion and duplication carriers within 24 hours, **e113**
- multiplex ligation dependent probe amplification (MLPA)**, Rapid, high throughput prenatal detection of aneuploidy using a novel quantitative method (MLPA), **907**
- muscular dystrophy**, A homozygous nonsense mutation in the *Fukutin* gene causes a Walker-Warburg syndrome phenotype, **845**
- Multiple exon skipping and RNA circularisation contribute to the severe phenotypic expression of exon 5 dystrophin deletion, **e100**
- Novel sarcoglycan gene mutations in a large cohort of Italian patients, **e67**
- mutation**, Association of germline mutations and polymorphisms of the *RET* proto-oncogene with idiopathic congenital central hypoventilation syndrome in 33 patients, **e10**
- High frequency of novel germline mutations in the *VHL* gene in the heterogeneous population of Brazil, **e31**
- Identification of eight novel *NSD1* mutations in Sotos syndrome, **e126**
- Independent *NF1* mutations in two large families with spinal neurofibromatosis, **122**
- Molecular study of frequency of mosaicism in neurofibromatosis 2 patients with bilateral vestibular schwannomas, **109**
- Wolcott-Rallison syndrome: pathogenic insights into neonatal diabetes from new mutation and expression studies of *EIF2AK3*, **685**
- mutation analysis**, Comparison of fluorescent SSCP and denaturing HPLC analysis with direct sequencing for mutation screening in hypertrophic cardiomyopathy, **e59**
- CYP1B1* gene analysis in primary congenital glaucoma in Indonesian and European patients, **e9**

- Novel *PHF6* mutation p.D333del causes Börjeson-Forsman-Lehmann syndrome, **e50**
- Sanfilippo syndrome type D: identification of the first mutation in the N-acetylglucosamine-6-sulphatase gene, **192**
- mutation screening**, An alternative to FISH: detecting deletion and duplication carriers within 24 hours, **e113**
- Spectrum of *NSD1* mutations in Sotos and Weaver syndromes, **436**
- mutational scanning**, Rapid, inexpensive scanning for all possible *BRCA1* and *BRCA2* gene sequence variants in a single assay: implications for genetic testing, **e33**
- mutational spectrum**, Mutations in *USH2A* in Spanish patients with autosomal recessive retinitis pigmentosa: high prevalence and phenotypic variation, **e8**
- mutations**, A synonymous codon change in the *LMNA* gene alters mRNA splicing and causes limb girdle muscular dystrophy type 1B, **e115**
- Molecular study of three cases of odontohypophosphatasia resulting from heterozygosity for mutations in the tissue non-specific alkaline phosphatase gene, **605**
- Myhre syndrome**, Myhre syndrome: new reports, review, and differential diagnosis, **546**
- myocardial infarction**, Association of single nucleotide polymorphisms in the oxidised LDL receptor 1 (*OLR1*) gene in patients with acute myocardial infarction, **933**
- Haemochromatosis (*HFE*) gene C282Y mutation and the risk of coronary artery disease and myocardial infarction: a study in 1279 patients undergoing coronary angiography, **e58**
- myoinositol**, Genetic variation analysis of *MLP*, *TFAP2A*, and *CSK* in patients with neural tube defects, **e43**
- myotonic dystrophy**, A tRNA^{Ala} mutation causing mitochondrial myopathy clinically resembling myotonic dystrophy, **752**
- N-acetylglucosamine-6-sulphatase**, Sanfilippo syndrome type D: identification of the first mutation in the N-acetylglucosamine-6-sulphatase gene, **192**
- NADH**, The mitochondrial DNA G13513A MELAS mutation in the NADH dehydrogenase 5 gene is a frequent cause of Leigh-like syndrome with isolated complex I deficiency, **188**
- nail patella syndrome**, Nail patella syndrome: a review of the phenotype aided by developmental biology, **153**
- neocentromere**, Anaphoid de novo marker chromosome inv dup(3)(q28qter) with neocentromere in a dysmorphic and developmentally retarded girl, **e27**
- neural tube defects**, Geographical and ethnic variation of the 677C>T allele of 5,10 methylenetetrahydrofolate reductase (*MTHFR*): findings from over 7000 newborns from 16 areas world wide, **619**
- neurobeachin**, The neurobeachin gene is disrupted by a translocation in a patient with idiopathic autism, **352**
- neurocristopathy**, Investigation of germline *GFRA4* mutations and evaluation of the involvement of *GFRA1*, *GFRA2*, *GFRA3*, and *GFRA4* sequence variants in Hirschsprung disease, **e18**
- neuroferritinopathy**, Neuroferritinopathy in a French family with late onset dominant dystonia, **e69**
- neurofibroma**, Identification of a splice acceptor site mutation in *p16^{INK4A}/p14^{ARF}* within a breast cancer, melanoma, neurofibroma prone kindred, **e102**
- neurofibromas**, Evaluation of genotype-phenotype correlations in neurofibromatosis type 1, **e109**
- neurofibromatosis 2**, Genotype-phenotype correlations for cataracts in neurofibromatosis 2, **758**
- Somatic mosaicism in neurofibromatosis 2: prevalence and risk of disease transmission to offspring, **459**
- neurofibromatosis 2 (NF2)**, Molecular study of frequency of mosaicism in neurofibromatosis 2 patients with bilateral vestibular schwannomas, **109**
- neurofibromatosis type 1**, From Aldrovandi's "Homuncio" (1592) to Buffon's girl (1749) and the "Wart Man" of Tilesius (1793): antique illustrations of mosaicism in neurofibromatosis?, **227**
- Independent *NF1* mutations in two large families with spinal neurofibromatosis, **122**
- Recurrent mutations in the *NF1* gene are common among neurofibromatosis type 1 patients, **e82**
- neuromuscular junction**, Possible founder effect of rapsyn N88K mutation and identification of novel rapsyn mutations in congenital myasthenic syndromes, **e81**
- neuronal migration defect**, Neuronal migration defect in a *BRCA1* gene carrier: possible focal nullisomy?, **e24**
- neuropathy**, Diagnosis of gene dosage alterations at the *PMP22* gene using MAPH, **e123**
- neutropenia**, Diagnostic criteria, clinical characteristics, and natural history of Cohen syndrome, **233**
- new approaches**, New approaches to investigating heterogeneity in complex traits, **553**
- NF1**, Evaluation of genotype-phenotype correlations in neurofibromatosis type 1, **e109**
- Heterogeneity of breakpoints in non-LCR-mediated large constitutional deletions of the 17q11.2 *NF1* tumour suppressor region, **e116**
- Identification of a mutation that perturbs *NF1* gene splicing using genomic DNA samples and a minigene assay, **220**
- NF1* mutations and clinical spectrum in patients with spinal neurofibromas, **368**
- Recurrent mutations in the *NF1* gene are common among neurofibromatosis type 1 patients, **e82**
- NF1 microdeletion**, Mitotic recombination mediated by the *JJAZF1* (*KIAA0160*) gene causing somatic mosaicism and a new type of constitutional *NF1* microdeletion in two children of a mosaic female with only few manifestations, **520**
- NF1 mutation**, *NF1* mutations and clinical spectrum in patients with spinal neurofibromas, **368**
- NF2**, Genotype-phenotype correlations for cataracts in neurofibromatosis 2, **758**
- Identification of recurrent regions of chromosome loss and gain in vestibular schwannomas using comparative genomic hybridisation, **802**
- Neurofibromatosis 2 in a patient with a *de novo* balanced reciprocal translocation 46,X,t(X;22)(p11.2;q11.2), **682**
- NF2 mutations**, Somatic mosaicism in neurofibromatosis 2: prevalence and risk of disease transmission to offspring, **459**
- non-carrier**, Experience of discharge from colonoscopy of mutation negative HNPCC family members, **e55**
- non-specific mental retardation**, Rare polymorphic variants of the *AGTR2* gene in boys with non-specific mental retardation, **357**
- non-specific X linked mental retardation**, A third MRX family (MRX68) is the result of mutation in the long chain fatty acid-CoA ligase 4 (*FACL4*) gene: proposal of a rapid enzymatic assay for screening mentally retarded patients, **11**
- nonsyndromic mental retardation**, Mapping of a new locus for autosomal recessive non-syndromic mental retardation in the chromosomal region 19p13.12-p13.2: further genetic heterogeneity, **729**
- non-syndromic sensorineural hearing loss**, DFNA49, a novel locus for autosomal dominant non-syndromic hearing loss, maps proximal to DFNA7/DFNM1 region on chromosome 1q21-q23, **832**
- Noonan syndrome**, Correlation between *PTPN11* gene mutations and congenital heart defects in Noonan and LEOPARD syndromes, **704**
- Notch signalling pathway**, Novel mutations in *DLL3*, a somitogenesis gene encoding a ligand for the Notch signalling pathway, cause a consistent pattern of abnormal vertebral segmentation in spondylocostal dysostosis, **333**
- novel mtDNA deletion**, Compensatory amplification of mtDNA in a patient with anovel deletion/duplication and high mutant load, **e125**
- NSD1**, Familial Sotos syndrome is caused by a novel 1 bp deletion of the *NSD1* gene, **51**
- Identification of eight novel *NSD1* mutations in Sotos syndrome, **e126**
- Sotos syndrome and haploinsufficiency of *NSD1*: clinical features of intragenic mutations and submicroscopic deletions, **285**
- Spectrum of *NSD1* mutations in Sotos and Weaver syndromes, **436**
- N-terminal domain**, A novel mutation (R218Q) at the boundary between the N-terminal and the first transmembrane domain of the glycine receptor in a case of sporadic hyperekplexia, **e71**
- NYS4**, Familial vestibulocerebellar disorder maps to chromosome 13q31-q33: a new nystagmus locus, **37**
- nystagmus**, Familial vestibulocerebellar disorder maps to chromosome 13q31-q33: a new nystagmus locus, **37**
- obesity**, APOE and TGF- β 1 genes are associated with obesity phenotypes, **918**
- Cryptic terminal deletion of chromosome 9q34: a novel cause of syndromic obesity in childhood?, **300**
- Effect of the peroxisome proliferator activated receptor- γ gene Pro12Ala variant on body mass index: a meta-analysis, **773**
- Okiihiro syndrome**, Mutations at the *SALL4* locus on chromosome 20 result in a range of clinically overlapping phenotypes, including Okiihiro syndrome, Holt-Oram syndrome, acro-renal-ocular

- syndrome, and patients previously reported to represent thalidomide embryopathy, **473**
- oligophrenin-1**, Mutations in the oligophrenin-1 gene (*OPHN1*) cause X linked congenital cerebellar hypoplasia, **441**
- OLR1**, Association of 3'-UTR polymorphisms of the oxidised LDL receptor 1 (*OLR1*) gene with Alzheimer's disease, **424**
- OLR1/LOX 1**, Association of single nucleotide polymorphisms in the oxidised LDL receptor 1 (*OLR1*) gene in patients with acute myocardial infarction, **933**
- oncogene**, Chromosome 2 (2p16) abnormalities in Carney complex tumours, **268**
- opsismodysplasia**, Clinical, radiological, and chondro-osseous findings in opsismodysplasia: survey of a series of 12 unreported cases, **195**
- optic atrophy**, Frequency of rare mitochondrial DNA mutations in patients with suspected Leber's hereditary optic neuropathy, **e85**
The role of *APOE* in the phenotypic expression of Leber hereditary optic neuropathy, **e41**
- OPTINEURIN**, The M98K variant of the OPTINEURIN (*OPTN*) gene modifies initial intraocular pressure in patients with primary open angle glaucoma, **842**
- optineurin mutations**, Prevalence of optineurin sequence variants in adult primary open angle glaucoma: implications for diagnostic testing, **e101**
- ossification delay**, Clinical, radiological, and chondro-osseous findings in opsismodysplasia: survey of a series of 12 unreported cases, **195**
- osteopetrosis**, A phenocopy of *CAII* deficiency: a novel genetic explanation for inherited infantile osteopetrosis with distal renal tubular acidosis, **115**
- otacoustic emissions**, Non-syndromic recessive auditory neuropathy is the result of mutations in the otoferlin (*OTOF*) gene, **45**
- otoferlin**, Non-syndromic recessive auditory neuropathy is the result of mutations in the otoferlin (*OTOF*) gene, **45**
- ovarian cancer**, Sensitivity of *BRCA1/2* mutation testing in 466 breast/ovarian cancer families, **e107**
- overgrowth**, Sotos syndrome and haploinsufficiency of *NSD1*: clinical features of intragenic mutations and submicroscopic deletions, **285**
- overgrowth syndromes**, Spectrum of *NSD1* mutations in Sotos and Weaver syndromes, **436**
- oxidative stress**, Effect of *TNF* and *LTA* polymorphisms on biological markers of response to oxidative stimuli in coal miners: a model of gene-environment interaction, **96**
Haptoglobin and its association with the HELLP syndrome, **214**
- P53**, Germline *TP53* mutations in breast cancer families with multiple primary cancers: is *TP53* a modifier of *BRCA1*?, **e34**
- P63**, *P63* mutations are not a major cause of non-syndromic split hand/foot malformation, **55**
- pachygyria**, Partial hexasomy 15pter→15q13 including *SNRPN* and *D15S10*: first molecular cytogenetically proven case report, **e28**
- palmoplantar keratoderma**, Identification of a locus for type I punctate palmoplantar keratoderma on chromosome 15q22-q24, **872**
- pancreatic cancer**, Identification of a splice acceptor site mutation in *p16^{INK4A}/p14^{ARF}* within a breast cancer, melanoma, neurofibroma prone kindred, **e102**
- paraganglioma**, Hereditary pheochromocytomas and paragangliomas: a study of five susceptibility genes, **e75**
- parathyroid carcinoma**, *HRPT2* mutations are associated with malignancy in sporadic parathyroid tumours, **657**
- parental origin**, Telomeric 22q13 deletions resulting from rings, simple deletions, and translocations: cytogenetic, molecular, and clinical analyses of 32 new observations, **690**
- partial deletion**, Partial deletion of the critical 1.5 Mb interval in Williams-Beuren syndrome, **e99**
- partial monosomy 6q**, A familial unbalanced subtelomeric translocation resulting in monosomy 6q27→qter, **e48**
- partial trisomy**, Partial trisomy of chromosome 22 resulting from an interstitial duplication of 22q11.2 in a child with typical cat eye syndrome, **e62**
- partial trisomy 16p**, A familial unbalanced subtelomeric translocation resulting in monosomy 6q27→qter, **e48**
- Pas1-4**, Single nucleotide polymorphism (SNP) analysis of mouse pulmonary adenoma susceptibility loci 1-4 for identification of candidate genes, **e36**
- paternal inheritance**, De novo germline *PTEN* mutation in a man with Lhermitte-Duclos disease which arose on the paternal chromosome and was transmitted to his child with polydactyly and Wormian bones, **e92**
- pathology review**, A twofold increase in *BRCA* mutation related prostate cancer among Ashkenazi Israelis is not associated with distinctive histopathology, **787**
- PAX2 gene**, Distal trisomy 6p and 20q owing to the concurrent transposition of distal 6p and 20q to the 22q telomere: a genomic polymorphism?, **e94**
- PAX9**, A unique form of hypodontia seen in Vietnamese patients: clinical and molecular analysis, **e79**
- PCAP locus**, No fumarate hydratase (*FH*) mutations in hereditary prostate cancer, **e19**
- Pelger-Huët homozygosity**, Congenital abnormalities reported in Pelger-Huët homozygosity as compared to Greenberg/HEM dysplasia: highly variable expression of allelic phenotypes, **937**
- Pendred syndrome**, Origins and frequencies of *SLC26A4* (*PDS*) mutations in east and south Asians: global implications for the epidemiology of deafness, **242**
- penetrance**, Genetic study of transthyretin amyloid neuropathies: carrier risks among French and Portuguese families, **e120**
SPINK1 mutations and phenotypic expression in patients with pancreatitis associated with trypsinogen mutations, **e40**
- Periventricular nodular heterotopia**, Patient with bilateral periventricular nodular heterotopia and polymicrogyria with apparently balanced reciprocal translocation t(1;6)(p12;p12.2) that interrupts the mannosidase alpha, class 1A, and glutathione S-transferase A2 genes, **e128**
- peroxisome proliferator activated receptor-γ**, Effect of the peroxisome proliferator activated receptor-γ gene Pro12Aa variant on body mass index: a meta-analysis, **773**
- Peutz-Jeghers syndrome**, Laugier-Hunziker syndrome: an important differential diagnosis for Peutz-Jeghers syndrome, **e77**
- pheochromocytoma**, Hereditary pheochromocytomas and paragangliomas: a study of five susceptibility genes, **e75**
- phenotype**, The genetics of inherited macular dystrophies, **641**
- phenotype/genotype correlation**, Hereditary haemorrhagic telangiectasia: a questionnaire based study to delineate the different phenotypes caused by endoglin and *ALK1* mutations, **585**
- phenotype-genotype correlations**, Telomeric 22q13 deletions resulting from rings, simple deletions, and translocations: cytogenetic, molecular, and clinical analyses of 32 new observations, **690**
- PHF6**, Novel *PHF6* mutation p.D333del causes Börjeson-Forsman-Lehmann syndrome, **e50**
- physical mapping**, Disruption of a novel member of a sodium/hydrogen exchanger family and *DOCK3* is associated with an attention deficit hyperactivity disorder-like phenotype, **733**
- pigmentary retinopathy**, Diagnostic criteria, clinical characteristics, and natural history of Cohen syndrome, **233**
- PIK3CG**, Association of *INPP1*, *PIK3CG*, and *TSC2* gene variants with autistic disorder: implications for phosphatidylinositol signalling in autism, **e119**
- PMP22**, Diagnosis of gene dosage alterations at the *PMP22* gene using *MAPH*, **e123**
- pneumoconiosis**, Effect of *TNF* and *LTA* polymorphisms on biological markers of response to oxidative stimuli in coal miners: a model of gene-environment interaction, **96**
- poly Q in AIB1**, Somatic instability of the DNA sequences encoding the polymorphic polyglutamine tract of the *AIB1* gene, **885**
- polyalanine tract**, A novel duplication in the *HoxA13* gene in a family with atypical hand-foot-genital syndrome, **e49**
- polycystic kidney disease**, Large deletion causing the *TSC2-PKD1* contiguous gene syndrome without infantile polycystic disease, **e17**
- polyglutamine tract in AIB1**, Somatic instability of the DNA sequences encoding the polymorphic polyglutamine tract of the *AIB1* gene, **885**
- Polymicrogyria**, Patient with bilateral periventricular nodular heterotopia and polymicrogyria with apparently balanced reciprocal translocation t(1;6)(p12;p12.2) that interrupts the mannosidase alpha, class 1A, and glutathione S-transferase A2 genes, **e128**
- polymorphism**, A common variant in the *ABCA1* gene is associated with a lower risk for premature coronary heart disease in familial hypercholesterolaemia, **163**
Association of 3'-UTR polymorphisms of the oxidised LDL receptor 1 (*OLR1*) gene with Alzheimer's disease, **424**
Association of germline mutations and polymorphisms of the *RET* proto-oncogene with idiopathic congenital central hypoventilation syndrome in 33 patients, **e10**

- Association of the *CD14* gene -159C polymorphism with progression of IgA nephropathy, **104**
- Germline *TP53* mutations in breast cancer families with multiple primary cancers: is *TP53* a modifier of *BRCA1*?, **e34**
- Maternal 677CT/1298AC genotype of the *MTHFR* gene as a risk factor for cleft lip, **e64**
- Renoprotective efficacy of renin-angiotensin inhibitors in IgA nephropathy is influenced by *ACE* A2350G polymorphism, **e130**
- Ser19→Trp polymorphism within the apolipoprotein AV gene in hypertriglyceridaemic people, **e105**
- population**, Geographical and ethnic variation of the 677C>T allele of 5,10 methylenetetrahydrofolate reductase (*MTHFR*): findings from over 7000 newborns from 16 areas world wide, **619**
- population based register**, Patients with the R133C mutation: is their phenotype different from patients with Rett syndrome with other mutations?, **e52**
- positional cloning**, The neurobeachin gene is disrupted by a translocation in a patient with idiopathic autism, **352**
- practice guidelines**, Little value from including cousins in individual risk assessment of hereditary breast cancer: a simulation study, **e25**
- Prader-Willi**, Prader-Willi syndrome: the psychopathological phenotype in uniparental disomy, **e112**
- Prader-Willi syndrome**, Chromosome 15 maternal uniparental disomy and psychosis in Prader-Willi syndrome, **72**
- Exclusion of maternal uniparental disomy of chromosome 14 in patients referred for Prader-Willi syndrome using a multiplex methylation polymerase chain reaction assay, **e46**
- Microarray analysis of gene/transcript expression in Prader-Willi syndrome: deletion versus UPD, **568**
- Prader-Willi/Angelman syndrome**, A supernumerary marker chromosome 15 tetrasomic for the Prader-Willi/Angelman syndrome critical region in a patient with a severe phenotype, **e84**
- predisposition**, Y chromosome haplotypes and testicular cancer in the English population, **e20**
- pre-eclampsia**, Haptoglobin and its association with the HELLP syndrome, **214**
- pregnancy**, A second heterozygous *MDR3* nonsense mutation associated with intrahepatic cholestasis of pregnancy, **e32**
- ABCB4* gene sequence variation in women with intrahepatic cholestasis of pregnancy, **e70**
- premature epimetaphyseal fusion**, Acrocapitofemoral dysplasia: an autosomal recessive skeletal dysplasia with cone shaped epiphyses in the hands and hips, **201**
- pre-mRNA**, Identification of a mutation that perturbs *NF1* gene splicing using genomic DNA samples and a minigene assay, **220**
- prenatal diagnosis**, Rapid, high throughput prenatal detection of aneuploidy using a novel quantitative method (MLPA), **907**
- pre-test probability assessment**, Pre-test prediction models of *BRCA1* or *BRCA2* mutation in breast/ovarian families attending familial cancer clinics, **503**
- primary congenital glaucoma**, *CYP11B1* gene analysis in primary congenital glaucoma in Indonesian and European patients, **e9**
- primary congenital lymphoedema**, Identification of eight novel *VEGFR-3* mutations in families with primary congenital lymphoedema, **697**
- primary tumour**, Differential targets of CpG island hypermethylation in primary and metastatic head and neck squamous cell carcinoma (HNSCC), **25**
- PRODH**, The severe form of type I hyperprolinaemia results from homozygous inactivation of the *PRODH* gene, **e7**
- progression**, Association of the *CD14* gene -159C polymorphism with progression of IgA nephropathy, **104**
- promoter polymorphism**, Promoter polymorphisms of the *CD14* gene in Italian patients with coeliac disease, **e108**
- prostate cancer**, A twofold increase in *BRCA* mutation related prostate cancer among Ashkenazi Israelis is not associated with distinctive histopathology, **787**
- A Y chromosomal influence on prostate cancer risk: the multi-ethnic cohort study, **815**
- BRCA1* and *BRCA2* mutations have no major role in predisposition to prostate cancer in Finland, **e98**
- RNASEL* mutations in hereditary prostate cancer, **e21**
- The *RNASEL* 471delAAAG allele and prostate cancer in Ashkenazi Jewish men, **e22**
- protease**, Pathogenic mutations but not polymorphisms in congenital and childhood onset autosomal recessive deafness disrupt the proteolytic activity of *TMPRSS3*, **629**
- protein trafficking**, A copper treatable Menkes disease mutation associated with defective trafficking of a functional Menkes copper ATPase, **290**
- Proteus syndrome**, *PTEN* hamartoma tumour syndrome: variability of an entity, **e11**
- Proteus-like syndrome**, *PTEN* hamartoma tumour syndrome: variability of an entity, **e111**
- proton pump**, A phenocopy of *CAII* deficiency: a novel genetic explanation for inherited infantile osteopetrosis with distal renal tubular acidosis, **115**
- pseudo-TORCH syndrome**, Cree encephalitis is allelic with Aicardi-Goutières syndrome: implications for the pathogenesis of disorders of interferon alpha metabolism, **183**
- psoriasis**, Haplotype analysis of distantly related populations implicates corneodesmosin in psoriasis susceptibility, **447**
- PSORS1**, Haplotype analysis of distantly related populations implicates corneodesmosin in psoriasis susceptibility, **447**
- psychiatric disorders**, Disruption of the neuronal *PAS3* gene in a family affected with schizophrenia, **325**
- psychiatric gene**, Disruption of the neuronal *PAS3* gene in a family affected with schizophrenia, **325**
- psychological impact**, Quality of life in a family based genetic cascade screening programme for familial hypercholesterolaemia: a longitudinal study among participants, **e3**
- psychosis**, Chromosome 15 maternal uniparental disomy and psychosis in Prader-Willi syndrome, **72**
- Prader-Willi syndrome: the psychopathological phenotype in uniparental disomy, **e112**
- PTEN**, De novo germline *PTEN* mutation in a man with Lhermitte-Duclos disease which arose on the paternal chromosome and was transmitted to his child with polydactyly and Wormian bones, **e92**
- PTEN hamartoma tumour syndrome**, *PTEN* hamartoma tumour syndrome: variability of an entity, **e111**
- PTPN11**, Correlation between *PTPN11* gene mutations and congenital heart defects in Noonan and LEOPARD syndromes, **704**
- pulmonary hypertension**, Molecular and functional analysis identifies ALK-1 as the predominant cause of pulmonary hypertension related to hereditary haemorrhagic telangiectasia, **865**
- puromycin**, Frequent genomic disorganisation of *MLH1* in hereditary non-polyposis colorectal cancer (HNPCC) screened by RT-PCR on puromycin treated samples, **e30**
- quality of life**, Quality of life in a family based genetic cascade screening programme for familial hypercholesterolaemia: a longitudinal study among participants, **e3**
- ragged red fibres**, A tRNA^{Ala} mutation causing mitochondrial myopathy clinically resembling myotonic dystrophy, **752**
- Rapp-Hodgkin syndrome**, Rapp-Hodgkin and AEC syndromes due to a new frameshift mutation in the *TP63* gene, **e133**
- RAPSN**, Possible founder effect of rapsyn N88K mutation and identification of novel rapsyn mutations in congenital myasthenic syndromes, **e81**
- rapsyn**, Possible founder effect of rapsyn N88K mutation and identification of novel rapsyn mutations in congenital myasthenic syndromes, **e81**
- RBMY**, The human Y chromosome's azoospermia factor b (AZFb) region: sequence, structure, and deletion congruity in infertile men, **18**
- RDA**, Appraisal of genetic and epigenetic congruity of a monozygotic twin pair discordant for schizophrenia, **e16**
- reading disability**, A genome scan for developmental dyslexia confirms linkage to chromosome 2p11 and suggests a new locus on 7q32, **340**
- receptors**, Association of genetic variants in the HDL receptor, SR-B1, with abnormal lipids in women with coronary artery disease, **453**
- recessive**, Molecular pathology and genetics of congenital hepatorenal fibrocystic syndromes, **311**
- recessive non-syndromic hearing loss**, Non-syndromic recessive auditory neuropathy is the result of mutations in the otoferlin (*OTOF*) gene, **45**
- recurrent mutation**, Recurrent mutations in the *NF1* gene are common among neurofibromatosis type 1 patients, **e82**
- renal**, Analysis of the Birt-Hogg-Dubé (*BHD*) tumour suppressor gene in sporadic renal cell carcinoma and colorectal cancer, **820**
- renal failure**, Nail patella syndrome: a review of the phenotype aided by developmental biology, **153**

- renal tubular acidosis**, A phenocopy of CAII deficiency: a novel genetic explanation for inherited infantile osteopetrosis with distal renal tubular acidosis, **115**
- Rendu-Osler-Weber**, Autosomal dominant hereditary benign telangiectasia maps to the *CMC1* locus for capillary malformation on chromosome 5q14, **849**
- repeat elements**, Characterisation of repeat and palindrome elements in patients harbouring single deletions of mitochondrial DNA, **e86**
- respiratory chain deficiency**, A tRNA^{Ala} mutation causing mitochondrial myopathy clinically resembling myotonic dystrophy, **752**
- respiratory tract infections**, Clinical and immunohistochemical evidence for an X linked retinitis pigmentosa syndrome with recurrent infections and hearing loss in association with an *RPGR* mutation, **e118**
- RET co-receptors**, Investigation of germline *GFRA4* mutations and evaluation of the involvement of *GFRA1*, *GFRA2*, *GFRA3*, and *GFRA4* sequence variants in Hirschsprung disease, **e18**
- RET gene**, Chinese patients with sporadic Hirschsprung's disease are predominantly represented by a single *RET* haplotype, **e122**
- RET proto-oncogene**, Association of germline mutations and polymorphisms of the *RET* proto-oncogene with idiopathic congenital central hypoventilation syndrome in 33 patients, **e10**
- Single nucleotide polymorphic alleles in the 5' region of the *RET* proto-oncogene define a risk haplotype in Hirschsprung's disease, **714**
- retardation**, Molecular characterisation of a 3.5 Mb interstitial 14q deletion in a child with several phenotypic anomalies, **e47**
- retinal dystrophies**, Evidence of *RPGRIP1* gene mutations associated with recessive cone-rod dystrophy, **616**
- retinitis pigmentosa**, Clinical and immunohistochemical evidence for an X linked retinitis pigmentosa syndrome with recurrent infections and hearing loss in association with an *RPGR* mutation, **e118**
- Genetic homogeneity and phenotypic variability among Ashkenazi Jews with Usher syndrome type III, **767**
- Mutations in *USH2A* in Spanish patients with autosomal recessive retinitis pigmentosa: high prevalence and phenotypic variation, **e8**
- Study of the involvement of the *RGR*, *CRPB1*, and *CRB1* genes in the pathogenesis of autosomal recessive retinitis pigmentosa, **e89**
- Rett syndrome**, Developmental delay and the methyl binding genes, **e13**
- Heterogeneity in residual function of MeCP2 carrying missense mutations in the methyl CpG binding domain, **487**
- Mutation analysis in the *MECP2* gene and genetic counselling for Rett syndrome, **380**
- No correlation between phenotype and genotype in boys with a truncating *MECP2* mutation, **e5**
- Patients with the R133C mutation: is their phenotype different from patients with Rett syndrome with other mutations?, **e52**
- reverse chromosome painting**, Array painting: a method for the rapid analysis of aberrant chromosomes using DNA microarrays, **664**
- RGR**, Study of the involvement of the *RGR*, *CRPB1*, and *CRB1* genes in the pathogenesis of autosomal recessive retinitis pigmentosa, **e89**
- rhabdomyosarcoma**, Is the locus for Costello syndrome on 11p?, **469**
- ring (22)**, Telomeric 22q13 deletions resulting from rings, simple deletions, and translocations: cytogenetic, molecular, and clinical analyses of 32 new observations, **690**
- risk**, Evaluation of breast cancer risk assessment packages in the family history evaluation and screening programme, **807**
- risk assessment**, Little value from including cousins in individual risk assessment of hereditary breast cancer: a simulation study, **e25**
- risk communication**, Women's preferences and consultants' communication of risk in consultations about familial breast cancer: impact on patient outcomes, **e56**
- RMRP**, Genetic changes in the RNA components of RNase MRP and RNase P in Schmid metaphyseal chondrodysplasia, **741**
- RNase MRP**, Genetic changes in the RNA components of RNase MRP and RNase P in Schmid metaphyseal chondrodysplasia, **741**
- RNase P**, Genetic changes in the RNA components of RNase MRP and RNase P in Schmid metaphyseal chondrodysplasia, **741**
- RNASEL**, The *RNASEL* 471delAAAG allele and prostate cancer in Ashkenazi Jewish men, **e22**
- rod-cone dystrophy**, Hypogonadotropic hypogonadism, short stature, cerebellar ataxia, rod-cone retinal dystrophy, and hypersegmented neutrophils: a novel disorder or a new variant of Boucher-Neuhauser syndrome?, **e2**
- RPE65 founder mutation**, A Tyr368His *RPE65* founder mutation is associated with variable expression and progression of early onset retinal dystrophy in 10 families of a genetically isolated population, **709**
- RPGR**, A new genetic locus for X linked progressive cone-rod dystrophy, **418**
- Clinical and immunohistochemical evidence for an X linked retinitis pigmentosa syndrome with recurrent infections and hearing loss in association with an *RPGR* mutation, **e118**
- RPGR gene (retinitis pigmentosa GTPase regulator)**, *RPGR* mutation associated with retinitis pigmentosa, impaired hearing, and sinorespiratory infections, **609**
- RPGRIP1**, Evidence of *RPGRIP1* gene mutations associated with recessive cone-rod dystrophy, **616**
- SALL1**, High incidence of the R276X *SALL1* mutation in sporadic but not familial Townes-Brocks syndrome and report of the first familial case, **e127**
- SALL4**, Mutations at the *SALL4* locus on chromosome 20 result in a range of clinically overlapping phenotypes, including Okhiro syndrome, Holt-Oram syndrome, acro-renal-ocular syndrome, and patients previously reported to represent thalidomide embryopathy, **473**
- Sanfilippo syndrome type D**, Sanfilippo syndrome type D: identification of the first mutation in the N-acetylglucosamine-6-sulphatase gene, **192**
- sarcoglycan**, Novel sarcoglycan gene mutations in a large cohort of Italian patients, **e67**
- sarcoglycanopathies**, Genotype-phenotype correlations in 35 Brazilian families with sarcoglycanopathies including the description of three novel mutations, **e12**
- schizophrenia**, Appraisal of genetic and epigenetic congruity of a monozygotic twin pair discordant for schizophrenia, **e16**
- Significant frequency deviation of the class I polymorphism HLA-A10 in schizophrenic patients, **217**
- schwannoma**, Identification of recurrent regions of chromosome loss and gain in vestibular schwannomas using comparative genomic hybridisation, **802**
- screening**, A genetic screening programme for Tay-Sachs disease and cystic fibrosis for Australian Jewish high school students, **e45**
- segmental heterodisomy**, Comprehensive microsatellite marker analysis contradicts previous report of segmental maternal heterodisomy of chromosome 14, **e26**
- segmental neurofibromatosis**, From Aldrovandi's "Homuncio" (1592) to Buffon's girl (1749) and the "Wart Man" of Tilesius (1793): antique illustrations of mosaicism in neurofibromatosis?, **227**
- seizures**, Partial hexasomy 15pter→15q13 including *SNRPN* and D15S10: first molecular cytogenetically proven case report, **e28**
- sequencing**, *SHOX* mutations detected by FISH and direct sequencing in patients with short stature, **e11**
- serpins**, The effect of sequence variations within the coding region of the C1 inhibitor gene on disease expression and protein function in families with hereditary angio-oedema, **e114**
- sex ratio**, Clinical aspects, prenatal diagnosis, and pathogenesis of trisomy 16 mosaicism, **175**
- SHANK3**, Molecular characterisation of the 22q13 deletion syndrome supports the role of haploinsufficiency of *SHANK3/PROSAP2* in the major neurological symptoms, **575**
- short limb dwarfism**, Acrocapitofemoral dysplasia: an autosomal recessive skeletal dysplasia with cone shaped epiphyses in the hands and hips, **201**
- short stature**, *SHOX* mutations detected by FISH and direct sequencing in patients with short stature, **e11**
- SHOX**, *SHOX* mutations detected by FISH and direct sequencing in patients with short stature, **e11**
- Silver-Russell syndrome**, Genomic characterisation of *C7orf10* in Silver-Russell syndrome patients, **e44**
- Searching for genomic variants in the *MESTIT1* transcript in Silver-Russell syndrome patients, **e65**
- Silver-Russell syndrome (SRS)**, The imprinted region on human chromosome 7q32 extends to the carboxypeptidase A gene cluster: an imprinted candidate for Silver-Russell syndrome, **249**
- single deletions**, Characterisation of repeat and palindrome elements in patients harbouring single deletions of mitochondrial DNA, **e86**
- single nucleotide polymorphism**, Application of haplotype pair analysis for the identification of hemizygous loci, **346**
- Single nucleotide polymorphic alleles in the 5' region of the *RET* proto-oncogene define a risk haplotype in Hirschsprung's disease, **714**
- single strand conformation polymorphism analysis**, Comparison of fluorescent SSCP and denaturing HPLC analysis with direct

- sequencing for mutation screening in hypertrophic cardiomyopathy, **e59**
- SIP1**, Characterisation of deletions of the *ZFX1B* region and genotype-phenotype analysis in Mowat-Wilson syndrome, **601**
- site-specific reversion**, In vivo reversion to normal of inherited mutations in humans, **721**
- skin prick test**, A common IL-1 complex haplotype is associated with an increased risk of atopy, **e66**
- SLC26A4**, Origins and frequencies of *SLC26A4* (*PDS*) mutations in east and south Asians: global implications for the epidemiology of deafness, **242**
- sleep**, β_1 -adrenergic antagonists and melatonin reset the clock and restore sleep in a circadian disorder, Smith-Magenis syndrome, **74**
Melatonin supplementation for severe and intractable sleep disturbance in young people with genetically determined developmental disabilities: short review and commentary, **793**
- SLFN gene cluster**, Heterogeneity of breakpoints in non-LCR-mediated large constitutional deletions of the 17q11.2 NF1 tumour suppressor region, **e116**
- SMA**, Prevalence of *SMN1* deletion and duplication in carrier and normal populations: implication for genetic counselling, **e39**
- Smith-Magenis syndrome**, β_1 -adrenergic antagonists and melatonin reset the clock and restore sleep in a circadian disorder, Smith-Magenis syndrome, **74**
- SMN quantification**, Prevalence of *SMN1* deletion and duplication in carrier and normal populations: implication for genetic counselling, **e39**
- SMN1 deletion**, Prevalence of *SMN1* deletion and duplication in carrier and normal populations: implication for genetic counselling, **e39**
- SMN1 duplication**, Prevalence of *SMN1* deletion and duplication in carrier and normal populations: implication for genetic counselling, **e39**
- SNPs**, Association of single nucleotide polymorphisms in the oxidised LDL receptor 1 (*OLR1*) gene in patients with acute myocardial infarction, **933**
- somatic mosaicism**, In vivo reversion to normal of inherited mutations in humans, **721**
- somitogenesis**, Novel mutations in *DLL3*, a somitogenesis gene encoding a ligand for the Notch signalling pathway, cause a consistent pattern of abnormal vertebral segmentation in spondylocostal dysostosis, **333**
- Sotos syndrome**, Identification of eight novel *NSD1* mutations in Sotos syndrome, **e126**
Sotos syndrome and haploinsufficiency of *NSD1*: clinical features of intragenic mutations and submicroscopic deletions, **285**
- Southern blot**, Mutational and gross deletion study of the *MEN1* gene and correlation with clinical features in Spanish patients, **e72**
- spasticity**, Progressive cerebellocerebral atrophy: a new syndrome with microcephaly, mental retardation, and spastic quadriplegia, **e96**
- spastin**, Meta-analysis of age at onset in spastin-associated hereditary spastic paraplegia provides no evidence for a correlation with mutational class, **e106**
Science in motion: common molecular pathological themes emerge in the hereditary spastic paraplegias, **81**
- spatial impairment**, Using case study comparisons to explore genotype-phenotype correlations in Williams-Beuren syndrome, **136**
- spina bifida**, Genetic variation analysis of *MLP*, *TFAP2A*, and *CSK* in patients with neural tube defects, **e43**
- spinal neurofibroma**, Heterogeneity of breakpoints in non-LCR-mediated large constitutional deletions of the 17q11.2 NF1 tumour suppressor region, **e116**
- spinal neurofibromatosis**, Independent *NF1* mutations in two large families with spinal neurofibromatosis, **122**
- spinal tumours**, *NF1* mutations and clinical spectrum in patients with spinal neurofibromas, **368**
- SPINK1**, *SPINK1* mutations and phenotypic expression in patients with pancreatitis associated with trypsinogen mutations, **e40**
- splice mutation**, The *BRCA2* variant 8204G>A is a splicing mutation and results in an in frame deletion of the gene, **e23**
- splicing**, Identification of a mutation that perturbs *NF1* gene splicing using genomic DNA samples and a minigene assay, **220**
- splicing defect**, Recurrent mutations in the *NF1* gene are common among neurofibromatosis type 1 patients, **e82**
- splicing mutation**, Transcript analysis of the cystic fibrosis splicing mutation 1525-1G>A shows use of multiple alternative splicing sites and suggests a putative role of exonic splicing enhancers, **e88**
- split hand/foot malformation**, *P63* mutations are not a major cause of non-syndromic split hand/foot malformation, **55**
- spondylocostal dysostosis**, Novel mutations in *DLL3*, a somitogenesis gene encoding a ligand for the Notch signalling pathway, cause a consistent pattern of abnormal vertebral segmentation in spondylocostal dysostosis, **333**
- startle disease**, A novel mutation (R218Q) at the boundary between the N-terminal and the first transmembrane domain of the glycine receptor in a case of sporadic hyperekplexia, **e71**
- steroid 21-hydroxylase deficiency**, Mapping of a de novo unequal crossover causing a deletion of the steroid 21-hydroxylase (*CYP21A2*) gene and a non-functional hybrid tenascin-X (*TNXX*) gene, **e53**
- sterol pathway**, Greenberg dysplasia (HEM) and lethal X linked dominant Conradi-Hünermann chondrodysplasia punctata (CDPX2): presentation of two cases with overlapping phenotype, **e129**
- submicroscopic deletion**, Telomeres: a diagnosis at the end of the chromosomes, **385**
- subtelomeric cryptic rearrangement**, Distal trisomy 6p and 20q owing to the concurrent transposition of distal 6p and 20q to the 22q telomere: a genomic polymorphism?, **e94**
- subtelomeric rearrangements**, A familial unbalanced subtelomeric translocation resulting in monosomy 6q27→qter, **e48**
- sulphate transporter**, Recessive multiple epiphyseal dysplasia (rMED): phenotype delineation in eighteen homozygotes for *DTDST* mutation R279W, **65**
- supernumerary chromosome**, A supernumerary marker chromosome 15 tetrasomic for the Prader-Willi/Angelman syndrome critical region in a patient with a severe phenotype, **e84**
- supernumerary marker chromosome (SMC)**, Partial hexasomy 15pter→15q13 including *SNRPN* and D15S10: first molecular cytogenetically proven case report, **e28**
- susceptibility**, Does apolipoprotein E polymorphism influence susceptibility to malaria?, **348**
- susceptibility allele**, *CHEK2* 1100delC and colorectal cancer, **e110**
- susceptibility genes**, Hereditary pheochromocytomas and paragangliomas: a study of five susceptibility genes, **e75**
- syndrome**, Molecular characterisation of the 22q13 deletion syndrome supports the role of haploinsufficiency of *SHANK3/PROSAP2* in the major neurological symptoms, **575**
- syndromic Hirschsprung disease**, Characterisation of deletions of the *ZFX1B* region and genotype-phenotype analysis in Mowat-Wilson syndrome, **601**
- synonymous codon change**, A synonymous codon change in the *LMNA* gene alters mRNA splicing and causes limb girdle muscular dystrophy type 1B, **e115**
- Tay-Sachs disease**, A genetic screening programme for Tay-Sachs disease and cystic fibrosis for Australian Jewish high school students, **e45**
- TBX5**, Novel *TBX5* mutations and molecular mechanism for Holt-Oram syndrome, **e29**
- TDI**, The metabotropic glutamate receptor 8 gene at 7q31: partial duplication and possible association with autism, **e42**
- TECTA**, Distinctive audiometric profile associated with DFNB21 alleles of *TECTA*, **360**
- tectorial membrane**, Distinctive audiometric profile associated with DFNB21 alleles of *TECTA*, **360**
- telomere**, Telomeric 22q13 deletions resulting from rings, simple deletions, and translocations: cytogenetic, molecular, and clinical analyses of 32 new observations, **690**
- telomeres**, Telomeres: a diagnosis at the end of the chromosomes, **385**
- telomeric rearrangement**, Cryptic terminal deletion of chromosome 9q34: a novel cause of syndromic obesity in childhood?, **300**
- testicular cancer**, Y chromosome haplotypes and testicular cancer in the English population, **e20**
- tetrasomy**, A supernumerary marker chromosome 15 tetrasomic for the Prader-Willi/Angelman syndrome critical region in a patient with a severe phenotype, **e84**
- TFAP2A**, Genetic variation analysis of *MLP*, *TFAP2A*, and *CSK* in patients with neural tube defects, **e43**
- TGFB3**, Linkage and linkage disequilibrium searched for between non-syndromic cleft palate and four candidate loci, **464**
- TGF β** , Molecular and functional analysis identifies ALK-1 as the predominant cause of pulmonary hypertension related to hereditary haemorrhagic telangiectasia, **865**

- TGF- β 1**, APOE and TGF- β 1 genes are associated with obesity phenotypes, **918**
- TGF- β pathway**, Germline mutations in the TGF- β and Wnt signalling pathways are a rare cause of the "multiple" adenoma phenotype, **e35**
- TGF- β receptors**, Visceral manifestations in hereditary haemorrhagic telangiectasia type 2, **494**
- thalidomide**, Mutations at the *SALL4* locus on chromosome 20 result in a range of clinically overlapping phenotypes, including Okihiro syndrome, Holt-Oram syndrome, acro-renal-ocular syndrome, and patients previously reported to represent thalidomide embryopathy, **473**
- TMPRSS3**, Pathogenic mutations but not polymorphisms in congenital and childhood onset autosomal recessive deafness disrupt the proteolytic activity of TMPRSS3, **629**
- TNF**, Effect of *TNF* and *LTA* polymorphisms on biological markers of response to oxidative stimuli in coal miners: a model of gene-environment interaction, **96**
- TNF- α** , Inflammation, genetics, and longevity: further studies on the protective effects in men of *IL-10* -1082 promoter SNP and its interaction with *TNF- α* -308 promoter SNP, **296**
- TNXB**, Mapping of a de novo unequal crossover causing a deletion of the steroid 21-hydroxylase (*CYP21A2*) gene and a non-functional hybrid tenascin-X (*TNXB*) gene, **e53**
- tooth agenesis**, A unique form of hypodontia seen in Vietnamese patients: clinical and molecular analysis, **e79**
- tortuosity**, Homozygosity mapping of a gene for arterial tortuosity syndrome to chromosome 20q13, **747**
- Townes-Brocks syndrome**, High incidence of the R276X *SALL1* mutation in sporadic but not familial Townes-Brocks syndrome and report of the first familial case, **e127**
- TP63**, Rapp-Hodgkin and AEC syndromes due to a new frameshift mutation in the TP63 gene, **e133**
- trafficking defect**, *TBX5* mutations and molecular mechanism for Holt-Oram syndrome, **e29**
- transcription factor**, Disruption of the neuronal PAS3 gene in a family affected with schizophrenia, **325**
- translocation**, Identification of a 650 kb duplication at the X chromosome breakpoint in a patient with 46,X,t(X;8)(q28;q12) and non-syndromic mental retardation, **169**
Patient with bilateral periventricular nodular heterotopia and polymicrogyria with apparently balanced reciprocal translocation t(1;6)(p12;p12.2) that interrupts the mannosidase alpha, class 1A, and glutathione S-transferase A2 genes, **e128**
The neurobeachin gene is disrupted by a translocation in a patient with idiopathic autism, **352**
- transthyretin**, Genetic study of transthyretin amyloid neuropathies: carrier risks among French and Portuguese families, **e120**
- triglycerides**, Ser19→Trp polymorphism within the apolipoprotein AV gene in hypertriglyceridaemic people, **e105**
- trisomy 10**, Partial trisomy 10q with mild phenotype caused by an unbalanced X;10 translocation, **e61**
- trisomy mosaicism**, Clinical aspects, prenatal diagnosis, and pathogenesis of trisomy 16 mosaicism, **175**
- trypsin**, *SPINK1* mutations and phenotypic expression in patients with pancreatitis associated with trypsinogen mutations, **e40**
- TSC2**, Association of INPP1, PIK3CG, and TSC2 gene variants with autistic disorder: implications for phosphatidylinositol signalling in autism, **e119**
- TSC2-PKD1 contiguous gene syndrome**, Large deletion causing the *TSC2-PKD1* contiguous gene syndrome without infantile polycystic disease, **e17**
- tuberous sclerosis**, Association of INPP1, PIK3CG, and TSC2 gene variants with autistic disorder: implications for phosphatidylinositol signalling in autism, **e119**
- tuberous sclerosis complex**, Large deletion causing the *TSC2-PKD1* contiguous gene syndrome without infantile polycystic disease, **e17**
- tumour**, Single nucleotide polymorphism (SNP) analysis of mouse pulmonary adenoma susceptibility loci 1–4 for identification of candidate genes, **e36**
- tumour suppressor**, *HRPT2* mutations are associated with malignancy in sporadic parathyroid tumours, **657**
- tumour suppressor gene**, Chromosome 2 (2p16) abnormalities in Carney complex tumours, **268**
- Turkish**, High frequency of T9 and *CFTR* mutations in children with idiopathic bronchiectasis, **530**
- two dimensional scanning**, Rapid, inexpensive scanning for all possible *BRCA1* and *BRCA2* gene sequence variants in a single assay: implications for genetic testing, **e33**
- uncultured amniocytes**, Rapid, high throughput prenatal detection of aneuploidy using a novel quantitative method (MLPA), **907**
- uniparental disomy**, Comprehensive microsatellite marker analysis contradicts previous report of segmental maternal heterodisomy of chromosome 14, **e26**
Death in adults with Prader-Willi syndrome may be correlated with maternal uniparental disomy, **e63**
Exclusion of maternal uniparental disomy of chromosome 14 in patients referred for Prader-Willi syndrome using a multiplex methylation polymerase chain reaction assay, **e46**
Paternal uniparental disomy in monozygotic twins discordant for hemihypertrophy, **223**
- uniparental disomy 13**, Uniparental disomy of chromosome 13q causing homozygosity for the 35delG mutation in the gene encoding connexin26 (*GJB2*) results in prelingual hearing impairment in two unrelated Spanish patients, **636**
- UPD**, Prader-Willi syndrome: the psychopathological phenotype in uniparental disomy, **e112**
- UPD(14)**, Comprehensive microsatellite marker analysis contradicts previous report of segmental maternal heterodisomy of chromosome 14, **e26**
- USH2A**, Mutations in *USH2A* in Spanish patients with autosomal recessive retinitis pigmentosa: high prevalence and phenotypic variation, **e8**
- Usher syndrome**, Clinical and immunohistochemical evidence for an X linked retinitis pigmentosa syndrome with recurrent infections and hearing loss in association with an *RPGR* mutation, **e118**
Genetic homogeneity and phenotypic variability among Ashkenazi Jews with Usher syndrome type III, **767**
- usherin**, Mutations in *USH2A* in Spanish patients with autosomal recessive retinitis pigmentosa: high prevalence and phenotypic variation, **e8**
- validation**, Evaluation of breast cancer risk assessment packages in the family history evaluation and screening programme, **807**
- valves**, Canine tricuspid valve malformation, a model of human Ebstein anomaly, maps to dog chromosome 9, **320**
- variable outcome**, Clinical, radiological, and chondro-osseous findings in opsismodysplasia: survey of a series of 12 unreported cases, **195**
- vascular disorder**, Visceral manifestations in hereditary haemorrhagic telangiectasia type 2, **494**
- VEGFR-3**, Identification of eight novel *VEGFR-3* mutations in families with primary congenital lymphoedema, **697**
- vestibular schwannoma**, Neurofibromatosis 2 in a patient with a de novo balanced reciprocal translocation 46,X,t(X;22)(p11.2;q11.2), **682**
Somatic mosaicism in neurofibromatosis 2: prevalence and risk of disease transmission to offspring, **459**
- VHL gene**, High frequency of novel germline mutations in the *VHL* gene in the heterogeneous population of Brazil, **e31**
- visceral manifestations**, Visceral manifestations in hereditary haemorrhagic telangiectasia type 2, **494**
- visuospatial and visuoconstructive abilities**, Unusual cognitive and behavioural profile in a Williams syndrome patient with atypical 7q11.23 deletion, **526**
- Walker-Warburg syndrome**, A homozygous nonsense mutation in the *Fukutin* gene causes a Walker-Warburg syndrome phenotype, **845**
- Weill-Marchesani syndrome**, In frame fibrillin-1 gene deletion in autosomal dominant Weill-Marchesani syndrome, **34**
- Williams syndrome**, Unusual cognitive and behavioural profile in a Williams syndrome patient with atypical 7q11.23 deletion, **526**
- Williams-Beuren syndrome**, Partial deletion of the critical 1.5 Mb interval in Williams-Beuren syndrome, **e99**
Using case study comparisons to explore genotype-phenotype correlations in Williams-Beuren syndrome, **136**
- willingness to pay**, Genetic information but not termination: pregnant women's attitudes and willingness to pay for carrier screening for deafness genes, **e80**
- Wnt pathway**, Germline mutations in the TGF- β and Wnt signalling pathways are a rare cause of the "multiple" adenoma phenotype, **e35**

- X**, Neurofibromatosis 2 in a patient with a *de novo* balanced reciprocal translocation 46,X,t(X;22)(p11.2;q11.2), **682**
- X; autosome translocation**, Partial trisomy 10q with mild phenotype caused by an unbalanced X;10 translocation, **e61**
- X chromosome**, Confirmation linkage study in support of the X chromosome harbouring a QTL underlying human height variation, **825**
Identification of a 650 kb duplication at the X chromosome breakpoint in a patient with 46,X,t(X;8)(q28;q12) and non-syndromic mental retardation, **169**
- X inactivation**, Fabry disease: molecular studies in Italian patients and X inactivation analysis in manifesting carriers, **e103**
Neurofibromatosis 2 in a patient with a *de novo* balanced reciprocal translocation 46,X,t(X;22)(p11.2;q11.2), **682**
Partial trisomy 10q with mild phenotype caused by an unbalanced X;10 translocation, **e61**
- X linked cone-rod dystrophy**, A new genetic locus for X linked progressive cone-rod dystrophy, **418**
- X linked mental retardation**, Rare polymorphic variants of the *AGTR2* gene in boys with non-specific mental retardation, **357**
- X linked retinitis pigmentosa**, *RPGR* mutation associated with retinitis pigmentosa, impaired hearing, and sinorespiratory infections, **609**
- Y chromosome**, A Y chromosomal influence on prostate cancer risk: the multi-ethnic cohort study, **815**
The human Y chromosome's azoospermia factor b (AZFb) region: sequence, structure, and deletion analysis in infertile men, **18**
- Y chromosome haplotypes**, Y chromosome haplotypes and testicular cancer in the English population, **e20**
- ZFX1B**, Characterisation of deletions of the *ZFX1B* region and genotype-phenotype analysis in Mowat-Wilson syndrome, **601**