ORIGINAL ARTICLE

Non-random asynchronous replication at 22q11.2 favours unequal meiotic crossovers leading to the human 22q11.2 deletion

A Baumer*, M Riegel*, A Schinzel

J Med Genet 2004;41:413–420. doi: 10.1136/jmg.2003.016352

See end of article for authors' affiliations

Correspondence to:
Dr A Baumer, Institute of
Medical Genetics,
University of Zurich,
Schorenstrasse 16,
CH-8603 Schwerzenbach,
Switzerland; baumer@
medgen.unizh.ch

Revised version received
16 January 2004
Accepted for publication
18 February 2004

Background: Analyses of the replication timing at 22q11.2 were prompted by our finding of a statistically significant bias in the origin of the regions flanking the deletion site in patients with 22q11.2 deletions, the proximal region being in the majority of cases of grandmaternal origin. We hypothesised that asynchronous replication may be involved in the formation of the 22q11.2 deletion, the most frequently occurring interstitial deletion in humans, by favouring the mispairing of low-copy repeats.

Methods: Replication timing during S phase at 22q11.2 was investigated by fluorescent *in situ* hybridisation on interphase nuclei. We report on the detection of non-random asynchronous replication at the human chromosome region 22q11.2, an autosomal locus believed not to contain imprinted genes.

Results: Asynchronous replication at 22q11.2 was observed without exception in all 20 tested individuals; these comprised individuals with structurally normal chromosomes 22 (10 cases), individuals with translocations involving the locus 22q11.2 (eight cases), and patients with a 22q11.2 deletion (two cases). The non-random nature of the asynchronous replication was observed in all individuals for whom the chromosomes 22 were distinguishable. The earlier replicating allele was found to be of paternal origin in all cases where the parental origin of the translocation or deletion was known.

Replication of the genome is a highly regulated process. Whilst the vast majority of chromosome loci replicate synchronously, asynchronous replication in normal cells is a phenomenon known to be associated with allelic exclusion. Indeed, asynchronous replication has been reported for the X chromosomes in females,¹ at autosomal regions containing imprinted genes,^{2–4} at olfactory receptor genes,⁵ and the murine B cell receptor loci.⁶ In contrast to the random allelic exclusion of the two X chromosomes and the non-imprinted autosomal loci, each parental allele at loci containing imprinted genes is specifically either early or later replicating. We report on the detection of non-random asynchronous replication at the human chromosome region 22q11.2. The locus is believed not to contain imprinted genes based on the normal phenotypes associated with uniparental disomy 22⁷ and the lack of parent-of-origin specific phenotypes in patients with hemizygous 22q11.2 deletions. Allelic exclusion at B cell receptor gene loci in mice was recently reported.⁶ Interestingly, a number of genes involved in the immune response are contained in the human 22q11.2 region; however, the findings of Mostoslavsky *et al*⁸ show a random pattern of allelic exclusion.

Based on the results presented here we suggest that the asynchronous replication at 22q11.2 is associated with the formation of the human 22q11.2 deletion. The 22q11.2 deletions appear to arise by mispairing of highly homologous regions, low-copy repeats (LCR), flanking a region of about 3 Mb.⁸ The proximal breakpoint is common to the vast majority of 22q11.2 deletions (approximately 97%), whereas at least three different distal breakpoints have been described, one being by far the most common.⁸ A statistically significant number of 22q11.2 deletions appear to arise by unequal meiotic crossovers.⁹

A deletion of 1.5 Mb at 7q11.23 represents a further frequently observed microdeletion in humans, associated with Williams-Beuren syndrome, which we have previously shown to be also associated with unequal meiotic recombination events.^{9–10} Furthermore, chromosome 7 contains

imprinted genes at 7p11.2–p12 and at 7q32. A recent study showed that a polymorphism at 7q11.23 consisting of an inversion was detected in 33% of parents who transmitted the abnormal chromosome.¹¹ Similarly, an inversion polymorphism was detected in a significant number of mothers of Angelman syndrome patients with a maternal 15q11–q13 deletion.¹² However, an inversion of the critical region was not detected in the parents of 18 patients with the 22q11.2 deletion.¹³ The inversions at 7q11.23 and 15q11–q13 could possibly represent a risk factor for the occurrence of a deletion by reducing the recombination frequency. Indeed, mispairing of repeat elements appears to be favoured by reduced meiotic recombination frequencies. Evidence has been shown of reduced recombination frequencies at 17p12 associated with the CMT1A duplication/HNPP deletion¹⁴ and at 17p11.2 associated with the Smith-Magenis syndrome deletion.¹⁵

METHODS

Microsatellite analysis

We included in the study all patients with a *de novo* hemizygous 22q11.2 deletion for whom parental and grandparental blood samples were available (20 cases, table 1). Patients 1–8, 18, and 19 were already included in a previous study.⁹ The first case refers to two affected monozygous twins, who are considered as a single proband for the purposes of this study. DNA was extracted using standard procedures from peripheral blood lymphocytes of the patients, their parents, and after having established the parental origin of the deletion also from the relevant grandparents. Informed consent was obtained from the adults and the parents of the underage patients. The origin of the deletion was analysed by microsatellite analysis based on the alleles of the patients and their parents. The following markers were used: *D22S1638*, *D22S941*, *D22S944*, and

Abbreviation: LCR, low-copy repeats

D22S264. The grandparental samples were included in the investigations of the haplotypes of the regions flanking the breakpoints. Microsatellites at the proximal region included: *F8VWFP*, *D22S427*, and *D22S420*; microsatellites at the distal breakpoint region included: *D22S303*, *D22S343*, *D22S306*, and *D22S539*. Microsatellite analysis was performed as described in a previous publication.⁹

Fluorescent in situ analysis

Peripheral blood samples drawn from 20 individuals were analysed after obtaining informed consent: 10 healthy control individuals (cases 21–30), eight unaffected individuals with balanced translocations (*t*(6;22), *t*(11;22) and *t*(19;22)) involving the 22q11.2 region (cases 31–38), and two patients with the common 22q11.2 deletion (cases 13 and 17).

Analysis of the replication pattern by BrdU incorporation was performed following standard procedures.¹⁶ Probes were used mapping within the deleted region (LSI DiGeorge/VCFs Region Probe (TUPLE 1; Vysis), *ZNF74*, and “*D22S936*”) and at flanking positions (“*D22S427*” and *ARSA* (Vysis)). The probes “*D22S427*”, *ZNF74* and “*D22S936*” each consist of pools of five different PCR products of about 400–500 bp, amplified from a region covering about 10 kb of genomic sequence. The probes are denoted based on known markers contained in the respective regions.^{8,17} Oligonucleotide sequences and PCR conditions are provided in table 2. Different combinations of probes were used (tables 3 and 4), which were scored individually and in pairs. The location of the signals in the nuclei^{18,19} as well as the distance between the signals allowed differentiation between the two chromosomes 22 (examples are shown in fig 1).

Probes used for assaying the replication timing at the control regions 7q11.23, 5p15.2, 15q12, and Xp22.3 were LSI Elastin/D7S486 (elastin gene), LSI D5S23, LSI SNRPN/D15Z1/PML Probe (*SNRPN*), and LSI Steroid Sulfatase (*STS*), respectively (all probes were purchased from Vysis).

RESULTS

Bias in the origin of the regions flanking the 22q11.2 deletions

We initially postulated that the human autosomal region 22q11.2 replicates asynchronously based on the results we

obtained by studying the mode and time of formation of 22q11.2 deletions. The study involved determination of the parental origin of the de novo hemizygous 22q11.2 deletions in 20 probands followed by analysis of the haplotype segregation at the regions flanking the deletion breakpoints. This latter analysis revealed the presence of an interchromosomal recombination in 17 cases, whereas an intrachromosomal rearrangement was detected only in three cases (table 1). The average distance between the informative markers closest to the deletion breakpoint was approximately 8.8 cM. Thus, the chances of observing 17 or more meiotic crossovers in this region in a group of 20 control individuals would be $p = 1.0 \times 10^{-15}$. The results obtained in this study based on an increased number of 22q11.2 deletion cases corroborate our previous finding⁹ that a significant number of 22q11.2 deletions are associated with an unequal meiotic recombination. Interestingly, a statistically significant bias in the parental origin of the proximal and distal positions emerged from the data collected: in 15 of the 17 cases where an interchromosomal crossover occurred at the deletion site, the proximal region was derived from the grandmother and only in two cases from the grandfather ($p = 1.1 \times 10^{-3}$, $\chi^2 = 9.95$).

We suggested that the most plausible reason for this bias is an asynchronous replication of the 22q11.2 region.

Proposed model for the formation of the human 22q11.2 deletion

Schematically illustrated in fig 2 is the proposed model for a mispairing event occurring during meiosis in a parental gamete. Assuming that the initial configuration leading to a mispairing event occurs while the replication forks have already passed both parental LCRI and LCRII on the paternal homologue but have not yet passed the LCRII on the maternal allele, then mispairing of the paternal LCRII with the maternal LCRI would be favoured as compared to the mispairing of the maternal LCRII and the paternal LCRI. The mispairing would then be the basis for an unequal crossing-over event later in meiosis, giving rise to a 22q11.2 deletion in the zygote with a predominant grandmaternal origin of the proximal region and grandpaternal origin of the distal region.

Table 1 Summary of the microsatellite results obtained for the patients with the 22q11.2 deletion

Case	Gender	Parental origin of the deletion	Grandparental origin of the proximal region*	Grandparental origin of the distal region*
Cases with a meiotic recombination at the deletion site				
1	Female†	Maternal	Grandpaternal (<i>F8VWFP</i>)	Grandmaternal (<i>D22S306</i>)
2	Male	Paternal	Grandmaternal (<i>D22S420</i>)	Grandpaternal (<i>D22S306</i>)
3	Female	Paternal	Grandmaternal (<i>D22S427</i>)	Grandpaternal (<i>D22S303</i>)
4	Female	Maternal	Grandmaternal (<i>D22S427</i>)	Grandpaternal (<i>D22S303</i>)
5	Male	Paternal	Grandmaternal (<i>D22S420</i>)	Grandpaternal (<i>D22S303</i>)
6	Female	Paternal	Grandmaternal (<i>D22S420</i>)	Grandpaternal (<i>D22S343</i>)
7	Male	Maternal	Grandmaternal (<i>D22S420</i>)	Grandpaternal (<i>D22S306</i>)
8	Female	Paternal	Grandmaternal (<i>D22S427</i>)	Grandpaternal (<i>D22S343</i>)
9	Female	Paternal	Grandmaternal (<i>D22S427</i>)	Grandpaternal (<i>D22S306</i>)
10	Female	Maternal	Grandmaternal (<i>D22S427</i>)	Grandpaternal (<i>D22S306</i>)
11	Male	Maternal	Grandmaternal (<i>D22S420</i>)	Grandpaternal (<i>D22S303</i>)
12	Female	Maternal	Grandpaternal (<i>D22S420</i>)	Grandmaternal (<i>D22S303</i>)
13	Female	Maternal	Grandmaternal (<i>D22S420</i>)	Grandpaternal (<i>D22S539</i>)
14	Male	Maternal	Grandmaternal (<i>D22S427</i>)	Grandpaternal (<i>D22S306</i>)
15	Male	Paternal	Grandmaternal (<i>D22S427</i>)	Grandpaternal (<i>D22S343</i>)
16	Female	Maternal	Grandmaternal (<i>D22S427</i>)	Grandpaternal (<i>D22S303</i>)
17	Female	Paternal	Grandmaternal (<i>D22S427</i>)	Grandpaternal (<i>D22S303</i>)
Cases with an intrachromosomal rearrangement at the deletion site				
18	Male	Maternal	Grandmaternal (<i>D22S427</i>)	Grandmaternal (<i>D22S303</i>)
19	Female	Paternal	Grandpaternal (<i>D22S427</i>)	Grandpaternal (<i>D22S343</i>)
20	Male	Maternal	Grandpaternal (<i>D22S427</i>)	Grandpaternal (<i>D22S306</i>)

*The informative microsatellite markers closest to the deletion breakpoint are shown in brackets; †Monozygous twins.

Figure 1 Representative examples of interphase nuclei analysed by fluorescent in situ hybridisation. Nuclei showing a double signal on one chromosome and a single signal on the other chromosome are illustrated for: (A) the LSI DiGeorge/VCFs Region Probe (case 21); (B) control region 7q11.23, LSI Williams Syndrome probe (case 21); (C–E) LSI DiGeorge/VCFs Region Probe (red signals) and ARSA, 22q13.3 (green signals) for (C) case 31, (D) case 35, and (E) case 38; (F–G) “D22S427” (red signals) and ZNF74 (green signals) for cases 22 and 32, respectively; (H–I) “D22S427” and “D22S936” (green signals) for cases 22 and 35, respectively; (J–L) “D22S427” and ARSA (green signals) for cases 22, 35, and 17, respectively. In panels (C) to (L), the signals representing the structurally normal chromosome 22 are circled as well as the derivative chromosomes.

Asynchronous replication at human 22q11.2

Direct evidence for the non-random asynchronous replication during S phase at 22q11.2 was obtained by fluorescent in situ hybridisation on interphase nuclei. Asynchronous replication is inferred when more than about 25% of nuclei show the presence of two distinct signals (ds) on the homologue that has already gone through replication, while only one signal (ss) is observed on the homologue that has not yet replicated.^{2–5,6} The replication timing at 22q11.2 and 22q13.3 was analysed in peripheral blood cells of 20 individuals, in 10 of whom it was possible to differentiate between the two chromosomes 22 in interphase nuclei, owing either to balanced translocations involving a chromosome 22 (eight individuals) or to a 22q11.2 deletion (two cases).

Analysis of the replication timing in the 10 control individuals with structurally normal chromosomes 22 revealed a high number of nuclei showing a ds/ss pattern (table 3). The mean value obtained for the DiGeorge/VCFs probe was 84.6% with a 95% confidence interval (95% CI) ranging from 82% to 87.2%. The mean value obtained for the ARSA probe was 76.6% with a 95% CI ranging from 73.3 to 79.9. The difference between these values and the generally accepted threshold value of about 25% for asynchronous replication are strongly significant ($p<0.0001$).

Similar high percentages of nuclei with ds/ss were detected in individuals with translocations and deletions (table 4), the only exception being the significantly lower values obtained with 22q probes mapping to the derivative chromosome 11 in the three individuals with a paternal

Table 2 Oligonucleotide sequences used for the generation of FISH probes

Probe	Oligonucleotide pairs		Annealing °C, PCR enhancers if needed and product size
	Forward (5'–3')	Reverse (5'–3')	
"D22S427"	A: ccaagcgttgggattac	A: caccatgtgtcagatgc	56°C, 486 bp
	B: gaccacagagactgaagac	B: gggctcagcctcagacag	56°C+10% DMSO, 477 bp
	C: ccgttcattgattatgtgg	C: cgatctgagctcactgcaac	56°C, 476 bp
	D: gcttgctgcatltaacatt	D: gactcggctctcggtctac	56°C, 448 bp
	E: gagtcgaaggacttgggagt	E: ggagggtgtgacaaaactg	56°C, 494 bp
ZNF74	A: attgggggtagcagctcttt	A: gggcaacatctggactctc	56°C, 479 bp
	B: tagctactgcagccttcaa	B: gtggttaccaaggaggagagg	56°C, 425 bp
	C: gtccacatgccttgaaaa	C: caagagccaaactctgcaaaa	56°C, 499 bp
	D: ggatttgggtccagctcta	D: ccagaacagggtctcagaca	56°C, 453 bp
	E: gtgtgttgccctattgct	E: gttagggaggccagttagg	56°C, 448 bp
"D22S936"	A: caccctcactgcttgagt	A: agattgtgccactcactcc	56°C, 455 bp
	B: ctgcagagccattttgggt	B: aatcaaaagcctgaatgtgg	56°C, 429 bp
	C: gccacccaggataatcttc	C: gggcaacaagagcaaaactc	56°C, 477 bp
	D: caggggttaatttgaagga	D: agctcagctctggctctgac	56°C, 500 bp
	E: taagaaccagcctcggtgt	E: tgggtgtggaagagaaggac	56°C, 480 bp

Table 3 Interphase nuclei of control individuals with structurally normal chromosomes 22 showing a double (ds) and a single signal (ss) for probes within the commonly deleted 22q11.2 region and at flanking positions

Case, gender*	FISH probes†	Percentage of interphase nuclei‡ showing either ds/ss, ds/ds, or ss/ss for individual probes			2×ds on one chr. 22 and 2×ss on the other chr. 22 for combinations of two probes§
		ds/ss	ds/ds	ss/ss	
21, F	"D22S427"	72% (201)	7% (201)	21% (201)	
	LSI DiGeorge/VCFS	88% (115)	2% (115)	10% (115)	
	ZNF74	67% (101)	15% (101)	18% (101)	
	ARSA	74% (201)	2% (201)	24% (201)	
	"D22S427" and ZNF74				52% (101)
22, F	"D22S427" and ARSA				57% (96)
	LSI DiGeorge/VCFS and ARSA				61% (100)
	"D22S427"	67% (303)	10% (303)	23% (303)	
	LSI DiGeorge/VCFS	89% (115)	4% (115)	7% (115)	
	ZNF74	58% (100)	7% (100)	35% (100)	
	"D22S936"	70% (91)	3% (91)	27% (91)	
	ARSA	69% (209)	8% (209)	23% (209)	
	"D22S427" and ZNF74				38% (98)
	"D22S427" and "D22S936"				64% (91)
	"D22S427" and ARSA				46% (106)
23, M	LSI DiGeorge/VCFS and ARSA				49% (100)
	LSI DiGeorge/VCFS	84% (106)	4% (106)	12% (106)	
	ARSA	79% (108)	4% (108)	17% (108)	
24, F	LSI DiGeorge/VCFS and ARSA				60% (101)
	LSI DiGeorge/VCFS	79% (111)	6% (111)	15% (111)	
25, F	ARSA	75% (113)	10% (113)	15% (113)	
	LSI DiGeorge/VCFS and ARSA				53% (105)
26, F	LSI DiGeorge/VCFS	88% (103)	2% (103)	10% (103)	
	ARSA	76% (104)	5% (104)	19% (104)	
27, M	LSI DiGeorge/VCFS and ARSA				57% (103)
	LSI DiGeorge/VCFS	81% (117)	2% (117)	17% (117)	
28, M	ARSA	78% (115)	2% (115)	20% (115)	
	LSI DiGeorge/VCFS and ARSA				50% (113)
29, M	LSI DiGeorge/VCFS	84% (117)	0% (117)	16% (117)	
	ARSA	78% (118)	3% (118)	19% (118)	
30, M	LSI DiGeorge/VCFS and ARSA				49% (113)
	LSI DiGeorge/VCFS	85% (114)	2% (114)	13% (114)	
31, F	ARSA	83% (115)	2% (115)	15% (115)	
	LSI DiGeorge/VCFS and ARSA				58% (112)
32, M	LSI DiGeorge/VCFS	88% (104)	3% (104)	9% (104)	
	ARSA	83% (104)	4% (104)	13% (104)	
33, F	LSI DiGeorge/VCFS and ARSA				53% (99)
	LSI DiGeorge/VCFS	80% (108)	1% (108)	19% (108)	
34, M	ARSA	71% (108)	2% (108)	27% (108)	
	LSI DiGeorge/VCFS and ARSA				51% (107)

*M, male; F, female; †The relative location of the probes is as follows: "D22S427": proximal to the LCRI; LSI DiGeorge/VCFS: within the deleted region, proximal to the breakpoint of the t(11;22); ZNF74: within the commonly deleted region, distal to the breakpoint of the t(11;22); "D22S936": within the commonly deleted region, proximal to the LCRI¹⁷; ARSA: distal to the LCRI, 22q13.3; ‡In brackets: total number of interphase nuclei analysed; §Out of the 10 possible combinations of ds and ss for two probes on the two chromosomes 22, only the combination ds/ds on one chromosome 22 and ss/ss on the other chromosome 22 is indicated being the most frequently detected combination in each case.

origin of the translocation (individuals 34, 35, and 36). The mean percentage of ds/ss obtained with the LSI DiGeorge/VCFS probe for the eight individuals with translocations was 83.9% (95% CI: 76.9 to 90.8). The mean percentages of ds/ss obtained with the ARSA probe in the individuals with a paternal origin of the translocation (individuals 34–36) was 35.7% (95% CI: 17.0 to 54.3) and 82.4% in the remaining five individuals with translocations (95% CI: 72.4 to 92.4).

The values observed in the 10 cases with structurally normal chromosomes 22 were comparable to those obtained in individuals with a translocation and those with a 22q11.2 deletion. Two-sample *t* tests were performed and resulted in non-significant differences between the following groups: (a) results for the LSI DiGeorge/VCFS probe in the 10 control individuals versus the eight individuals with translocations: *p* = 0.81; (b) results for the ARSA probe in the 10 control individuals versus five individuals with translocations (not considering individuals 34–36): *p* = 0.095. On the other hand, the probes mapping to the derivative chromosome 11 in the three individuals with a paternal origin of the translocation showed a strongly significant reduction in the percentage of nuclei with ds/ss (two-sample *t* test with the results for the

ARSA probe in the 10 control individuals 21–30 versus individuals 34–36: *p* < 0.0001).

On the basis of these results it appears that the asynchronous replication at this region is a general phenomenon and is not an effect of either translocations or deletions. Differences in the replication timing were only seen for 22q regions translocated onto another chromosome in three cases. No major changes in the replication timing at 22q11.2 were observed in the other cases with translocations and deletions compared to control individuals. This legitimates the use of such structurally abnormal chromosomes 22 for investigation of the randomness of the asynchronous replication. It should be stressed that our results cannot exclude minor changes in the replication timing due to translocations or deletions, however, the investigation of such minor changes is beyond the scope of this study.

The three further shorter probes we generated (that is, "D22S427" proximal to the LCRI; and ZNF74 and "D22S936", both mapping within the commonly deleted region distal to the breakpoint of the t(11;22)^{8,17}), used on a subset of individuals, resulted in percentages of nuclei with ds/ss comparable to those obtained with the commercially

Table 4 Interphase nuclei of individuals with a structurally abnormal chromosome 22 showing a double (ds) and a single signal (ss) for probes within the commonly deleted 22q11.2 region and at flanking positions

Case, gender*	Chromosomes 22 (denoted A and B)	FISH probes†	Derivative chromosomes on which the 22q probes hybridise (chromosomes "A")	Interphase nuclei showing ds/ss for single probes§ or 2×ds on one chr. 22 and 2×ss on the other chr. 22 for combinations of two probes¶			Interpretation and comments
				ds on A	ds on B	Total‡	
31, F	A: t(6;22)(q26;q11.2) mat; B: structurally normal	LSI DiGeorge/VCFS ARSA LSI DiGeorge/VCFS and ARSA	der 22 der 6	24% 24% 19%	59% 55% 49%	83% (122) 79% (122) 68%	Non-random asynchronous replication: paternal chromosome 22 early replicating
32, F	A: t(11;22)(q23;q11.2); B: structurally normal	"D22S427" LSI DiGeorge/VCFS ZNF74 "D22S936" ARSA "D22S427" and ZNF74 "D22S427" and "D22S936" "D22S427" and ARSA LSI DiGeorge/VCFS and ARSA	der 22 der 22 der 11 der 11 der 11 der 11	4% 26% 6% 0 5% 2% 0 0 0	69% 63% 47% 62% 65% 45% 52% 61% 63%	73% (307) 89% (102) 53% (115) 62% (107) 70% (169) 47% (105) 52% (107) 61% (88) 63% (81)	Non-random asynchronous replication
33, F	A: t(11;22)(q23;q11.2); B: structurally normal	LSI DiGeorge/VCFS ARSA LSI DiGeorge/VCFS and ARSA	der 22 der 11	24% 24% 22%	69% 65% 64%	93% (98) 89% (93) 86% (93)	Non-random asynchronous replication
34, M	A: t(11;22)(q23;q11.2) pat; B: structurally normal	LSI DiGeorge/VCFS ARSA LSI DiGeorge/VCFS and ARSA	der 22 der 11	48% 21% 10% (52%) A: ds-ss; B: ss-ss)	18% 19% 0	66% (152) 40% (102) 10% (102)	Non-random asynchronous replication: paternal chromosome 22 early replicating
35, M	First son of case 34; A: t(11;22)(q23;q11.2) pat; B: structurally normal	"D22S427" LSI DiGeorge/VCFS ZNF74 "D22S936" ARSA "D22S427" and ZNF74 "D22S427" and "D22S936" "D22S427" and ARSA LSI DiGeorge/VCFS and ARSA	der 22 der 22 der 11 der 11 der 11	59% 61% 10% 3% 14% 2% 2% 3% 11% (63%) A: ds-ss; B: ss-ss)	11% 20% 17% 6% 13% 3% 3% 2% 0	70% (316) 81% (123) 27% (109) 9% (104) 27% (210) 5% (106) 5% (104) 5% (103) 11% (107)	Non-random asynchronous replication for the probes proximal to the translocation breakpoint: paternal chromosome 22 early replicating; Segment of chromosome 22 distal to the translocation breakpoint no longer asynchronous replicating
36, M	Second son of case 34; A: t(11;22)(q23;q11.2) pat; B: structurally normal	LSI DiGeorge/VCFS ARSA LSI DiGeorge/VCFS and ARSA	der 22 der 11	50% 10% 10% (55%) A: ds-ss; B: ss-ss)	32% 30% 0	82% (135) 40% (100) 10% (100)	Non-random asynchronous replication: paternal chromosome 22 early replicating
37, F	A: t(19;22)(p13.3;q11.2); B: structurally normal	LSI DiGeorge/VCFS ARSA LSI DiGeorge/VCFS and ARSA	der 22 der 19	26% 29% 25%	62% 56% 50%	88% (100) 85% (100) 75% (100)	Non-random asynchronous replication
38, F	Daughter of case 37; A: t(19;22)(p13.3;q11.2); B: structurally normal	LSI DiGeorge/VCFS ARSA LSI DiGeorge/VCFS and ARSA	der 22 der 19	29% 28% 24%	60% 61% 53%	89% (100) 89% (100) 77% (100)	Non-random asynchronous replication: paternal chromosome 22 early replicating
13, F	A: maternal 22q11.2 deletion; B: structurally normal	"D22S427" ARSA "D22S427" and ARSA"	der 22 der 22	3% 6% 2%	66% 62% 51%	69% (117) 68% (117) 53% (117)	Non-random asynchronous replication: paternal chromosome 22 early replicating
17, F	A: paternal 22q11.2 deletion; B: structurally normal	"D22S427" ARSA "D22S427" and ARSA"	der 22 der 22	65% 60% 50%	2% 8% 1%	67% (78) 68% (78) 51% (78)	Non-random asynchronous replication: paternal chromosome 22 early replicating

*M, male; F, female; †The relative location of the probes is as follows: "D22S427": proximal to the LCRI; LSI DiGeorge/VCFS: within the deleted region, proximal to the breakpoint of the t(11;22); ZNF74: within the commonly deleted region, distal to the breakpoint of the t(11;22); "D22S936": within the commonly deleted region, proximal to the LCRII (8,17); ARSA: distal to the LCRII, 22q13.3; ‡In brackets: total number of interphase nuclei analysed; §The cells with ds/ss, ds/ds and ss/ss were scored, but only the percentage with ds/ss is indicated, being the value of relevance in this context; ¶Out of the 16 possible combinations of ds and ss for two probes on the two chromosomes 22, the combination ds/ds on one chromosome 22 and ss/ss on the other chromosome 22 is indicated being the most frequently detected combination in most cases (additional data are indicated in cases 34–36 where another combination was more frequent).

available probes (tables 3 and 4). Illustrative examples of interphase nuclei showing double/single signals are shown in fig 1.

The use of combinations of two different probes revealed that the double signals for both probes were most frequently observed on the same chromosome (tables 3 and 4). Thus, the entire region ranging from at least 22q11.2 (proximal to the LCRI) to 22q13.3 appears to replicate asynchronously.

Four different control regions showed the expected proportion of ds/ss in interphase nuclei in two control individuals with structurally normal chromosomes. The control regions where synchronous replication is expected were the *ELN* locus on chromosome 7q11.23 (ds/ss in 19% and 21% of interphase nuclei in cases 21 and 22, respectively, example in fig 1B), and a probe at 5p15.2 spanning *D5S23-D5S721* (ds/ss: 13% and 18%, respectively, in cases 21 and 22, data not shown). Control regions where asynchronous replication is expected were the LSI Prader-Willi/Angelman Region Probe (*SNRPN*) at 15q12 (ds/ss: 42% and 39%, respectively, in individuals 21 and 22, data not shown), and the LSI Steroid Sulfatase (*STS*) probe at Xp22.3 (ds/ss: 36% and 38%, respectively, in individuals 21 and 22, data not shown). The number of interphase nuclei investigated in each case ranged from 100 to 115.

Non-random asynchronous replication: early replication of the paternal 22q11.2 allele

According to our proposed model (fig 2) we would expect the asynchronous replication at 22q11.2 to be non-random. The total percentages of nuclei with ds/ss are indicated in table 4 for the 10 individuals with a structurally abnormal chromosome 22 (cases 31–38, 13, and 17). In addition, the proportion is given as to which chromosome 22 carries the double signal (that is, the derivative chromosomes or the structurally normal 22). In the case of a random asynchronous replication double signals would be evenly distributed between the two chromosomes 22. This was not the case for any of the 10 individuals listed in table 4. The LSI DiGeorge/

VCFS probe showed in all eight cases with translocations a preferential occurrence of the ds on either the structurally normal chromosome 22 (five cases) or the derivative chromosomes 22 (the three cases with a paternal origin of the translocation). The bias in the distribution of the ds in favour of a specific chromosome 22 was statistically significant: (a) eight cases (31–38), LSI DiGeorge/VCFS probe: the mean value of the higher percentages was 59% (95% CI: 53.2 to 64.8) and for the lower percentages 24.9% (95% CI: 21.1 to 28.7), paired-sample *t* test: $p < 0.0001$; (b) five cases (31–33, 37, and 38), *ARSA* probe: the mean value of the higher percentages was 60.4% (95% CI: 54.5 to 66.3) and for the lower percentages 22.0% (95% CI: 9.9 to 34.1), paired-sample *t* test: $p = 0.003$. A non-random pattern of asynchronous replication was also observed in the two patients with a 22q11.2 deletion, individuals 13 and 17, with an early replication of the structurally normal chromosome 22 in the former case and of the chromosome bearing the deletion in the latter case.

Importantly, the paternal allele was earlier replicating in all cases where the parental origin of the chromosomes 22 was known (five cases with a translocation and two cases with a 22q11.2 deletion; table 4).

DISCUSSION

We detected comparable levels of asynchronous replication at 22q11.2 in all tested individuals, that is, control individuals and carriers of either translocations or deletions involving the 22q11.2 region. Based on cases with distinguishable chromosomes 22 we show a non-random nature of the asynchronous replication. In all cases where the origin of the structurally abnormal chromosome 22 was known we detected an earlier replication of the paternal alleles. We hypothesise that a non-random asynchronous replication in this region represents a risk factor for the formation of the 22q11.2 deletion by increasing the probability of an initial mispairing of the parental alleles at the highly homologous low-copy repeats. These initial abnormal conformations may lead, later in meiosis, to an unequal meiotic crossover and thus to the 22q11.2 deletion.

The replication timing results presented here were performed on peripheral blood cells. Our hypothesis would imply that the non-random asynchronous replication also occurs in gametogenesis. In mice it has been shown that the non-random asynchronous replication at imprinted genes is established in gametogenesis and is maintained throughout development in a wide range of tissues.²⁰ The random asynchronous replication observed for example at the murine olfactory receptor loci and B cells occurs before meiosis in gametes, is erased after fertilisation, and is then reset before implantation of the embryo.⁶ Our results represent the first example of non-random asynchronous replication apparently not associated with imprinted genes. It would be interesting to investigate whether this phenomenon also occurs in mice or other model organisms in order to investigate in detail the setting of the non-random asynchronous replication at such loci.

A growing number of researchers have recently reported on their results regarding replication timing using cell cycle fractionation.^{21–22} The replication timing results obtained using different techniques are generally in agreement.^{21–22} Exceptions have been reported at heterochromatic regions and at timing-switch boundaries. At such regions there appears to be a delayed resolution of the sister chromatids, which would appear in FISH analysis as a singlet and would be misleading as to the actual replication timing.^{4–21} A delayed chromatid resolution could also be due amongst other factors to the pausing of the replication fork or to long replicons that are still undergoing replication. Indeed, the FISH based

Figure 2 Schematic representation of mispairing in a situation with asynchronous replication. (A) The genomic segment encompassing the 22q11.2 deletion is schematically illustrated. The proximal block of low-copy repeats (LCRI) and the distal repeat (LCRII) are the positions at which mispairing occurs. For the sake of clarity only one of the distal breakpoints is depicted. (B) In a situation where the two parental alleles (Mat: maternal, Pat: paternal) replicate synchronously, or in the case of random asynchronous replication, the repeats LCRI and LCRII from both chromosomes are free to mispair with one another. No bias in the origin of the proximal and distal regions would be expected. (C) In a situation where the paternal chromosome replicates earlier, mispairing of the paternal LCRII with the maternal LCRI would be possible at a time when the maternal LCRII has not yet gone through replication. A bias in favour of a grandmaternal origin of the proximal region flanking the 22q11.2 deletion would be expected in the patients.

method is not a measurement of replication itself but rather of the sister chromatid resolution. However, FISH analysis remains appropriate for analyses of the replication timing when these are aimed at detecting asynchrony and particularly a possible non-randomness of the phenomenon. In this study we obtained an extremely high level of nuclei showing ds/ss using probes for 22q11.2–q13.3, whilst eight measurements at control regions resulted in ratios of ds/ss in agreement with published data.^{2–5,6} It is possible that, in a subset of nuclei, replication at 22q had actually already occurred on both homologues but one chromosome showed a delayed chromatid resolution. This is in itself intriguing as the parental alleles clearly behave differently in the S phase, thus excluding that the high level of ds/ss is simply due to artefacts such as properties of probes or domain specific resolution.

The sizes of the mammalian replication units have been estimated to range from about 50 kb up to 1 Mb.²³ Replication at each fork is estimated to proceed at about 2 kb per minute in mammalian cells.²⁴ The replication timing appears to be organised in large-scale replication domains of about 1 Mb and sequence elements controlling the initiation of replication may be located at many kb distant from the primary origin.^{25,26} The long arm of chromosome 22 appears to have large segments of several Mb which replicate early in S phase.²² Thus, the replicons covering the 22q11.2 commonly deleted region appear to be part of a large-scale replication domain. We attempted to extrapolate the relative position of the relevant time-switching element(s) from the data obtained for the patients with a 22q11.2 deletion and the patients with a t(11;22). Asynchronous replication was maintained in both patients with a 22q11.2 deletion, indicating that the relevant time-switching element(s) appears to lie outside the deleted region. Analysis of cases with a translocation (11;22) showed that early replication of the structurally normal chromosome 22 was observed for all tested 22q probes in individual 32. On the other hand, early replication of the paternal alleles of individuals 34–36 was restricted to the probes mapping to the derivative 22 chromosome (that is, proximal to the translocation breakpoint: “D22S427” and LSI DiGeorge/VCFS probe).¹⁷ The chromosome 22 segments distal to the breakpoint, which were translocated onto chromosome 11, showed a strongly significant decrease in ds/ss signals (table 4). Therefore, we suggest that the relevant time-switching element(s) responsible for the non-random asynchronous replication at 22q11.2 lies proximal to the LCRI.

Several recent publications report on the intimate coupling of DNA replication, genome stability, and meiotic recombinations in yeast.^{27–31} Although a tight coupling of replication and meiotic recombination has not yet been shown in mammals, certain key proteins in yeast, such as Rec8p, are conserved in a wide range of organisms including humans.³²

One of the control regions investigated in this study was the *ELN* locus on chromosome 7q11.23, which is of particular interest for several reasons. As mentioned above, the 1.5 Mb deletion at human 7q11.23 represents a further frequently observed interstitial deletion, which is also known to be associated with unequal meiotic recombination events.^{9–10} We performed haplotype analysis for 36 patients with de novo 7q11.23 deletions. A meiotic recombination was detected in 23 out of 36 cases. Unlike the situation observed for the 22q11.2 deletions, the flanking regions at 7q11.23 originated almost equally from the two grandparents (the proximal region was derived from the grandmother in 13 cases and from the grandfather in 10 cases, data not shown). Clearly, haplotype analysis at the breakpoints would only show an uneven distribution of the origin of the proximal and distal regions in cases with a non-random allelic exclusion, but not

in the case of a random asynchronous replication. Although the percentage of cells with ds/ss we obtained for the control individuals was in the range indicative of synchronous replication, it would be very interesting to investigate the replication timing in individuals with the inversion polymorphism.

With an incidence of about 1 in 4000 newborns,³³ the 22q11.2 deletion is the most frequently occurring microdeletion in humans. Whether or not asynchronous replication, random or non-random, also favours the mispairing of other low-copy repeats remains to be investigated.

ACKNOWLEDGEMENTS

We wish to thank the patients, their parents, and all control individuals for their cooperation.

We are grateful to Professor T Gasser and Dr V Rousson, Department of Biostatistics, University of Zurich, for their assistance with the statistical evaluations.

Authors' affiliations

A Baumer, M Riegel, A Schinzel, Institute of Medical Genetics, University of Zurich, Schwerzenbach, Switzerland

This study was supported in part by the Swiss National Foundation (grant numbers: 3100-056956.99 and 32-56051.98).

Conflict of interest: none declared.

*The authors AB and MR contributed equally to this work.

REFERENCES

- Schmidt M, Migeon BR. Asynchronous replication of homologous loci on human active and inactive X chromosomes. *Proc Natl Acad Sci U S A* 1990;**87**(10):3685–9.
- Kisberg D, Selig S, Brandeis M, Simon I, Keshet I, Driscoll DJ, Nicholls RD, Cedar H. Allele-specific replication timing of imprinted gene regions. *Nature* 1993;**364**(6436):459–63.
- Elstradiadis A. A new whiff of monoallelic expression. *Curr Biol* 1995;**5**:21–4.
- Selig S, Okumura K, Ward DC, Cedar H. Delineation of DNA replication time zones by fluorescence in situ hybridization. *EMBO J* 1992;**11**(3):1217–25.
- Chess A, Simon I, Cedar H, Axel R. Allelic inactivation regulates olfactory receptor gene expression. *Cell* 1994;**78**(5):823–34.
- Mostoslavsky R, Singh N, Tenzen T, Goldmit M, Gabay C, Elizur S, Qi P, Reubinoff BE, Chess A, Cedar H, Bergman Y. Asynchronous replication and allelic exclusion in the immune system. *Nature* 2001;**414**(6860):221–5.
- Schinzel AA, Basaran S, Bernasconi F, Karaman B, Yuksel-Apak M, Robinson WP. Maternal uniparental disomy 22 has no impact on the phenotype. *Am J Hum Genet* 1994;**54**(1):21–4.
- Shaikh TH, Kurahashi H, Saita SC, O'Hare AM, Hu P, Roe BA, Driscoll DA, McDonald-McGinn DM, Zackai EH, Budarf ML, Emanuel BS. Chromosome 22-specific low copy repeats and the 22q11.2 deletion syndrome: genomic organization and deletion endpoint analysis. *Hum Mol Genet* 2000;**9**(4):489–501.
- Baumer A, Dutly F, Balmer D, Riegel M, Tükel T, Krajewska-Walasek M, Schinzel AA. High level of unequal meiotic crossovers at the origin of the 22q11.2 and 7q11.23 deletions. *Hum Mol Genet* 1998;**7**(5):887–94.
- Dutly F, Schinzel A. Unequal interchromosomal rearrangements may result in elastin gene deletions causing the Williams-Beuren syndrome. *Hum Mol Genet* 1996;**5**(12):1893–8.
- Osborne LR, Li M, Pober B, Chitayat D, Bodurtha J, Mandel A, Costa T, Grebe T, Cox S, Tsui LC, Scherer SW. A 1.5 million-base pair inversion polymorphism in families with Williams-Beuren syndrome. *Nat Genet* 2001;**29**(3):321–5.
- Gimelli G, Pujana MA, Patricelli MG, Russo S, Giordano D, Larizza L, Cheung J, Armengol L, Schinzel A, Estivill X, Zuffardi O. Genomic inversions of human chromosome 15q11–q13 in mothers of Angelman syndrome patients with class II (BP2/3) deletions. *Hum Mol Genet* 2003;**12**(8):849–58.
- Gebhardt GS, Devriendt K, Thoelen R, Swillen A, Pijls E, Gewillig M, Frys JP, Vermeesch JR. No evidence for a parental inversion polymorphism predisposing to rearrangements at 22q11.2 in the DiGeorge/velocardiofacial syndrome. *Eur J Hum Genet* 2003;**11**(2):109–111.
- Inoue K, Dewar K, Katsanis N, Reiter LT, Lander ES, Devon KL, Wyman DW, Lupski JR. The 1.4-Mb CMT1A duplication/HNPP deletion genomic region reveals unique genome architectural features and provides insights into the recent evolution of new genes. *Genome Res* 2001;**11**(6):1018–33.
- Bi W, Yan J, Stankiewicz P, Park SS, Walz K, Boerkoel CF, Potocki L, Shaffer LG, Devriendt K, Nowaczyk MJ, Inoue K, Lupski JR. Genes in a refined Smith-Magenis syndrome critical deletion interval on chromosome 17p11.2 and the syntenic region of the mouse. *Genome Res* 2002;**12**(5):713–28.
- Wegner RD. *Diagnostic cytogenetics*. Berlin: Springer, 1999.
- Kurahashi H, Shaikh TH, Hu P, Roe BA, Emanuel BS, Budarf ML. Regions of genomic instability on 22q11 and 11q23 as the etiology for the recurrent constitutional t(11;22). *Hum Mol Genet* 2000;**9**(11):1665–70.

- 18 **Hadlaczky G**, Went M, Ringertz NR. Direct evidence for the non-random localization of mammalian chromosomes in the interphase nucleus. *Exp Cell Res* 1986;**167**(1):1–15.
- 19 **Hager HD**, Schroeder-Kurth TM, Vogel F. Position of chromosomes in the human interphase nucleus. An analysis of nonhomologous chromatid translocations in lymphocyte cultures after Trenimon treatment and from patients with Fanconi's anemia and Bloom's syndrome. *Hum Genet* 1982;**61**(4):342–56.
- 20 **Simon I**, Tenzen T, Reubinoff BE, Hillman D, McCarrey JR, Cedar H. Asynchronous replication of imprinted genes is established in the gametes and maintained during development. *Nature* 1999;**401**(6756):929–32.
- 21 **Azuara V**, Brown KE, Williams RRE, Webb N, Dillon N, Festenstein R, Buckle V, Merckenschlager M, Fisher AG. Heritable gene silencing in lymphocytes delays chromatid resolution without affecting the timing of DNA replication. *Nat Cell Biol* 2003;**5**(7):668–74.
- 22 **Woodfine K**, Fiegler H, Beare DM, Collins JE, McCann OT, Young BD, Debernardi S, Mott R, Dunham I, Carter NP. Replication timing of the human genome. *Hum Mol Genet* 2004;**13**(2):191–202.
- 23 **Liapunova NA**. Organization of replication units and DNA replication in mammalian cells as studied by DNA fiber autoradiography. *Int Rev Cytol* 1994;**154**:261–6.
- 24 **Edenberg HJ**, Huberman JA. Eukaryotic chromosome replication. *Annu Rev Genet* 1975;**9**:245–84.
- 25 **Nogami M**, Nogami O, Kagotani K, Okumura M, Taguchi H, Ikemura T, Okumura K. Intranuclear arrangement of human chromosome 12 correlates to large-scale replication domains. *Chromosoma* 2000;**108**:514–22.
- 26 **Cimbora DM**, Schübeler D, Reik A, Hamilton J, Francastel C, Epner EM, Groudine M. Long-distance control of origin choice and replication timing in the human beta-globin locus are independent of the locus control region. *Mol Cell Biol* 2000;**20**(15):5581–91.
- 27 **Chen C**, Kolodner RD. Gross chromosomal rearrangements in *Saccharomyces cerevisiae* replication and recombination defective mutants. *Nat Genet* 1999;**23**:81–5.
- 28 **Myung K**, Pennaneach V, Kats ES, Kolodner RD. *Saccharomyces cerevisiae* chromatin-assembly factors that act during DNA replication function in the maintenance of genome stability. *Proc Natl Acad Sci U S A* 2003;**100**(11):6640–5.
- 29 **Borde V**, Goldman ASH, Lichten M. Direct coupling between meiotic DNA replication and recombination initiation. *Science* 2000;**290**:806–9.
- 30 **Murakami H**, Borde V, Shibata T, Lichten M, Ohta K. Correlation between premeiotic DNA replication and chromatin transition at yeast recombination initiation sites. *Nucleic Acids Res* 2003;**31**(14):4085–90.
- 31 **Cha RS**, Weiner BM, Keeney S, Dekker J, Kleckner N. Progression of meiotic DNA replication is modulated by interchromosomal interaction proteins, negatively by Spo11p and positively by Rec8p. *Genes Dev* 2000;**14**:493–503.
- 32 **Parisi S**, McKay M, Molnar M, Thompson M, van der Spek P, van Drunen-Schoenmaker E, Kanaar R, Lehmann E, Hoeijmakers J, Kohli J. Rec8p, a meiotic recombination and sister chromatid cohesion phosphoprotein of the Rad21p family conserved from fission yeast to humans. *Mol Cell Biol* 1999;**19**:3515–28.
- 33 **Devriendt K**, Fryns JP, Mortier G, van Thienen MN, Keymolen K. The annual incidence of DiGeorge/velocardiofacial syndrome. *J Med Genet* 1998;**35**(9):789–90.