

## Contents



### Original articles

- 609** Correlations between clinical severity, genotype and muscle pathology in limb girdle muscular dystrophy type 2A  
*M Fanin, L Nardetto, A C Nascimbeni, E Tasca, M Spinazzi, R Padoan, C Angelini*
- 615** Congenital ichthyosis: mutations in *ichthyin* are associated with specific structural abnormalities in the granular layer of epidermis  
*J Dahlqvist, J Klar, I Hausser, I Anton-Lamprecht, M Hellström Pigg, T Gedde-Dahl Jr, A Gånemo, A Vahlquist, N Dahl*
- 621** Mutation detection in the *ABCC6* gene and genotype–phenotype analysis in a large international case series affected by pseudoxanthoma elasticum  
*E G Pfendner, O M Vanakker, S F Terry, S Vourthis, P E McAndrew, M R McClain, S Fratta, A-S Marais, S Hariri, P J Coucke, M Ramsay, D Viljoen, P F Terry, A De Paepe, J Uitto, L G Bercovitch*
- 629** Molecular karyotyping in patients with mental retardation using 100K single-nucleotide polymorphism arrays  
*J Hoyer, A Dreweke, C Becker, I Göhring, C T Thiel, M M Peippo, R Rauch, M Hofbeck, U Trautmann, C Zweier, M Zenker, U Hüffmeier, C Kraus, A B Ekici, F Rüschemdorf, P Nürnberg, A Reis, A Rauch*

### Short report

- 637** Isolated imprinting mutation of the *DLK1/GTL2* locus associated with a clinical presentation of maternal uniparental disomy of chromosome 14  
*I K Temple, V Shrubbs, M Lever, H Bullman, D J G Mackay*

### Letters to JMG

- 641** Phenotypic diversity of Menkes disease in mottled mice is associated with defects in localisation and trafficking of the ATP7A protein  
*B-E Kim, M J Petris*
- 647** Deletion of a 760 kb region at 4p16 determines the prenatal and postnatal growth retardation characteristic of Wolf-Hirschhorn syndrome  
*D Concolino, E Rossi, P Strisciuglio, M A Iembo, R Giorda, R Ciccone, R Tenconi, O Zuffardi*

- 651** *SOS1* is the second most common Noonan gene but plays no major role in cardio-facio-cutaneous syndrome  
*M Zenker, D Horn, D Wieczorek, J Allanson, S Pauli, I van der Burgt, H-G Doerr, H Gaspar, M Hofbeck, G Gillissen-Kaesbach, A Koch, P Meinecke, S Mundlos, A Nowka, A Rauch, S Reif, C von Schnakenburg, H Seidel, L-E Wehner, C Zweier, S Bauhuber, V Matejas, C P Kratz, C Thomas, K Kutsche*
- 657** Mutation analysis of *NPHP6/CEP290* in patients with Joubert syndrome and Senior-Løken syndrome  
*J Helou, E A Otto, M Attanasio, S J Allen, M A Parisi, I Glass, B Utsch, S Hashmi, E Fazzi, H Omran, J F O'Toole, J A Sayer, F Hildebrandt*
- 664** Stability of the m.8993T→G mtDNA mutation load during human embryofetal development has implications for the feasibility of prenatal diagnosis in NARP syndrome  
*J Steffann, N Gigarel, J Corcos, M Bonnière, F Encha-Razavi, M Sinico, S Prevot, Y Dumez, A Yamgnane, R Frydman, A Munnich, J P Bonnefont*
- 670** The *IRF5* polymorphism in type 1 diabetes  
*H-Q Qu, L Marchand, R Grabs, C Polychronakos*

### Miscellaneous

- 628** Correction

### Online mutation report

- e89** The T/G–13915 variant upstream of the lactase gene (*LCT*) is the founder allele of lactase persistence in an urban Saudi population  
*F Imtiaz, E Savilahti, A Sarnesto, D Trabzuni, K Al-Kahtani, I Kagevi, M S Rashed, B F Meyer, I Järvelä*

### Correspondence

- e90** Anthropometric evaluation of children with *SHOX* mutations can be used as indication for genetic studies in children of short stature  
*A A L Jorge, I J P Arnhold*
- e91** Anthropometric evaluation of children with *SHOX* mutations can be used as indication for genetic studies in children of short stature: reply from authors  
*W Blum, G Rappold*