



Clinical spectrum of *SIX3*-associated mutations in holoprosencephaly: correlation between genotype, phenotype and function

F Lacbawan,^{1,2} B D Solomon,¹ E Roessler,¹ K El-Jaick,¹ S Domené,¹ J I Vélez,¹ N Zhou,¹ D Hadley,¹ J Z Balog,¹ R Long,¹ A Fryer,³ W Smith,⁴ S Omar,⁵ S D McLean,⁶ K Clarkson,⁷ A Lichty,⁷ N J Clegg,⁸ M R Delgado,⁸ E Levey,⁹ E Stashinko,⁹ L Potocki,¹⁰ M I VanAllen,¹¹ J Clayton-Smith,¹² D Donnai,¹² D W Bianchi,¹³ P B Juliusson,¹⁴ P R Njølstad,^{14,15} H G Brunner,¹⁶ J C Carey,¹⁷ U Hehr,¹⁸ J Müsebeck,¹⁹ P F Wieacker,²⁰ A Postra,²¹ R C M Hennekam,²² M-J H van den Boogaard,²³ A van Haeringen,²⁴ A Paulussen,²⁵ J Herbergs,²⁵ C T R M Schrandt-Stumpel,²⁵ A R Janecke,²⁶ D Chitayat,²⁷ J Hahn,²⁸ D M McDonald-McGinn,²⁹ E H Zackai,²⁹ W B Dobyns,³⁰ M Muenke¹

► Additional data are published online only at <http://jmg.bmj.com/content/vol46/issue6>

For numbered affiliations see end of article

Correspondence to:
Maximilian Muenke, MD,
Medical Genetics Branch,
National Human Genome
Research Institute, National
Institutes of Health, 35 Convent
Drive, MSC 3717, Building 35,
Room 1B-203, Bethesda, MD
20892-3717; mamuenke@mail.nih.gov

The first two authors contributed equally to this work.

Received 31 October 2008
Revised 23 December 2008
Accepted 21 January 2009
Published Online First 2 April 2009

ABSTRACT

Background: Holoprosencephaly (HPE) is the most common structural malformation of the human forebrain. There are several important HPE mutational target genes, including the transcription factor *SIX3*, which encodes an early regulator of Shh, Wnt, Bmp and Nodal signalling expressed in the developing forebrain and eyes of all vertebrates.

Objective: To characterise genetic and clinical findings in patients with *SIX3* mutations.

Methods: Patients with HPE and their family members were tested for mutations in HPE-associated genes and the genetic and clinical findings, including those for additional cases found in the literature, were analysed. The results were correlated with a mutation-specific functional assay in zebrafish.

Results: In a cohort of patients ($n = 800$) with HPE, *SIX3* mutations were found in 4.7% of probands and additional cases were found through testing of relatives. In total, 138 cases of HPE were identified, 59 of whom had not previously been clinically presented. Mutations in *SIX3* result in more severe HPE than in other cases of non-chromosomal, non-syndromic HPE. An over-representation of severe HPE was found in patients whose mutations confer greater loss of function, as measured by the functional zebrafish assay. The gender ratio in this combined set of patients was 1.5:1 (F:M) and maternal inheritance was almost twice as common as paternal. About 14% of *SIX3* mutations in probands occur de novo. There is a wide intrafamilial clinical range of features and classical penetrance is estimated to be at least 62%.

Conclusions: Our data suggest that *SIX3* mutations result in relatively severe HPE and that there is a genotype–phenotype correlation, as shown by functional studies using animal models.

Holoprosencephaly (HPE) is the most common structural malformation of the human forebrain and occurs after failed or abbreviated midline cleavage of the developing brain (cortex and subcortical structures) during the third and fourth weeks of gestation. HPE occurs in up to 1 in 250

gestations, but only 1 in 8000 live births.^{1,2} Classically, three degrees of severity, defined by the extent of brain malformation, have been described. In the most severe form, alobar HPE, there is a single ventricle and no interhemispheric fissure. The olfactory bulbs and tracts and the corpus callosum are typically absent and dark-grey nuclei are not separated. In semilobar HPE, the most common type of HPE in neonates who survive to clinical examination, findings include partial cortical separation with absent or hypoplastic olfactory structures and corpus callosum. In lobar HPE, findings include separate ventricles but incomplete frontal cortical separation. Recently, an additional milder form, called middle interhemispheric variant (MIHV) has been delineated, in which the posterior frontal and parietal lobes are incompletely separated and the corpus callosum may be hypoplastic.^{3–6} Of patients with HPE who survived the neonatal period, alobar, semilobar and lobar HPE occurred in 21%, 60% and 19%, respectively.⁷ A separate study showed that of patients with non-chromosomal, non-syndromic HPE (including both living children and deceased fetuses), alobar, semilobar and lobar HPE occurred in 22%, 45% and 33%, respectively.⁸

Clinical features may be qualitatively predicted by the specific neuroanatomical abnormalities present. These features can include characteristic craniofacial anomalies, ophthalmological abnormalities such as colobomata or microphthalmia, severe mental retardation or developmental delay, pituitary dysfunction including diabetes insipidus, oromotor dysfunction, dysautonomia and seizures. Severely affected patients do not typically survive beyond early infancy; however, less severely affected patients may have normal life-spans. Though there are exceptions, more severe brain anomalies correlate with more severe clinical sequelae and shorter life-spans.^{7–10}

Craniofacial findings tend to correlate with the type and severity of brain anomalies. In HPE caused by single-gene mutations, facial findings may




Figure 1 Patients with mutations in *SIX3*, arranged by HPE type (neuroimaging does not correspond with patients). All patients had point mutations in *SIX3* with the exception of the patient shown in the lower row, second from left, who had a complex cytogenetic rearrangement including deletion of 2p21. MRI on patients in row 2 courtesy of the Carter Centers.^{19–48} MIHV, middle interhemispheric variant.

additionally correlate with the causative gene. Patients with alobar HPE may be found to have cyclopia or synophthalmia (fusion of the optic vesicles and incomplete development of separate eyes), proboscis (a tubular nasal appendage appearing above the fused eyes), severe microcephaly and bilateral cleft lip and palate. Features in less severely affected patients may include microcephaly, hypotelorism, a flat nasal bridge and cleft lip or palate. Findings at the least severe end of the phenotypically recognisable spectrum (sometimes termed a “microform” of HPE, seen in patients without the cardinal central nervous system findings of HPE) may include solitary maxillary central incisor, hypotelorism and microcephaly. However, exceptions are often seen; a patient with severe HPE may have relatively subtle facial dysmorphism and can have macrocephaly (as opposed to the more common microcephaly) due to hydrocephalus.^{4–9} Overall, facial anomalies have been reported in approximately 80% of patients with HPE and often lead to the diagnosis.⁸

HPE is aetiologically associated with teratogens such as maternal diabetes mellitus and alcohol, and has been reported in cases of prenatal exposure to pharmaceutical agents such as retinoic acid and the statin class of drugs, and to infections including cytomegalovirus, toxoplasmosis, and rubella.^{9–11–12} Up to half of patients with HPE have a numerical or structural chromosomal abnormality, whereas up to a quarter have HPE as part of a recognisable syndrome.^{15–15} HPE may also be due to single-gene mutations; < 25% of cases of HPE result from single-gene mutations in the currently commercially tested HPE-associated genes *SHH*, *ZIC2*, *SIX3* and *TGIF*.⁸

Sonic Hedgehog (*SHH*) was the first causative gene identified in human HPE.¹⁶ Since then, single mutations in at least 10 other genes have been purported to cause HPE.^{9–17–24} In families in which these mutations were found, HPE seems to be inherited in an autosomal dominant fashion. Large kindreds segregating HPE-associated mutations show both incomplete penetrance and highly variable expressivity, suggesting additional environmental or genetic influences superimposed on the haploinsufficient state.²³ According to this “multiple-hit” model, a mutation in a HPE-associated gene is necessary but not sufficient for HPE, and other genes or environmental factors are required for complete phenotypic feature.²⁵

SIX3 has been reported as the third most common cause of HPE due to single-gene mutations.⁸ The vertebrate *Six* genes encode a family of related transcription factors that are orthologues of the *sine oculis* (“without eyes”) gene in *Drosophila*, and are expressed in the developing fly’s visual system, suggesting partially conserved phylogenetic roles. Indeed, vertebrate *Six3* has been shown to be involved in midline forebrain and eye formation in several organisms including mouse, chick, *Xenopus* and zebrafish.^{26–28} Proteins encoded by the *Six* genes characteristically contain a DNA-binding homeobox domain, and an upstream SIX domain that can recruit additional factors to accomplish transcriptional activation or repression.^{29–30} Known biological properties of vertebrate *Six3* include transcriptional repression of BMP, Wnt and Nodal targets through complex(s) formed with the general co-repressor Groucho.^{31–33} It also forms a complex with a different factor, geminin, which can influence a cell’s fate towards differentiation rather than proliferation in the early expanding forebrain territory.³⁴ Presumably through a distinct set of co-factors, the *Six3* transcription factor can instead activate lens specification genes during eye formation.³⁵ Finally, *Six3* acts as a direct regulator of Sonic hedgehog expression in the ventral forebrain.^{36–37}

SIX3 was first identified as a candidate gene because of patients with HPE who had cytogenetic anomalies involving 2p21. *SIX3*, which has the appropriate spatial and temporal expression pattern to result in HPE, was the most attractive candidate gene in the interval.^{19–38}

We report our findings in 59 new patients with *SIX3* mutations and an additional 79 cases collected from the literature, and correlate the clinical phenotypes with *SIX3* genotypes and functional studies in the zebrafish.^{19–58–51} The clinical findings echo the incomplete penetrance and highly variable expressivity seen in HPE due to mutations in other genes. Recent work in animal models specifically supports the multiple-hit model in mammalian *SIX3*-associated HPE.³⁶ It has been suggested that *SIX3* mutations result in a more severe holoprosencephaly phenotype than do mutations in the other HPE-related genes although a mechanism that would explain this is not yet clear.^{4–9–48}

Table 1 Patient characteristics, mutations and functional data

Patient*	HPE type	DNA alteration (location) in <i>SIX3</i>	Predicted protein alteration in <i>SIX3</i>	Functional studies ^{†,47}	Gender	Proband/relative	Ref.
1	Not specified	c.109G→T (NTD)	p.G37C	0.5–0.9	Female	Proband	NA
2‡	Lobar	c.109G→T (NTD)	p.G37C	0.5–0.9	Male	Proband	42
3a‡	MIHV	c.109G→T (NTD)	p.G37C	0.5–0.9	Female	Proband	NA
3b‡	None	c.109G→T (NTD)	p.G37C	0.5–0.9	Female	Mother	NA
4a‡	Alobar	c.206G→A (NTD)	c.G69D	>0.9	Female	Proband	48
		c.406dupGC (SD)	FS	<0.5			
4b‡	None	c.206G→A (NTD)	c.G69D	>0.9	Male	Father	48
		c.406dupGC (SD)	FS	<0.5			48
4c	None	c.206G→A (NTD)	c.G69D	>0.9	Female	Paternal grandmother	48
		c.406dupGC (SD)	FS	<0.5	Female		48
5a‡	Alobar	c.214G→C (NTD)	p.A72P	<0.5	Female	Proband	48
5b‡	None	c.214G→C (NTD)	p.A72P	<0.5	Female	Mother	48
6	Alobar	c.235A→G (NTD)	p.M79V	Unknown	Not specified	Proband	47
7a‡	Alobar	Not specified (SD)	p.V92G	<0.5	Male	Proband	45
7b‡	Alobar	Not specified (SD)	p.V92G	<0.5	Male	Brother	45
7c‡	Alobar	Not specified (SD)	p.V92G	<0.5	Male	Brother	45
7d‡	Microform	Not specified (SD)	p.V92G	<0.5	Female	Mother	45
8‡	Semilobar	c.275T→G (SD)	p.V92G	<0.5	Female	Proband	43
9‡	Semilobar	c.278C→A (SD) (<i>PTCH1</i> : c.1165G→A)	p.A93D (<i>PTCH1</i> : p.A393T)	<0.5	Female	Proband	22 42
10a‡	Alobar	c.311A→G (SD)	p.D104G	0.5–0.9	Male	Proband	48
10b‡	None	c.311A→G (SD)	p.D104G	0.5–0.9	Male	Father	48
11‡	Not specified	c.313A→G (SD)	p.I105V	>0.9	Not specified	Proband	43
12‡	Lobar	c.338G→A (SD)	p.W113X	<0.5	Male	Proband	42
13a‡	Not specified	c.339G→A (SD)	p.W113X	<0.5	Male	Proband	NA
13b‡	Not specified	Not tested	Not tested	Not tested	Male	Brother	NA
13c‡	Microform	c.339G→A (SD)	p.W113X	<0.5	Female	Mother	NA
14a‡	Alobar	c.339G→T (SD)	p.W113C	<0.5	Female	Proband	42
14b‡	Microform	c.339G→T (SD)	p.W113C	<0.5	Female	Mother	42
14c‡	Microform	c.339G→T (SD)	p.W113C	<0.5	Female	Maternal aunt	42
14d‡	Microform	c.339G→T (SD)	p.W113C	<0.5	Male	Maternal uncle	42
14e‡	None	c.339G→T (SD)	p.W113C	<0.5	Female	Maternal aunt	42
14f‡	Microform	c.339G→T (SD)	p.W113C	<0.5	Female	Maternal grandmother	42
14g‡	Microform	Linkage only	Linkage only	Linkage only	Male	Maternal great uncle	42
14h‡	None	Linkage only	Linkage only	Linkage only	Female	Maternal first cousin	42
14i‡	Not specified	Linkage only	Linkage only	Linkage only	Female	Maternal half-sister	42
14j	Not specified	Not tested	Not tested	Not tested	Female	Maternal first cousin	42
14k‡	Not specified	Not tested	Not tested	Not tested	Male	Maternal first cousin	42
14l‡	Not specified	Not tested	Not tested	Not tested	Female	Maternal first cousin	42
14m	Not specified	Not tested	Not tested	Not tested	Female	Maternal first cousin, once removed	42
14n	Not specified	Not tested	Not tested	Not tested	Male	Maternal great uncle	42
14o	Not specified	Not tested	Not tested	Not tested	Not specified	Maternal great-grandparent	42
15‡	Semilobar	c.341C→T (SD)	p.S114L	<0.5	Female	Proband	42
16a‡	Lobar	c.385G→T (SD)	p.E129X	<0.5	Male	Proband	42
16b‡	Alobar	c.385G→T (SD)	p.E129X	<0.5	Female	Sister	42
16c‡	Microform	c.385G→T (SD)	p.E129X	<0.5	Male	Father	42
17	Not specified	c.389C→A (SD)	p.S130X	<0.5	Male	Proband	42
18a‡	Alobar	c.404G→C (SD)	p.R135P	<0.5	Male	Proband	48

Continued

Table 1 Continued

Patient*	HPE type	DNA alteration (location) in <i>SIX3</i>	Predicted protein alteration in <i>SIX3</i>	Functional studies ^{†,47}	Gender	Proband/relative	Ref.
18b‡	None	c.404G→C (SD)	p.R135P	<0.5	Female	Mother	48
19	Not specified	c.404_407dupGCGC (SD)	FS	<0.5	Female	Proband	42
20‡	Semilobar	c.405_409CGCCG (SD)	FS	<0.5	Male	Proband	42
21a	Not specified	c.463_465delCAC (SD)	p.H155del	Unknown	Not specified	Proband	47
21b	Not specified	c.463_465delCAC (SD)	p.H155del	Unknown	Female	Mother	47
22‡	Not specified	c.469T→A (SD)	p.F157I	<0.5	Male	Proband	42
23a‡	Alobar	c.507delG (SD)	FS	<0.5	Female	Proband	48
23b‡	None	c.507delG (SD)	FS	<0.5	Male	Father	48
24a‡	Lobar	c.515C→T (SD)	p.A172V	0.5–0.9	Male	Proband	42
24b‡	Semilobar	Not tested	Not tested	Not tested	Female	Sister	42
24c‡	Not specified	Not tested	Not tested	Not tested	Female	Mother	42
25a	Alobar	c.518A→C (SD)	p.H173P	0.5–0.9	Female	Proband	45
25b	None	c.518A→C (SD)	p.H173P	0.5–0.9	Female	Mother	45
26a	Semilobar	c.518A→C (SD)	p.H173P	0.5–0.9	Female	Proband	43
26b	None	c.518A→C (SD)	p.H173P	0.5–0.9	Female	Mother	43
26c	None	c.518A→C (SD)	p.H173P	0.5–0.9	Female	Maternal aunt	43
26d	Not specified	Not tested	Not tested	Not tested	Not specified	Maternal first cousin	43
26e	Not specified	Not tested	Not tested	Not tested	Not specified	Maternal first cousin	43
26f	Not specified	Not tested	Not tested	Not tested	Not specified	Maternal first cousin	43
27‡	Not specified	c.520T→C (SD)	p.Y174H	0.5–0.9	Female	Proband	42
28a‡	Alobar	c.542_576dup (SD)	FS	Unknown	Female	Proband	43 45
28b‡	None	c.542_576dup (SD)	FS	Unknown	Male	Father	43 45
29‡	Semilobar	c.551delC (SD)	FS	<0.5	Female	Proband	42
30a‡	Alobar	c.556_557dupGG (SD)	FS	<0.5	Female	Proband	44 45
30b‡	Alobar	c.556_557dupGG (SD)	FS	<0.5	Male	Brother	44 45
30c‡	Alobar	c.556_557dupGG (SD)	FS	<0.5	Female	Sister	44 45
30d‡	Not specified	c.556_557dupGG (SD)	FS	<0.5	Female	Sister	44 45
30e‡	Microform	c.556_557dupGG (SD)	FS	<0.5	Male	Father	44 45
31a‡	Alobar	Not tested	Not tested	Not tested	Female	Proband	42
31b‡	Semilobar	c.582dup C (SD)	FS	<0.5	Female	Sister	42
31c‡	Microform	c.582dup C (SD)	FS	<0.5	Male	Father	42
32a‡	Semilobar	c.605C→T (SD)	p.T202I	<0.5	Female	Proband	43 45
32b‡	None	c.605C→T (SD)	p.T202I	<0.5	Female	Mother	43 45
33a‡	Semilobar	c.619G→T (HD)	p.E207X	<0.5	Male	Proband	42
33b‡	Alobar	c.619G→T (HD)	p.E207X	<0.5	Female	Sister	42
34	Not specified	c.637T→G (HD)	p.F213V	<0.5	Male	Proband	42
35‡	Lobar	c.652C→T (HD)	p.R218W	0.5–0.9	Female	Proband	42
36‡	Not specified	c.653G→C (HD)	p.R218P	<0.5	Female	Proband	42
37‡	Not specified	c.676C→G (HD)	p.L226V	>0.9	Female	Proband	19 52
38	Not specified	c.680A→C (HD)	p.Q227P	0.5–0.9	Not specified	Proband	42
39‡	Microform	c.686C→T (HD)	p.P229L	Unknown	Male	Proband	46
40‡	Alobar	c.692C→G (HD)	p.P231R	<0.5	Female	Proband	45
41‡	Alobar	c.698_706del (HD)	p.N233_S235del	<0.5	Female	Proband	19 39
42	Not specified	c.GC718_719AA (HD)	p.A240K	<0.5	Not specified	Proband	42
43a‡	Not specified	c.721C→T (HD)	p.Q241X	Unknown	Male	Proband	42
43b	Not specified	Not tested	Not tested	Unknown	Not specified	Sibling	42
43c‡	None	c.721C→T (HD)	p.Q241X	Unknown	Female	Mother	42
44	Not specified	c.730G→T (HD)	p.G244C	<0.5	Not specified	Proband	42
45	Not specified	c.736delA (HD)	FS	Unknown	Not specified	Proband	47
46	Not specified	c.743_745delCAG (HD) (TGIF: c.478C→T)	p.248K delQ (TGIF: p.160S)	Unknown	Female	Proband	NA
47a‡	Microform	c.749T→C (HD)	p.V250A	0.5–0.9	Male	Proband	19
47b‡	None	c.749T→C (HD)	p.V250A	0.5–0.9	Female	Mother	19
47c‡	None	c.749T→C (HD)	p.V250A	0.5–0.9	Female	Maternal aunt	19
47d‡	Alobar	c.749T→C (HD)	p.V250A	0.5–0.9	Not specified	Sibling	19

Continued

Table 1 Continued

Patient*	HPE type	DNA alteration (location) in SIX3	Predicted protein alteration in SIX3	Functional studies† ⁴⁷	Gender	Proband/relative	Ref.
47e‡	Alobar	c.749T→C (HD)	p.V250A	0.5–0.9	Not specified	Sibling	19
47f	Alobar	c.749T→C (HD)	p.V250A	0.5–0.9	Not specified	Sibling	19
48	Not specified	c.762T→A (HD)	p.F254L	0.5–0.9	Male	Proband	42
49‡	Not specified	c.769C→T (HD)	p.R257W	<0.5	Female	Proband	42
50a	Semilobar	c.769C→T (HD)	p.R257W	<0.5	Female	Proband	43 45
50b	Alobar	Not tested	Not tested	Not tested	Male	Brother	43 45
50c‡	None	c.769C→T (HD)	p.R257W	<0.5	Male	Father	43 45
51a	Lobar	c.769C→G (HD)	p.R257G	Unknown	Not specified	Proband	47
51b	Not specified	c.769C→G (HD)	p.R257G	Unknown	Male	Father	47
52‡	Semilobar	c.770G→C (HD)	p.R257P	>0.9	Female	Proband	19 52
53	Not specified	c.773G→C (HD)	p.R258L	<0.5	Not specified	Proband	42
54	Not specified	c.785G→A (HD)	p.R262H	0.5–0.9	Not specified	Proband	42
55a	Not specified	Not tested	Not tested	Unknown	Not specified	Proband	47
55b	None	Not tested	Not tested	Unknown	Not specified	Obligate carrier	47
55c	Not specified	c.806G→C (CTD)	p.R269T	Unknown	Male	Grandfather	47
56a‡	Semilobar	c.806G→T (CTD)	p.R269M	Unknown	Female	Proband	47 51
56b‡	Alobar	Not tested	Not tested	Unknown	Male	Sibling	47
56c‡	Lobar	c.806G→T (CTD)	p.R269M	Unknown	Male	Sibling	47
56d‡	Microform	c.806G→T (CTD)	p.R269M	Unknown	Female	Mother	47 51
57	Semilobar	c.806+1G→T (CTD)	Splice	Unknown	Not specified	Proband	47
58‡	Not specified	c.G807C (CTD)	p.R269S	0.5–0.9	Male	Proband	42
59	Not specified	c.820_832delinsCTGGACCT (CTD)	p.A274X	Unknown	Male	Proband	NA
60	Semilobar	c.850G→C (CTD) (ZIC2: c.910T→C)	p.A284P (ZIC2: p.VW304R)	Unknown	Female	Proband	NA
61‡	Semilobar	c.890C→T (CTD)	p.P297L	0.5–0.9	Female	Proband	42
62‡	Semilobar	c.944C→T (CTD)	p.T315I	Unknown	Female	Proband	48
63	Not specified	Deletion	NA	Unknown	Not specified	Proband	47
64	Not specified	Microdeletion by qPCR	NA	Unknown	Not specified	Proband	40 41
65	Not specified	Microdeletion by qPCR	NA	Unknown	Not specified	Proband	40 41
66	Not specified	Microdeletion by qPCR	NA	Unknown	Not specified	Proband	41
67	Not specified	Microdeletion by qPCR	NA	Unknown	Female	Proband	41
68‡	Semilobar	del(2)(p2101p2109)	NA	Unknown	Female	Proband	38 49
69‡	None	del(2)(p21)	NA	Unknown	Male	Proband	38
70‡	Semilobar	del(2)(p21p23)	NA	Unknown	Male	Proband	38
71‡	Semilobar	del(2)(p16p22)	NA	Unknown	Female	Proband	38
72	Lobar	del(2)(p21p22.2)	NA	Unknown	Female	Proband	38 50
73‡	Alobar	del(2)(p21p22.1)	NA	Unknown	Male	Proband	38
74‡	Semilobar	inv ins(2)(p21q24q13)	NA	Unknown	Female	Proband	38
75‡	Semilobar	t(1;2)(p21;p21)	NA	Unknown	Male	Proband	38
76	Alobar	t(1;2)(p22.3;p21)	NA	Unknown	Male	Proband	38
77‡	Lobar	del(2)(p16.3p21)§	NA	Unknown	Male	Proband	NA

CTD, C-terminal domain; HD, homeodomain; HPE, holoprosencephaly; MIHV, middle interhemispheric variant; NA, not applicable (unpublished when seeming in the Reference column); NTD, N-terminal domain; qPCR, quantitative PCR; SD, SIX domain.

*Each kindred is numbered separately; within a kindred, individual members have separate letter identifier.

†Protein Activity Index.

‡Clinical information available.

§Complex rearrangement additionally resulting in chromosome 2 inversion, translocation involving chromosomes 7, 13 and 18 and deletions of regions of chromosomes 7 and 18.

METHODS

The study protocol was approved by the National Human Genome Research Institute, and informed consent was obtained from all participants or their guardians.

Of the 65 cases for whom inheritance was known, 57.8% had maternal inheritance, 26.6% had paternal inheritance, 13.8% were de novo and the condition in 2 maternal half-siblings in one family seems to be due either an undetected germline mutation or parental mosaicism (the mother had negative mutation testing on peripheral blood analysis). We found no significant association between HPE severity and the parent of origin.

Of the 113 patients for whom gender was known, 40.7% were male and 59.3% were female, giving male:female ratio of 1:1.5. By χ^2 analysis, there was a statistically significant difference in the increased prevalence of affected females overall ($\chi^2 = 3.903$, $p = 0.0482$). Of the 61 kindreds in which the gender of the proband was known, 59.0% were female and 41.0% were male, giving a male:female ratio of 1:1.4. There was no significant difference in the increased prevalence of affected female probands ($\chi^2 = 1.984$, $p = 0.159$).

Of those kindreds with molecularly identified specific mutations in the proband, 40.3% had multiple affected family members identified, though familial testing or clinical description

Table 2 Mutations (n = 63)

	n (%)
Location	
N-terminal domain	6 (9.5)
SIX domain	27 (42.9)
Homeodomain	22 (34.9)
C-terminal domain	8 (12.7)
Exon 1	57 (90.5)
Exon 2	6 (9.5)
Mutation type	
Missense	44 (69.8)
Nonsense	6 (9.5)
Duplication→ FS	6 (9.5)
Deletion→FS	3 (4.8)
Splice mutation	1 (1.6)
9 bp deletion → 3 aa deletion	1 (1.6)
3 bp deletion → 1 aa deletion	1 (1.6)
Deletion/insertion → nonsense	1 (1.6)

aa, Amino acid; FS, frameshift; MIHV, middle interhemispheric variant.

was not available in all cases. Of the cases with described phenotypes, the penetrance of recognised phenotypic effect of mutations in *SIX3* was estimated to be at least 82% (n = 111). Within kindreds with multiple affected members and where clinical description was available, penetrance was estimated at 73% (n = 61). However, testing was often performed only on individuals with the phenotype and their parents. Many others were ascertained only because of the presence of a relative with severe HPE and only on later examination were considered to have microform HPE (eg, hypotelorism). Considering these patients to be “unaffected” results in a penetrance estimate of 62% (n = 61) (fig 1, table 1).

Mutations

Of those with molecularly identified mutations, the 62 kindreds encompassed 63 mutations in *SIX3* (kindred 4 had two mutations in *SIX3*), of which 93.7% were unique. Three unrelated kindreds had the same mutation in the N-terminal domain, two had the same mutation in the SIX domain and two had the same mutation in the homeodomain. Three patients had mutations in

two HPE genes: one (patient 9) in both *SIX3* and *PTCH*, one (patient 46) in both *SIX3* and *TGIF* and one (patient 60) in both *SIX3* and *ZIC2* (tables 1, 2; figs 2, 3).

Clinical features

Patients with mutations in *SIX3* (not including cytogenetic cases) had a significantly different distribution of HPE types than previously published cases of non-chromosomal, non-syndromic HPE ($\chi^2 = 24.179$, $p < 1 \cdot 10^{-4}$).⁸ Overall, patients with *SIX3* mutations had a higher proportion of severe HPE (table 3).


The quality of patient data was highly variable, so it is difficult to accurately calculate the prevalence of specific phenotypic features. However, we present the most commonly reported craniofacial anomalies (table 4). Severe craniofacial findings such as cyclopia and proboscis were only reported with alobar HPE. Overall, the severity of facial dysmorphisms seemed to correlate with the degree of brain anomalies—for example, the degree of hypotelorism was more pronounced in patients with alobar HPE.

In terms of clinical features other than craniofacial anomalies, the most commonly reported findings, in decreasing order of prevalence are: mental retardation or developmental delay, seizure disorder and diabetes insipidus (supplemental tables 1a,b online).

Functional studies

Using the functional analysis developed by Domené *et al*,⁴² 46 mutations (representing 99 patients with HPE who had neuroimaging performed or who were neurologically normal with *SIX3* mutations) had functional studies performed using a zebrafish assay.⁴² Functional study results were divided into three categories, with protein activity described relative to a normal control (value of 1.0): (1) protein activity <0.5 (alleles with the least functional activity), (2) protein activity 0.5–0.9 (alleles with moderate functional activity) and (3) protein activity >0.9 (alleles with near-normal functional activity). One kindred (kindred 4) had two separate mutations in *SIX3* and was categorised as belonging to the functional group with the more severe impairment of the two.

Including those patients who had alobar, semilobar and lobar HPE types with a functional protein index <0.5, there was an

Figure 2 Results of mutation studies.




Figure 3 Known mutations in *SIX3*, showing holoprosencephaly and mutation type. MIHV, middle interhemispheric variant. Numbers refer to kindreds.

unequal distribution of HPE types ($\chi^2 = 12.071$, $p = 0.0039$), with more severe HPE over-represented. Of those with functional protein index <0.5 and HPE on neuroimaging, 60.7% had alobar, 32.1% had semilobar and 7.1% had lobar HPE. Within the group of patients whose mutations had a functional protein index of 0.5–0.9, a significantly unequal distribution of HPE types was not found ($\chi^2 = 0.727$, $p = 0.8054$). (fig 4, table 5).

Including those with the most severe types of HPE (alobar, semilobar and lobar) and functional protein index either <0.5 and 0.5–0.9, we performed the linear trend alternative to independence test,⁵² using SAS V9.1 software. As there was only one person with functional protein index >0.9 whose HPE type was known, this category was not included in the statistical analysis. Our results showed that, within the group of patients whose mutations had a functional protein index <0.9 there was a significant correlation between HPE types and the functional protein index ($\chi^2 = 12.99$, $p = 0.0003$). That is, the functional protein index is an explanatory variable with ordered categories for traditional HPE types.

DISCUSSION

We present 138 cases of HPE with mutations in *SIX3*, many of whom have not been previously clinically described; this is the largest comprehensive evaluation of a cohort of patients with mutations in a gene involved in HPE. Analysis of this group allows several conclusions to be made.

First, as has been previously posited, our study shows that mutations in *SIX3* correlate with more severe HPE than with non-chromosomal, non-syndromic HPE overall.^{4 9 48} Despite the overall high severity, the clinical features are highly variable. Within large kindreds, people with *SIX3* mutations include both phenotypically normal people and people with severe holoprosencephaly incompatible with life. The basis of this variable expressivity is largely unknown. One potential explanation is a multi-hit mechanism, examples of which are the three cases where mutations were found in *SIX3* and in one other HPE-associated gene. In these patients, mutations in *SIX3* may be necessary but not sufficient for HPE. Another insult, either genetic (eg, changes in at least one other HPE-associated gene) or environmental (eg, gestational diabetes mellitus) must also occur.

Second, mutations in *SIX3* result in holoprosencephaly through a loss-of-function mechanism. Of patients whose mutations result in the greatest functional impairment in the zebrafish assay, protein activity correlates with human HPE severity.⁴² This analysis allows for a more refined genotype–phenotype discussion. Instead of correlating the gene involved or the location or type of the mutation with severity of phenotype, we can begin to predict features based on functional analyses.

Third, these patients do confirm the idea that in HPE, “the face predicts the brain”. That is, more severe facial dysmorphisms (such as cyclopia) tend to correlate with more severe HPE.

Table 3 Distribution of HPE types (n = 138)

HPE type	n (% of total) (% of cases with known HPE type)
Alobar	27 (19.6, 29.3)
Semilobar	22 (15.9, 23.9)
Lobar	9 (6.5, 9.8)
MIHV	1 (0.7, 1.1)
Microform	13 (9.4, 14.1)
None*	20 (14.5, 21.7)
Not specified	46 (33.3)

MIHV, middle interhemispheric variant.

*No evidence for penetrance.

Table 4 Reported* phenotypic findings (n = 91)

Finding	Prevalence (%)
Craniofacial anomalies	
Hypotelorism	44.0
Microcephaly	36.3
Cleft lip and/or palate	35.2
Flat nasal bridge/absent nasal septum	17.6
Philtral agenesis/hypoplasia	13.2
Coloboma	9.9
Solitary maxillary central incisor	8.8
Cyclopia	6.6

*It is likely that many of these findings, such as hypotelorism, occur more often than was reported.




Figure 4 Functional results and HPE types, showing overrepresentation of severe HPE of whose mutation resulted in the greatest functional impairment by zebrafish assay.⁴² MIHV, middle interhemispheric variant.

Although this adage does not apply to all cases of HPE, it does seem to hold true in HPE due to *SIX3* mutations.

Fourth, we estimated penetrance at 82% of all cases, 73% of cases from well-described families with multiple mutation carriers and 62% of those diagnosed on clinical grounds alone. In the practice of clinical genetics, this lowest estimate could be considered the most accurate. However, this latter penetrance estimate may be inaccurate for at least two reasons. First, the quality of clinical data was variable and subtle signs of midline defects may have been missed, resulting in an under-estimate of penetrance. Second, as full testing was not available on family members in many cases, there may have been many cases with *SIX3* mutations who were not ascertained. This would result in an overestimate of penetrance. If more members of families could be tested, it would be possible to better calculate the penetrance, quantify expressivity and calculate the rate of sporadic mutations.

Fifth, it is interesting that more female patients have been reported with *SIX3* mutations. The female predominance has been reported in some, but not all studies of HPE.^{4 8} The fact that the difference does reach significance when all mutation

carriers are considered lends credence to the idea that being female is somehow protective. One explanation is that the *SIX3* mutations in males may be more likely to be embryonic-lethal. However, in patients who survived long enough to have the type of HPE identified, the correlation between gender and severity of HPE was not significant.

One shortcoming of this report is that most of the patients discussed here were not seen in person, although we did perform the laboratory analysis to identify the *SIX3* mutation in approximately half the cases. Details of the data available from referring clinicians varied greatly; in some cases, extensive medical records were sent, whereas in other cases, relatively little was available. For this reason, it is difficult to make certain genotype–phenotype correlations. However, it can be noted that in cases for which more clinical information was available, the severity of features often seemed more impressive. Thus, these data may under-represent the severity of the clinical features of *SIX3* mutations. On the other hand, many people who had either no or very subtle features were ascertained only because they were related to a person with the same mutation but much more severe presenting signs. Following this logic, these data may over-represent severity.

Despite the challenges interpreting the large and varied data, the number of patients and families described here greatly enriches our understanding of the spectrum of features in patients with mutations in *SIX3*. These considerations argue for the importance of a combined and comprehensive approach to clinical and genetic studies of complex genetic disorders such as HPE.

Author affiliations: ¹Medical Genetics Branch, National Human Genome Research Institute, National Institutes of Health, Bethesda, MD, USA; ²Department of Pathology, State University of New York-Downstate Medical Center, Brooklyn, NY, USA; ³Department of Clinical Genetics, Royal Liverpool Children's Hospital (Alder Hey), Liverpool, UK; ⁴Department of Genetics, Maine Medical Center, Portland, ME, USA; ⁵Department of Neonatology, College of Human Medicine, Michigan State University, MI, USA; ⁶Departments of Pediatrics and Genetics, San Antonio Military Medical Center, San Antonio, TX, USA; ⁷Department of Clinical Genetics, Greenwood Genetic Center, Columbia, SC, USA; ⁸Department of Neurology, Texas Scottish Rite Hospital for Children, University of Texas Southwestern Medical Center, Dallas, TX, USA; ⁹Kennedy Krieger Institute, Johns Hopkins University, Baltimore, MD, USA; ¹⁰Department of Molecular and Human Genetics, Baylor College of Medicine and Texas Children's Hospital, Houston, TX, USA; ¹¹Department of Medical Genetics, University of British Columbia, Vancouver, BC, Canada; ¹²Academic Department of Medical Genetics and Regional Genetic Services, St Mary's Hospital, University of Manchester, Manchester, UK; ¹³Division of Genetics, Department of Pediatrics, Tufts University School of Medicine, Boston, MA, USA; ¹⁴Department of Paediatrics, Haukeland University Hospital, Bergen, Norway; ¹⁵Department of Clinical Medicine, University of Bergen, Bergen, Norway; ¹⁶Department of Human Genetics, Radboud University Nijmegen Medical Center, Nijmegen, Netherlands; ¹⁷Division of Medical Genetics, University of Utah Medical Center, Salt Lake City, UT, USA; ¹⁸Department of Human Genetics, University of Regensburg, Regensburg, Germany; ¹⁹Center for Human Genetics, University of Bremen, Bremen, Germany; ²⁰Institute of Human Genetics, University of Munster, Munster, Germany; ²¹Department of Clinical Genetics, Academic Medical Center, Amsterdam, Netherlands; ²²Institute of Child Health, Great Ormond Street Hospital for Children, London, UK; ²³Department of Medical Genetics, University Medical Centre Utrecht, Netherlands; ²⁴Department of Human and Clinical Genetics, Leiden University Medical Center, Leiden, Netherlands; ²⁵Department of Clinical Genetics, Academic Hospital Maastricht, Maastricht, Netherlands; ²⁶Division of Clinical Genetics, Innsbruck Medical University, Innsbruck, Austria; ²⁷Prenatal Diagnostics and Medical Genetics, Mt. Sinai Hospital, Toronto, ON, Canada; ²⁸Department of Neurology, Stanford University School of Medicine, Palo Alto, CA, USA; ²⁹Division of Human Genetics, Children's Hospital of Philadelphia, Philadelphia, PA, USA; ³⁰Departments of Human Genetics, Neurology and Pediatrics, The University of Chicago, Chicago, IL, USA

Acknowledgements: We would like to express our deep gratitude to the patients and families who participated in these studies. We also thank all of the members of the Carter Centers for Brain Research in Holoprosencephaly and Related Malformations. We appreciate the contribution of Dr D A T Cummings for

Table 5 Functional result vs. HPE type (n = 99)

	<0.5	0.5-0.9	>0.9	Total
Alobar	17	5	0	22
Semilobar	9	3	1	13
Lobar	2	3	0	5
MIHV	0	1	0	1
Microform	10	1	0	11
None	9	7	0	16
Not specified	19	10	2	31
Total	66	30	3	99

MIHV, middle interhemispheric variant.

his assistance with statistical analysis and J Fekacs for her assistance with illustrations.

Funding: This research was supported by the Division of Intramural Research, National Human Genome Research Institute, National Institutes of Health, Department of Health and Human Services, USA

Competing interests: There are no competing interests.

Patient consent: Obtained.

REFERENCES

- Matsunaga E, Shiota K. Holoprosencephaly in human embryos: epidemiologic studies of 150 cases. *Teratology* 1977;**16**:261–72.
- Leoncini E, Baranello G, Orioli IM, Annerén G, Bakker M, Bianchi F, Bower C, Canfield MA, Castilla EE, Cocchi G, Correa A, De Vigan C, Doray B, Feldkamp ML, Gatt M, Irgens LM, Lowry RB, Maraschini A, Mc Donnell R, Morgan M, Mutchnick O, Poetsch S, Riley M, Ritvanen A, Gnansia ER, Scarano G, Sipek A, Tenconi R, Mastroiacovo P. Frequency of holoprosencephaly in the International Clearinghouse Birth Defects Surveillance Systems: Searching for population variations. *Birth Defects Res A Clin Mol Teratol* 2008;**82**:585–91.
- Muenke M, Beachy PA. Genetics of ventral forebrain development and holoprosencephaly. *Curr Opin Genet Dev* 2000;**10**:262–9.
- Cohen MM Jr. Holoprosencephaly: clinical, anatomic, and molecular dimensions. *Birth Defects Res A Clin Mol Teratol* 2006;**76**:658–73.
- Barkovich AJ, Simon EM, Clegg NJ, Kinsman SL, Hahn JS. Analysis of the cerebral cortex in holoprosencephaly with attention to the sylvian fissures. *AJNR Am J Neuroradiol* 2002;**23**:143–50.
- Barkovich AJ, Quint DJ. Middle interhemispheric fusion: an unusual variant of holoprosencephaly. *AJNR Am J Neuroradiol* 1993;**14**:431–40.
- Hahn JS, Barkovich AJ, Stashinko EE, Kinsman SL, Delgado MR, Clegg NJ. Factor analysis of neuroanatomical and clinical characteristics of holoprosencephaly. *Brain Dev* 2006;**28**:413–9.
- Lazaro L, Dubourg C, Pasquier L, Le Duff F, Blayau M, Durou MR, de la Pintièrre AT, Aguilera C, David V, Odent S. Phenotypic and molecular variability of the holoprosencephalic spectrum. *Am J Med Genet* 2004;**129A**:21–4.
- Dubourg C, Bendavid C, Pasquier L, Henry C, Odent S, David V. Holoprosencephaly. *Orphanet J Rare Dis* 2007;**2**:8.
- Plawner LL, Delgado MR, Miller VS, Levey EB, Kinsman SL, Barkovich AJ, Simon EM, Clegg NJ, Sweet VT, Stashinko EE, Hahn JS. Neuroanatomy of holoprosencephaly as predictor of function: beyond the face predicting the brain. *Neurology* 2002;**59**:1058–66.
- Edison RJ, Muenke M. Mechanistic and epidemiologic considerations in the evaluation of adverse birth outcomes following gestational exposure to statins. *Am J Med Genet* 2004;**131**:287–98.
- Edison RJ, Muenke M. Central nervous system and limb anomalies in case reports of first-trimester statin exposure. *N Engl J Med* 2004;**350**:1579–82. Erratum in: *N Engl J Med* 2005;**352**:2759.
- Cohen MM Jr. Perspectives on holoprosencephaly: Part I. Epidemiology, genetics, and syndromology. *Teratology* 1989;**40**:211–35.
- Cohen MM Jr, Sulik KK. Perspectives on holoprosencephaly: Part II. Central nervous system, craniofacial anatomy, syndrome commentary, diagnostic approach, and experimental studies. *J Craniofac Genet Dev Biol* 1992;**12**:196–244.
- Croen LA, Shaw GM, Lammer EJ. Holoprosencephaly: epidemiologic and clinical characteristics of a California population. *Am J Med Genet* 1996;**64**:465–72.
- Roessler E, Belloni E, Gaudenz K, Jay P, Berta P, Scherer SW, Tsui LC, Muenke M. Mutations in the human Sonic Hedgehog gene cause holoprosencephaly. *Nat Genet* 1996;**14**:357–60.
- Brown SA, Warburton D, Brown LY, Yu CY, Roeder ER, Stengel-Rutkowski S, Hennekam RC, Muenke M. Holoprosencephaly due to mutations in ZIC2, a homologue of Drosophila odd-paired. *Nat Genet* 1998;**20**:180–3.
- Wallis D, Muenke M. Mutations in holoprosencephaly. *Hum Mutat* 2000;**16**:99–108.
- Wallis DE, Roessler E, Hehr U, Nanni L, Wiltshire T, Richieri-Costa A, Gillissen-Kaesbach G, Zackai EH, Rommens J, Muenke M. Mutations in the homeodomain of the human SIX3 gene cause holoprosencephaly. *Nat Genet* 1999;**22**:196–8.
- Gripp KW, Wotton D, Edwards MC, Roessler E, Ades L, Meinecke P, Richieri-Costa A, Zackai EH, Massagué J, Muenke M, Elledge SJ. Mutations in TGIF cause holoprosencephaly and link NODAL signalling to human neural axis determination. *Nat Genet* 2000;**25**:205–8.
- de la Cruz JM, Bamford RN, Burdine RD, Roessler E, Barkovich AJ, Donnai D, Schier AF, Muenke M. A loss-of-function mutation in the CFC domain of TDGF1 is associated with human forebrain defects. *Hum Genet* 2002;**110**:422–8.
- Ming JE, Kaupus ME, Roessler E, Brunner HG, Golabi M, Tekin M, Stratton RF, Sujansky E, Bale SJ, Muenke M. Mutations in PATCHED-1, the receptor for SONIC HEDGEHOG, are associated with holoprosencephaly. *Hum Genet* 2002;**110**:297–301.
- Roessler E, Du YZ, Mullor JL, Casas E, Allen WP, Gillissen-Kaesbach G, Roeder ER, Ming JE, Ruiz i Altaba A, Muenke M. Loss-of-function mutations in the human GLI2 gene are associated with pituitary anomalies and holoprosencephaly-like features. *Proc Natl Acad Sci USA* 2003;**100**:13424–9.
- Roessler E, Ouspenskaia MV, Karkera JD, Vélez JL, Kantipong A, Lacbawan F, Bowers P, Belmont JW, Towbin JA, Goldmuntz E, Feldman B, Muenke M. Reduced NODAL signaling strength via mutation of several pathway members including FOXH1 is linked to human heart defects and holoprosencephaly. *Am J Hum Genet* 2008;**83**:18–29.
- Ming JE, Muenke M. Multiple hits during early embryonic development: digenic diseases and holoprosencephaly. *Am J Hum Genet* 2002;**71**:1017–32.
- Gestri G, Carl M, Appolloni I, Wilson SW, Barsacchi G, Andreazzoli M. Six3 functions in anterior neural plate specification by promoting cell proliferation and inhibiting Bmp4 expression. *Development* 2005;**132**:2401–13.
- Kobayashi M, Toyama R, Takeda H, Dawid IB, Kawakami K. Overexpression of the forebrain-specific homeobox gene six3 induces rostral forebrain enlargement in zebrafish. *Development* 1998;**125**:2973–82.
- Oliver G, Mailhos A, Wehr R, Copeland NG, Jenkins NA, Gruss P. Six3, a murine homologue of the sine oculis gene, demarcates the most anterior border of the developing neural plate and is expressed during eye development. *Development* 1995;**121**:4045–55.
- Kawakami K, Ohto H, Takizawa T, Saito T. Identification and expression of six family genes in mouse retina. *FEBS Lett* 1996;**393**:259–63.
- Granadino B, Gallardo ME, Lopez-Rios J, Sanz R, Ramos C, Ayuso C, Bovolenta P, Rodriguez de Cordoba S. Genomic cloning, structure, expression pattern, and chromosomal location of the human SIX3 gene. *Genomics* 1999;**55**:100–5.
- Inbal A, Kim SH, Shin J, Solnica-Krezel L. Six3 represses nodal activity to establish early brain asymmetry in zebrafish. *Neuron* 2007;**55**:407–15.
- Kobayashi M, Nishikawa K, Suzuki T, Yamamoto M. The homeobox protein Six3 interacts with the Groucho corepressor and acts as a transcriptional repressor in eye and forebrain formation. *Dev Biol* 2001;**232**:315–26.
- Lagutin OV, Zhu CC, Kobayashi D, Topczewski J, Shimamura K, Puelles L, Russell HR, McKinnon PJ, Solnica-Krezel L, Oliver G. Six3 repression of Wnt signaling in the anterior neuroectoderm is essential for vertebrate forebrain development. *Genes Dev* 2003;**17**:368–79.
- Del Bene F, Tessmar-Raible K, Wittbrodt J. Direct interaction of geminin and Six3 in eye development. *Nature* 2004;**427**:745–9.
- Liu W, Lagutin OV, Mende M, Streit A, Oliver G. Six3 activation of Pax6 expression is essential for mammalian lens induction and specification. *EMBO J* 2006;**25**:5383–95.
- Geng X, Speirs C, Lagutin O, Inbal A, Liu W, Solnica-Krezel L, Jeong Y, Epstein DJ, Oliver G. Haploinsufficiency of Six3 fails to activate Sonic hedgehog expression in the ventral forebrain and causes holoprosencephaly. *Dev Cell* 2008;**15**:236–47.
- Jeong Y, Leskow FC, El-Jaick K, Roessler E, Muenke M, Yocum A, Dubourg C, Li X, Geng X, Oliver G, Epstein DJ. Regulation of a remote Shh forebrain enhancer by the Six3 homeoprotein. *Nat Genet* 2008;**40**:1348–53.
- Schell U, Wienberg J, Köhler A, Bray-Ward P, Ward DE, Wilson WG, Allen WP, Lebel RR, Sawyer JR, Campbell PL, Aughton DJ, Punnett HH, Lammer EJ, Kao FT, Ward DC, Muenke M. Molecular characterization of breakpoints in patients with holoprosencephaly and definition of the HPE2 critical region 2p21. *Hum Mol Genet* 1996;**5**:223–9.
- Nanni L, Croen LA, Lammer EJ, Muenke M. Holoprosencephaly: molecular study of a California population. *Am J Med Genet* 2000;**90**:315–19.
- Bendavid C, Dubourg C, Gicquel I, Pasquier L, Saugier-Verber P, Durou MR, Jaillard S, Frebourg T, Haddad BR, Henry C, Odent S, David V. Molecular evaluation of fetuses with holoprosencephaly shows high incidence of microdeletions in the HPE genes. *Hum Genet* 2006;**119**:1–8.
- Bendavid C, Haddad BR, Griffin A, Huizinga M, Dubourg C, Gicquel I, Cavalli LR, Pasquier L, Shanske AL, Long R, Ouspenskaia M, Odent S, Lacbawan F, David V, Muenke M. Multicolour FISH and quantitative PCR can detect submicroscopic deletions in holoprosencephaly patients with a normal karyotype. *J Med Genet* 2006;**43**:496–500.
- Domené S, Roessler E, El-Jaick K, Boorech J, Vélez JL, Bale S, Lacbawan F, Muenke M, Feldman B. Mutations in the human SIX3 gene in holoprosencephaly are loss-of-function. *Hum Mol Genet* 2008;**17**:3919–28.
- Dubourg C, Lazaro L, Pasquier L, Bendavid C, Blayau M, Le Duff F, Durou MR, Odent S, David V. Molecular screening of SHH, ZIC2, SIX3, and TGIF genes in patients with features of holoprosencephaly spectrum: Mutation review and genotype-phenotype correlations. *Hum Mutat* 2004;**24**:43–51.
- Pasquier L, Dubourg C, Blayau M, Lazaro L, Le Marec B, David V, Odent S. A new mutation in the six-domain of SIX3 gene causes holoprosencephaly. *Eur J Hum Genet* 2000;**8**:797–800.
- Pasquier L, Dubourg C, Gonzales M, Lazaro L, David V, Odent S, Encha-Razavi F. First occurrence of aprosencephaly/atelencephaly and holoprosencephaly in a family with a SIX3 gene mutation and phenotype/genotype correlation in our series of SIX3 mutations. *J Med Genet* 2005;**42**:e4.
- El-Jaick KB, Fonseca RF, Moreira MA, Ribeiro MG, Bolognese AM, Dias SO, Pereira ET, Castilla EE, Orioli IM. Single median maxillary central incisor: new data and mutation review. *Birth Defects Res A Clin Mol Teratol* 2007;**79**:573–80.
- Paulussen A, Tserpelis D, Spierts S, Smeets H, Herbergs J. Holoprosencephaly mutations in the Dutch population. Poster presented at the European Society of Human Genetics, Barcelona, Spain, 31 May to 3 June, 2008
- Ribeiro LA, El-Jaick KB, Muenke M, Richieri-Costa A. SIX3 mutations with holoprosencephaly. *Am J Med Genet* 2006;**140**:2577–83.
- Wilson WG, Shanks DE, Sudduth KW, Couper KA, McIlhenny J. Holoprosencephaly and interstitial deletion of 2(p2101p2109). *Am J Med Genet* 1989;**34**:252–4.

50. **Sawyer JR**, Jones E, Hawks FF, Quirk JG Jr, Cunniff C. Duplication and deletion of chromosome band 2(p21p22) resulting from a familial interstitial insertion (2;11)(p21;p15). *Am J Med Genet* 1994;**49**:422–7.
51. **Zwetsloot CP**, Brouwer OF, Maaswinkel-Mooy PD. Holoprosencephaly: variation of expression in face and brain in three sibs. *J Med Genet* 1989;**26**:274–6.
52. **Laflamme C**, Filion C, Labelle Y. Functional characterization of SIX3 homeodomain mutations in holoprosencephaly: interaction with the nuclear receptor NR4A3/NOR1. *Hum Mutat* 2004;**24**:502–8.
53. **Agresti A**. *An introduction to categorical data analysis*. New York: John Wiley and Sons, 1996.

BMJ Careers Fair

2–3 October 2009, Business Design Centre, London, UK

9–10 October 2009, Thinktank, Birmingham, UK

BMJ is the largest organiser of medical recruitment fairs across the UK. This year we are organising two careers fairs, in partnership with the London Deanery on 2–3 October in London, and the West Midlands Deanery on 9–10 October in Birmingham.

Whatever your grade or specialty there is a careers fair for you. You can:

- ▶ attend seminars on topics such as CV writing, interview skills, planning your career and working abroad
- ▶ visit exhibition stands to get careers advice, find a new job, identify alternative career pathways

It's free to attend the exhibition if you register online in advance. There is a small fee for attending our seminar programme.

Register online today at www.careersfair.bmj.com