



## SHORT REPORT

# Exome sequencing reveals a novel Fanconi group defined by XRCC2 mutation

Hanan E Shamseldin,<sup>1</sup> Mohamed Elfaki,<sup>2</sup> Fowzan S Alkuraya<sup>1,2,3</sup>

► Additional materials are published online only. To view these files please visit the journal online (<http://jmg.bmj.com/content/49/3.toc>).

<sup>1</sup>Department of Genetics, King Faisal Specialist Hospital and Research Center, Riyadh, Saudi Arabia

<sup>2</sup>Department of Pediatrics, King Khalid University Hospital and College of Medicine, King Saud University, Riyadh, Saudi Arabia

<sup>3</sup>Department of Anatomy and Cell Biology, College of Medicine, Alfaisal University, Riyadh, Saudi Arabia

## Correspondence to

Dr Fowzan S Alkuraya, Developmental Genetics Unit, King Faisal Specialist Hospital and Research Center, MBC-03 PO Box 3354, Riyadh 11211, Saudi Arabia; [falkuraya@kfsshr.edu.sa](mailto:falkuraya@kfsshr.edu.sa)

Received 23 October 2011

Accepted 14 December 2011

Published Online First

9 January 2012

## ABSTRACT

**Background** Fanconi anaemia (FA) is a group of disorders characterised by progressive bone marrow failure and a characteristic but variable craniofacial and skeletal involvement. Recessive mutations in any of 15 genes linked to FA lead to the pathognomonic increased susceptibility to double-strand DNA breaks.

**Methods** Autozygome and exome analysis of a patient with classic FA phenotype

**Results** The authors identified a novel truncating mutation in *XRCC2*. Consistent with the proposed causal link to FA, this gene is an essential non-redundant component of the RAD51 family of homologous repair proteins and its deficiency in a murine model has been shown to lead to a highly similar phenotype to that of this patient both at the cellular and organismal level.

**Conclusion** This study implicates *XRCC2* in the pathogenesis of FA and calls for further investigation of the potential contribution of *XRCC2* mutations to the overall mutational load of FA.

Fanconi anaemia (FA) is a group of disorders characterised by progressive bone marrow failure and a characteristic but variable craniofacial and skeletal involvement. Consistent with the pathognomonic finding of double-strand DNA breaks in the chromosomes of these patients when cultured with DNA interstrand crosslinking agents, all FA genes (15 to date) are known to play a role in dsDNA breaks repair.<sup>1 2</sup> Although recessive mutations in these genes account for the majority of FA cases, the genetic heterogeneity of this disorder has not yet been fully captured.

In the highly consanguineous population of Saudi Arabia, we have shown marked allelic as well as genetic heterogeneity for a large number of Mendelian disorders, and how autozygome analysis can be a powerful tool to identify the underlying genetic defect.<sup>3</sup> In this communication, we describe a Saudi child with a novel FA group in whom the combined use of autozygome and exome analysis led to the identification of a *XRCC2* mutation as the likely cause of his phenotype.

## SUBJECTS AND METHODS

### Human subject

The patient had standard haematological and clinical genetics evaluation. He and his parents were enrolled in an Institutional Review Board-approved research protocol with written informed consent. This was followed by blood extraction in EDTA (for DNA extraction) and sodium-heparin (for Epstein–Barr virus-transformed lymphoblast cell line) tubes.

## Autozygome analysis

This was performed as described before.<sup>4 5</sup> Briefly, DNA was extracted from the patient and his parents and processed for genotyping using the Axiom platform following the manufacturer's protocol (Affymetrix, Santa Clara, CA, USA). Resulting genotypes were analysed using autoSNPa (<http://dna.leeds.ac.uk/autosnpa/>) to determine the overall pattern of autozygosity as judged by the runs of homozygosity (ROH) per individual (autozygome).<sup>6</sup> We have only considered runs of homozygosity that are 2 Mb or more and contained at least 175 single-nucleotide polymorphisms to increase the likelihood that these ROH represent autozygous (identical by descent) rather than identity by state.


## Exome analysis

Exome capture was performed using the TruSeq Exome Enrichment kit (Illumina, San Diego, CA, USA) following the manufacturer's protocol. Samples were prepared as an Illumina sequencing library, and in the second step the sequencing libraries were enriched for the desired target using the Illumina exome enrichment protocol. The captured libraries were sequenced using the Illumina HiSeq 2000 sequencer. The reads are mapped against UCSC hg19 (<http://genome.ucsc.edu/>) by Burrows–Wheeler Aligner (<http://bio-bwa.sourceforge.net/>). The single-nucleotide polymorphisms and indels are detected by SAMTOOLS (<http://samtools.sourceforge.net/>).

## RESULTS

### Case report

Index is a 2.5-year-old Saudi child who was born to young healthy first cousin Saudi parents. The pregnancy was unremarkable and he was born at term. Length and head circumference were abnormally low at birth (length 41 cm, weight 2.6 kg and head circumference 30.5 cm). His physical examination revealed left facial nerve palsy, microcephaly and bilaterally absent thumbs. Investigations in the nursery revealed ectopic left kidney, haemodynamically stable patent ductus arteriosus, normal blood indices and blood karyotype. Skeletal survey revealed complete absence of the first metacarpal and scaphoid bones bilaterally, absent left radius and hypoplastic right radius. On follow-up visits, he displayed severe growth deficiency and microcephaly. In addition, the diepoxybutane (DEB) test showed marked increase in the frequency of dsDNA breaks (figure 1).



**Figure 1** (A) Facial photograph showing facial palsy. (B) and (C) Hand photographs documenting lack of thumbs. (D) x-Ray of the left hand showing agenesis of the radius. (E) Chromosomal breakage study with diepoxybutane (DEB) showing the numerous sites of dsDNA breaks (arrows) typical of Fanconi anaemia. (F) Sequence chromatogram showing a novel homozygous *XRCC2* mutation in the patient.

### Exome and autozygome analysis

In view of the genetic heterogeneity of FA, we took advantage of the consanguineous nature of the parents to pursue autozygome analysis in order to identify the most likely candidate.<sup>7</sup> Supplementary figure S1 and table S1 (available online only) summarise the overall ROH pattern of the patient. *FANCL* was the only autosomal recessive FA gene to reside within ROH but we failed to identify any mutation in that gene at the genomic or complementary DNA level. This led us to hypothesise that this child's FA is caused by a recessive mutation of a novel FA gene. Therefore, we set out to perform whole exome sequencing followed by autozygome filtering.<sup>4–5</sup> Supplementary table S2 (available online only) summarises the raw data of our exome sequencing. Our filtration strategy (summarised in supplementary figure S2, available online only) allowed us to narrow the search to three coding variants: *MTBP* (NM\_022045:c.A397G; p.(N133D)), *RGS3* (NM\_144488: c.C2882A; p.(T961N)) and *XRCC2* (NM\_005431.1:c.643C>T; p.(Arg215\*)). The first two missense variants were deemed unlikely because they were detected in 1/270 and 3/192 control Saudi samples, respectively. The truncating nonsense mutation in *XRCC2* was therefore the only likely variant left based on our filtration strategy.

### DISCUSSION

*XRCC2* is one of five RAD51 paralogs that act non-redundantly in the same pathway of homologous recombination repair (HR)

of dsDNA breaks.<sup>8</sup> It has been shown that *XRCC2* enhances the binding of RAD51 to sites of dsDNA breaks, a key step in HR, and truncation mapping is consistent with the hypothesis that our patient's truncation is pathogenic as it removes an important C-terminus, lack of which fully abolishes *XRCC2* activity.<sup>9</sup> *XRCC2* deficiency has been repeatedly shown at the cellular level to result in impaired HR and increased susceptibility to dsDNA breaks, consistent with our patient's chromosomal phenotype.<sup>8</sup> Importantly, *Xrcc2* knockout mice have embryonic lethality and abnormal patterning leading to multiple congenital malformations (including microcephaly and skeletal anomalies) that resemble our patient's phenotype.<sup>10</sup> Finally, the severe proliferation disadvantage of fibroblasts derived from these mice is also consistent with our observation on skin-derived fibroblasts from our patient.<sup>11</sup>

Our results, therefore, strongly implicate *XRCC2* in the pathogenesis of FA and call for further investigation of the potential contribution of *XRCC2* mutations to the overall mutational load of FA. This case shows the power of next generation sequencing when combined with autozygome analysis in identifying novel genetic causes of Mendelian disorders even on simplex cases.

**Acknowledgements** The authors would like to thank the family for their enthusiastic participation. They also wish to thank the genotyping and sequencing core facilities at KFSHRC for their technical help.

**Funding** This study was funded in part by KACST grant 09-MED941-20 (FSA), and a Dubai-Harvard Foundation for Medical Research Collaborative Grant (FSA).

**Competing interests** None.

**Patient consent** Obtained.

**Ethics approval** Ethics approval was provided by the King Faisal Specialist Hospital and Research Center Institutional Review Board.

**Contributors** HS performed the autozygome and exome analysis. MF provided clinical data. FSA supervised the entire study, provided clinical data, analysed data and wrote the paper.

**Provenance and peer review** Not commissioned; externally peer reviewed.

## REFERENCES

1. **Kee Y**, D'Andrea AD. Expanded roles of the Fanconi anemia pathway in preserving genomic stability. *Genes Dev* 2010;**24**:1680–94.
2. **Shimamura A**, Alter BP. Pathophysiology and management of inherited bone marrow failure syndromes. *Blood Rev* 2010;**24**:101–22.
3. **Alkuraya FS**. Autozygome decoded. *Genet Med* 2010;**12**:765–71.
4. **Shaheen R**, Faqeh E, Sunker A, Morsy H, Al-Sheddi T, Shamseldin HE, Adly N, Hashem M, Alkuraya FS. Recessive mutations in DOCK6, encoding the guanidine nucleotide exchange factor DOCK6, lead to abnormal actin cytoskeleton organization and Adams–Oliver syndrome. *Am J Hum Genet* 2011;**89**:328–33.
5. **Aldahmesh MA**, Khan AO, Mohamed JY, Alkuraya H, Ahmed H, Bobis S, Al-Mesfer S, Alkuraya FS. Identification of ADAMTS18 as a gene mutated in Knobloch syndrome. *J Med Genet* 2011;**48**:597–601.
6. **Carr IM**, Flintoff KJ, Taylor GR, Markham AF, Bonthron DT. Interactive visual analysis of SNP data for rapid autozygosity mapping in consanguineous families. *Hum Mutat* 2006;**27**:1041–6.
7. **Alkuraya FS**. Homozygosity mapping: one more tool in the clinical geneticist's toolbox. *Genet Med* 2010;**12**:236–9.
8. **Johnson RD**, Liu N, Jasin M. Mammalian XRCC2 promotes the repair of DNA double-strand breaks by homologous recombination. *Nature* 1999;**401**:397–9.
9. **Tambini CE**, Spink KG, Ross CJ, Hill MA, Thacker J. The importance of XRCC2 in RAD51-related DNA damage repair. *DNA Repair (Amst)* 2010;**9**:517–25.
10. **Deans B**, Griffin CS, Maconochie M, Thacker J. Xrcc2 is required for genetic stability, embryonic neurogenesis and viability in mice. *EMBO J* 2000;**19**:6675–85.
11. **Deans B**, Griffin CS, O'Regan P, Jasin M, Thacker J. Homologous recombination deficiency leads to profound genetic instability in cells derived from Xrcc2-knockout mice. *Cancer Res* 2003;**63**:8181–7.

# DIFFERENTIAL DIAGNOSIS

Trustworthy guidance on your iPhone



New app available now

Find out more at [bestpractice.bmj.com/differentials](http://bestpractice.bmj.com/differentials)

