Editor's choice
Scan to access more
free content

SHORT REPORT

Loss-of-function de novo mutations play an important role in severe human neural tube defects

Philippe Lemay,¹ Marie-Claude Guyot,¹ Élisabeth Tremblay,¹
Alexandre Dionne-Laporte,² Dan Spiegelman,² Édouard Henrion,² Ousmane Diallo,²
Patrizia De Marco,³ Elisa Merello,³ Christine Massicotte,¹ Valérie Désilets,¹
Jacques L Michaud,¹ Guy A Rouleau,² Valeria Capra,³ Zoha Kibar¹

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/jmedgenet-2015-103027>).

¹CHU Ste-Justine Research Center, Université de Montréal, Montréal, Québec, Canada

²Montreal Neurological Institute, McGill University, Montréal, Québec, Canada

³Istituto Giannina Gaslini, Genoa, Italy

Correspondence to

Dr Zoha Kibar, Department of Neurosciences, University of Montréal, CHU Sainte Justine Research Center, 3175 Cote-Ste-Catherine, Room 5999C, Montréal, Québec, Canada H3T 1C5; zoha.kibar@recherche-ste-justine.qc.ca

Received 16 January 2015

Revised 26 February 2015

Accepted 4 March 2015

Published Online First

24 March 2015

ABSTRACT

Background Neural tube defects (NTDs) are very common and severe birth defects that are caused by failure of neural tube closure and that have a complex aetiology. Anencephaly and spina bifida are severe NTDs that affect reproductive fitness and suggest a role for de novo mutations (DNMs) in their aetiology.

Methods We used whole-exome sequencing in 43 sporadic cases affected with myelomeningocele or anencephaly and their unaffected parents to identify DNMs in their exomes.

Results We identified 42 coding DNMs in 25 cases, of which 6 were loss of function (LoF) showing a higher rate of LoF DNM in our cohort compared with control cohorts. Notably, we identified two protein-truncating DNMs in two independent cases in *SHROOM3*, previously associated with NTDs only in animal models. We have demonstrated a significant enrichment of LoF DNMs in this gene in NTDs compared with the gene specific DNM rate and to the DNM rate estimated from control cohorts. We also identified one nonsense DNM in *PAX3* and two potentially causative missense DNMs in *GRHL3* and *PTPRS*.

Conclusions Our study demonstrates an important role of LoF DNMs in the development of NTDs and strongly implicates *SHROOM3* in its aetiology.

Neural tube defects (NTDs) are a group of congenital malformations affecting 1–2 individuals per 1000 births.¹ They are caused by an incomplete closure of the neural tube during embryogenesis.¹ The most frequent forms of NTDs are anencephaly and myelomeningocele (MMC) (or spina bifida), which are caused by a closure defect in the brain and the spinal cord region, respectively.¹ Children affected with anencephaly die early in development or soon after birth, while children affected with MMC survive but are affected by developmental physical defects with varying degrees of severity. Most cases of NTDs are sporadic and non-syndromic.² Periconceptional folic acid intake has been shown to reduce prevalence of NTDs by 50–70%,³ but a large amount of cases remain resistant to this preventive treatment urging the need for identification of other causative factors and development of novel preventive and counselling strategies.

NTDs have a strong genetic component with an estimated heritability of 60%,¹ but so far the

genetics of the disease remains largely unknown. Previous linkage studies in NTDs have identified candidate regions on chromosomes 2, 7 and 10 but failed to identify any causative NTD gene.⁴ Few common variants in folic acid-related genes have also been shown to be associated with NTDs in certain populations, but these variants seem to contribute only to a small part of the aetiology of the disease.⁵ Previous gene identification studies in NTDs have mainly adopted a candidate gene approach and focused on folate-related genes and on candidate genes from animal studies.^{1 2 5} Animal models have demonstrated an important role of the planar cell polarity pathway in the aetiology of NTDs.¹ Subsequent investigation of genes of this pathway, including *VANGL1* and *VANGL2*, in human NTDs has implicated them as risk factors in a small fraction of patients.¹ Generally, candidate gene studies in NTDs have faced limited success in identifying major causative genes predisposing to NTDs, suggesting the need for novel approaches. Several recent studies strongly suggest that de novo mutations (DNMs) represent a common cause of birth defects and neurodevelopmental diseases.⁶ DNM could provide a mechanism by which early-onset reproductively lethal diseases remain frequent in the population. Therefore, these variants are strong candidates for causing diseases that occur sporadically and that have a reduced reproductive fitness.⁶ Severe forms of NTDs, such as anencephaly and MMC, fall in this category and hence investigation of DNMs may therefore increase the chance of identifying loss of function (LoF) DNMs implicated in NTDs.

Forty-three families each composed of one affected child and two unaffected parents with no family history of NTDs were recruited through the Montreal Ste-Justine Hospital Spina Bifida Center, the 3D study of the Integrated Research Network in Perinatology of Quebec and Eastern Ontario and the Istituto Giannina Gaslini in Genoa, Italy. Detailed information including folate status, tissue of origin and type of NTDs of this cohort is summarised in online supplementary table S1. Briefly, all 43 cases were affected with NTDs including 35 MMC and 8 anencephaly cases. A total of 21 cases were fetuses and 55.6% took folate periconceptionally. Tissues from fetuses were all obtained following induced abortions. The average maternal and paternal ages were 30.0 ± 4.8 years and 30.7 ± 5.9 years, respectively.

CrossMark

To cite: Lemay P, Guyot M-C, Tremblay É, et al. *J Med Genet* 2015;52:493–497.

Exome data from 43 NTD trios were aligned to the GRCh37 genome reference and called variants were filtered for minimal quality (genotype quality (GQ) >20, total variant reads >3, total reads >7) and for absence in parents and public databases (1000 genome at <http://www.1000genomes.org/home> and Exome Variant Server at <http://evs.gs.washington.edu/EVS/>) to identify DNMs. Details of the alignment strategy and single-nucleotide variant (SNV) annotation are included in online supplementary data. Seventy variants were identified and validated by Sanger sequencing to minimise false negative results. This

approach resulted in the identification of 42 coding mutations in 25 trios including 3 nonsense, 3 frameshift, 30 missense and 6 synonymous (table 1). The average DNM rate per base pair was 1.62×10^{-8} , which was consistent with published ratios.^{7,8}

We identified six LoF DNMs defined as nonsense, frameshift and splicing variants, in six NTD trios resulting in a per base DNMs rate of 0.23×10^{-8} . This rate was higher than two previously published per base LoF DNMs rates, 0.05×10^{-8} (ref. 8) and 0.17×10^{-8} (ref. 9) but only reached significance ($p=0.002$) when compared to the lower rate, suggesting that larger cohorts

Table 1 De novo mutations identified in NTD-affected trios and confirmed by Sanger sequencing.

Family	NTD type	Chr	Position	Genes	cDNA change	Amino acid change	Polyphen HDIV*
<i>Loss-of-function de novo mutations (nonsense, frameshift, splicing)</i>							
Pr394	MMC	4	77 662 169	SHROOM3	c.2843_2844insG	p.L948fs	NA
PrZRV	Anen	4	77 660 502	SHROOM3	c.1176C>G	p.Y392X	NA
Pr134	MMC	2	223 161 800	PAX3	c.218C>A	p.S73X	NA
Pr201	MMC	15	44 116 692	MFAP1	c.69del7	p.K23fs	NA
Pr389	MMC	X	41 202 544	DDX3X	c.620dupA	p.Q207fs	NA
PrYPT	MMC	7	73 279 630	WBSCR28	c.380G>A	p.W127X	NA
<i>Missense and synonymous de novo mutations</i>							
Pr548	MMC	1	24 668 728	GRHL3	c.1171C>T	p.R391C	1
Pr548	MMC	19	39 798 985	LRFN1	c.1604C>T	p.T535I	1
Pr548	MMC	9	124 751 686	TLL11	c.1327A>G	p.N443D	0.968
Pr125	MMC	19	5 214 591	PTPR5†	c.4475G>A	p.R1049Q	0.998
Pr125	MMC	2	27 435 209	ATRAID	c.138G>A	p.A46A	NA
Pr122	MMC	12	123 341 629	HIP1R	c.1682G>T	p.G561V	0.019
Pr122	MMC	17	71 232 301	C17orf80	c.1441C>T	p.R481W	0.099
Pr122	MMC	X	53 592 096	HUWE1	c.6812G>A	p.S2271N	0
Pr134	MMC	14	105 179 874	INF2†	c.2971C>T	p.R991W	1
Pr191	MMC	8	10 464 772	RP1L1†	c.6836C>T	p.P2279L	0.053
Pr191	MMC	20	62 371 335	SLC24A4RG	c.70C>T	p.R24C	0.426
Pr20	MMC	3	38 317 786	SLC22A13	c.1246G>A	p.V416M	0.948
Pr20	MMC	16	31 383 022	ITGAX	c.2077C>G	p.Q693E	0.001
Pr201	MMC	2	47 703 654	MSH2†	c.1956A>G	p.Q652Q	NA
Pr202	MMC	6	30 122 164	TRIM10	c.1028A>T	p.D343V	1
Pr202	MMC	7	23 775 208	STK31	c.535A>C	p.I179L	0.039
Pr25	MMC	9	134 183 554	PPAPDC3†	c.696C>T	p.I232I	NA
Pr263	MMC	6	1 390 351	FOXF2	c.169G>A	p.A57T	0.01
Pr28	MMC	8	124 333 387	ATAD2	c.4160G>A	p.S1387N	0.001
Pr282	MMC	19	808 439	PTBP1	c.1233C>A	p.N411K	0.013
Pr389	MMC	7	100 285 176	GIGYF1	c.325C>T	p.P109S	0.728
Pr402	MMC	17	19 319 353	RNF112	c.1761C>T	p.A587A	NA
Pr402	MMC	3	49 775 724	IP6K1	c.355C>T	p.R119C	1
Pr402	MMC	4	48 424 093	SLAIN2	c.1745G>C	p.X582S	NA
Pr530	MMC	3	57 616 163	DENND6A	c.1605A>C	p.E535D	0.997
Pr551	MMC	9	130 279 261	FAM129B	c. 848C>T	p.A283V	0.004
Pr553	MMC	19	39 329 153	HNRNPL	c.1441C>T	p.R481W	1
Pr554	Anen	3	142 741 447	U2SURP	c.961G>A	p.G321S	1
Pr554	Anen	5	93 966 388	ANKRD32	c.371T>C	p.F124S	0.999
Pr67	MMC	2	44 566 318	PREPL	c.937C>G	p.L313V	1
Pr67	MMC	5	176 314 262	HK3	c.1677G>A	p.V559V	NA
Pr67	MMC	9	33 264 606	BAG1	c.67G>A	p.A23T	0.897
PrKKS	MMC	15	43 621 819	LCMT2†	c.869T>C	p.I290T	0.201
PrKKS	MMC	7	11 076 097	PHF14	c.1655G>C	p.R552P	0.998
PrTVB	MMC	22	41 558 745	EP300†	c.3690A>G	p.Q1230Q	NA
PrVWA	MMC	9	120 475 791	TLR4	c.1385C>T	p.A462V	0.006

*Probably damaging (polyphen HDIV ≥ 0.957), possibly damaging ($0.453 \leq$ polyphen HDIV ≤ 0.956); benign (polyphen HDIV ≤ 0.452).

†Mutation previously reported in the ExAC database (<http://exac.broadinstitute.org/>). Reported mutations frequencies are PTPRS (frequency: 0.000008368); INF2 (frequency: 0.000009451); RP1L1 (frequency: 0.0001411); MSH2 (frequency: 0.0001730); PPAPDC3 (frequency: 0.0003366); LCMT2 (frequency: 0.00008281); EP300 (frequency: 0.0000082). Anen, anencephaly; MMC, myelomeningocele; NA, non-applicable; NTD, neural tube defect.

of MMC and anencephaly are needed to confirm these initial findings. This increased rate suggests that LoF DNMs are an important part of the pathogenicity of NTDs. Details of all statistical analysis can be found in online supplementary data.

Notably, we identified two LoF DNMs in *SHROOM3* (NM_020859.3) in two unrelated families: one nonsense variant c.1176C>G (p.Y392X) and one frameshift variant c.2843_2844insG leading to a premature stop codon (p.L948fs) (figure 1B). The c.1176C>G variant was detected in the PrZRV proband affected with anencephaly with cranial fossa agenesis and facial dysmorphism (table 1). The mother took folic acid periconceptionally (see online supplementary table S1). The c.2843_2844insG variant was detected in the Pr394 proband affected with a thoracic MMC and Chiari type IV malformation (table 1). The mother did not take folic acid periconceptionally (see online supplementary table S1). None of these mutations were reported in the ExAC database (<http://exac.broadinstitute.org/>). *SHROOM3* is an actin-binding protein that is known to be a key regulator of apical constriction, a process by which cells convert their shape from cuboidal to wedge-like due to a decrease in their apical area.⁹ This function is essential for hinge-point formation and bending of the neural tube during its formation and closure in both vertebrate embryos.⁹ Recessive mutations in *SHROOM3* were previously associated to

heterotaxy, which represents a multiple congenital anomaly syndrome resulting from abnormalities of the proper specification of left-right asymmetry during embryonic development.¹⁰ The LoF DNM in this gene described in our study was detected in two patients with NTD who showed no sign of heterotaxy, and hence these NTD mutations might act in a haploinsufficient or a dominant negative manner. *Shroom3* exists in two isoforms that were demonstrated to have similar functions and expression patterns in the mouse and frog models.^{11 12} In the mouse gene trap mutant, both isoforms are knocked out, leading to exencephaly and spina bifida in homozygous embryos at a penetrance of 100% and 23%, respectively.¹¹ In our cohort, both truncating DNMs detected in *SHROOM3* seem to affect both isoforms since they map at positions 1176 and 2843 bp respectively 5' or inside the ASD1 domain (figure 1A). These two DNMs clearly remove important functional domains and might confer nonsense-mediated RNA decay and hence they are most likely LoF mutations. Furthermore, previous studies have demonstrated the potential of a truncated version of *Shroom3* to act in a dominant negative manner in *Xenopus*.¹² This supports the potential pathogenicity of these heterozygous mutations that could result in limited apical constriction causing the NTD.

The phenotypic variation between the two probands (MMC and anencephaly) who carry LoF DNM in *SHROOM3* could be

Figure 1 De novo mutations (DNMs) identified in *SHROOM3* and in *PAX3* in trios affected with neural tube defects (NTDs). (A) Protein schematic representation of the long and predicted short isoforms of *SHROOM3* indicating the position of the two DNMs identified in NTD probands. (B) Chromatograms of the three individuals of family PrZRV and Pr394 (from top to bottom: proband, mother, father) surrounding the *SHROOM3* c.1176C>G nonsense variant and the c.2843_2844insG frameshift variant. (C) Protein schematic representation of the long and predicted short isoforms of *PAX3* indicating the position of the DNM identified in NTD probands. (D) Chromatograms of the three individuals of family Pr134 (from top to bottom: proband, mother, father) surrounding the *PAX3* c.218C>A nonsense variant. HOX, paired-type homeodomain; PAX, paired box domain.

caused by many factors including genetic modifiers and environmental factors. The two variants might also have different pathogenic effects on the protein with different outcomes in the implicated developmental pathway. These factors could act alone or in combination to affect the clinical expressivity of the NTD phenotype.

Considering the estimated LoF DNM rates between 0.05×10^{-8} and 0.17×10^{-8} SNV per base^{7,8} and the length of *SHROOM3* long isoform (NM_020859.3: 5991 bp), we expect to find respectively between 0.00026 and 0.00088 LoF DNM in this gene in our cohort. However, we detected two DNMs in our cohort of 43 trios, resulting in a frequency of DNM per bp of 3.88×10^{-6} . This represents a significant enrichment of LoF DNMs in *SHROOM3* under the two rates with respective Bonferroni corrected p values of 6.77×10^{-4} and 0.008 using a two-tailed binomial exact test. A recently published gene-specific per trio mutation rate for LoF DNMs was also used to test the significant enrichment of LoF DNMs in *SHROOM3*.¹³ Under the *SHROOM3* gene-specific rate of 1.34×10^{-5} , we expected to find 0.00058 LoF DNMs in our cohort, which was significantly lower than our observed rate (p Bonferroni=0.003). Details of all statistical analyses can be found in online supplementary data.

Also importantly we detected one de novo stopgain in the NTD-associated gene *PAX3*, c.218C>A (p.S73X), in the Pr134 case affected with MMC with type II Arnold–Chiari malformation, hydrocephalus and Waardenburg syndrome (WS) (figure 1D). The mother did not take folic acid or multivitamins periconceptionally (see online supplementary table S1). This mutation was not reported in the ExAC database. *PAX3* is a paired box transcription factor that is a key developmental regulator of the neural crest and its derivatives and that plays an important role during neurogenesis and myogenesis. It contains two DNA binding domains, the paired box domain (PAX) and a paired-type homeodomain (HOX) that interact cooperatively for DNA binding.¹⁴ The putative truncated protein caused by the DNM p.S73X lacks all functional domains including both PAX and HOX (figure 1C) and might confer nonsense-mediated RNA decay.

In the *Spotch* mouse model, homozygous LoF mutations of *Pax3* cause spina bifida and other neural crest abnormalities. In humans, the relative contribution of *PAX3* to the overall burden of NTDs remains unclear.¹⁴ Heterozygous mutations in *PAX3* are known to cause WS, an autosomal-dominant condition that affects neural crest-derived structures and that is occasionally associated with NTDs. Few individuals with both WS and NTDs were demonstrated to carry heterozygous *PAX3* mutations or deletions.¹⁴ Our report of a new protein truncating SNV in *PAX3* in one patient with NTD provides additional evidence for a pathogenic role of this gene in spina bifida.

While *SHROOM3* and *PAX3* are by far the strongest candidate genes identified in this study, other interesting DNMs have been identified in our cohort (table 1). We detected three LoF DNMs, p.W127X in *WBSR28*, p.K23fs in *MFAP1* and p.Q207fs in *DDX3X*, with no previous association to NTDs and that were not reported in the ExAC database (table 1, see online supplementary figure S1). The *WBSR28* DNM was detected in the PrYPT proband affected with lumbosacral MMC and type II Arnold–Chiari malformation. This gene encodes a putative transmembrane protein of unknown function that maps to the region deleted in the Williams–Beuren syndrome (table 1). This syndrome is characterised by a range of phenotypes including mental retardation, dysmorphic facies, heart abnormalities, short stature and infantile hypocalcaemia. *WBSR28* is not the

main candidate gene in this disease.¹⁵ The *MFAP1* DNM was found in the Pr201 proband affected with lumbosacral MMC. This gene encodes the microfibrillar-associated protein 1 that represents an uncharacterised protein found in some human spliceosomal fractions.¹⁶ The *DDX3X* DNM was found in the Pr389 proband affected with lumbosacral MMC and hydrocephalus. This gene belongs to the DEAD-box proteins, a large family of ATP-dependent RNA helicases that participate in all aspects of RNA metabolism.¹⁷ While none of these three genes represent strong candidates for NTDs based on published data, the presence of LoF DNMs in these genes still suggests them as potentially interesting NTD candidates. Additional genetic studies in larger cohorts and functional studies are needed to validate their role in NTDs.

Two other interesting missense DNM identified in two NTD trios implicated genes whose orthologues cause NTD in mice: c.1171C>T (p.R391C) in *GRHL3* (*GRAINYHEAD-LIKE 3*) and c.4475G>A (p.R1492Q) in *PTPRS* (*PROTEIN TYROSINE PHOSPHATASE, RECEPTOR TYPE, S*) (see online supplementary figure S2). The *GRHL3* mutation was identified in the Pr548 proband affected with MMC with type II Arnold–Chiari malformation and hydromyelia. This mutation was not previously reported in the ExAC database. *Grhl3* is a transcription factor that plays an important role in epidermal integrity and wound healing. Importantly, null alleles at this gene caused mainly severe spina bifida and occasionally exencephaly. *Grhl3* was tightly linked to Curly tail that represents one of the most well-established mouse models for NTDs.¹⁸ The p.R391C maps to the DNA binding domain of the GRHL3 protein. The modified amino acid is highly conserved (see online supplementary figure S2), and the mutation is defined as probably damaging by PolyPhen-2 HDIV (table 1). Surprisingly, the same DNM p.R391C was detected in Van der Woude syndrome, the most common syndromic form of cleft lip and palate.¹⁹ While this could point towards a chance finding, we hypothesise that additional genetic and/or environmental factors may modify the phenotypic expression of the same *GRHL3* mutation in different individuals. The p.R1492Q in *PTPRS* was found in the Pr125 proband affected with lumbosacral MMC with type II Arnold–Chiari malformation. The mutation was previously reported on one allele in the ExAC database (<http://exac.broadinstitute.org/>), resulting in a frequency of 0.000008368. This gene is a phosphatase involved in regulating cell proliferation, cell adhesion and nervous system maturation and causes exencephaly in mice.²⁰ *PTPRS* fibronectin and Ig-like domains are extracellular and mainly involved in cell interaction, while the catalytic phosphatase domain is intracellular and mainly involved in signal transmission. The p.R1049Q variant was identified in a proband affected with MMC (table 1) and resides in a conserved region that forms part of the phosphatase domain (see online supplementary figure S2). It modifies a conserved arginine to a glutamine, a non-conservative substitution that removes the positive charge of the amino acid side chain, changes its size and is predicted to be probably damaging by PolyPhen-2 (table 1).

Molecular genetic studies of bigger cohorts and careful phenotyping are still needed to better understand the mechanism of action of potential DNMs detected in this study and to assess their role in the pathogenicity in NTDs. While these DNMs are most likely highly penetrant, it is possible that they act in concert with other events to cause the disease. This is consistent with the two-hit model that was initially proposed as a cancer mechanism and later suggested as a potential mechanism in complex diseases.²¹ This model was more recently explored in

explaining variable expressivity in severe developmental genomic disorders.²¹ In this model, a secondary insult is necessary to result in a different or more severe clinical manifestation of a complex disease. This insult occurs during development and could be genetic or epigenetic (germ line or somatic) or environmental. The two ‘hits’ could act independently or additively to each other, resulting in a phenotype that differs from either hit alone. Alternatively, the two hits could involve gene(s) from the same or similar biochemical pathway and hence could interact in an epistatic manner.

The novel approach of whole-exome sequencing has long been due in the genetic investigation of NTDs. This study uses this powerful approach in this complex trait and has successfully identified potential candidate DNMs in novel genes in the development of human NTDs. We have demonstrated the presence of LoF variants in five genes, have reported more LoF DNMs in our cohort than expected and have identified two of those mutations in orthologues of mouse NTD genes, suggesting the involvement of those variants in the aetiology of human NTDs. We have also demonstrated the presence of two independent protein truncating variants in *SHROOM3* in 43 trios and presented a statistically significant enrichment of LoF DNMs in this gene compared with control and gene-specific DNMs rates. Our data strongly suggest that highly penetrant pathogenic variants in this gene may account for a significant part of the genetic aetiology of severe forms of NTD. Further studies of this gene in a bigger cohort of sporadic cases may help better assess the significance of these findings.

Acknowledgements We would like to thank all participants in this study.

Contributors PL, JLM, GAR, VC and ZK designed the study. PL, M-CG and ET validated the mutations. PL, AD-L, DS, EH and OD analysed the genetic data. PL ran the statistical analysis. PDM, EM, CM, VD, VC and ZK recruited patients and provided clinical information. PL and ZK wrote the manuscript.

Funding CHU Ste-Justine fundation; Fonds de Recherche du Québec—Santé; Canadian Institutes of Health Research. This project was conducted as part of the research programme of the Integrated Research Network in Perinatology of Quebec and Eastern Ontario (IRNPQEO). This work was supported by the Canadian Institutes of Health Research (CRI 88413 and MOP 130411).

Competing interests ZK has a salary award from the “Fonds de Recherche du Québec—Santé”. JLM is a National Scientist of the Fonds de Recherche du Québec—Santé. PL is supported by “Fondation du CHU Ste-Justine” and “Fonds de Recherche du Québec—Santé”. Authors have no competing interest.

Patient consent Obtained.

Ethics approval CHU Sainte Justine Hospital (Protocols’ numbers: 2598 and 2899) and Istituto Giannina Gaslini, Genoa, Italy (protocol number: 213/2013).

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- Bassuk AG, Kibar Z. Genetic basis of neural tube defects. *Semin Pediatr Neurol* 2009;16:101–10.
- Copp AJ, Stanier P, Greene ND. Neural tube defects: recent advances, unsolved questions, and controversies. *Lancet Neurol* 2013;12:799–810.
- [No authors listed]. Prevention of neural tube defects: results of the Medical Research Council Vitamin Study. MRC Vitamin Study Research Group. *Lancet* 1991;338:131–7.
- Stamm DS, Siegel DG, Mehlretter L, Connelly JJ, Trott A, Ellis N, Zismann V, Stephan DA, George TM, Vekemans M, Ashley-Koch A, Gilbert JR, Gregory SG, Speer MC. Refinement of 2q and 7p loci in a large multiplex NTD family. *Birt Defects Res A Clin Mol Teratol* 2008;82:441–52.
- Zhang T, Lou J, Zhong R, Wu J, Zou L, Sun Y, Lu X, Liu L, Miao X, Xiong G. Genetic variants in the folate pathway and the risk of neural tube defects: a meta-analysis of the published literature. *PLoS ONE* 2013;8:e59570.
- Veltman JA, Brunner HG. De novo mutations in human genetic disease. *Nat Rev Genet* 2012;13:565–75.
- Sanders SJ, Murtha MT, Gupta AR, Murdoch JD, Raubeson MJ, Willsey AJ, Ercan-Sencicek AG, DiLullo NM, Parikshak NN, Stein JL, Walker MF, Ober GT, Teran NA, Song Y, El-Fishawy P, Murtha RC, Choi M, Overton JD, Bjornson RD, Carriero NJ, Meyer KA, Bilguvar K, Mane SM, Sestan N, Lifton RP, Günel M, Roeder K, Geschwind DH, Devlin B, State MW. De novo mutations revealed by whole-exome sequencing are strongly associated with autism. *Nature* 2012;485:237–41.
- Iossifov I, O’Roak BJ, Sanders SJ, Ronemus M, Krumm N, Levy D, Stessman HA, Witherspoon KT, Vives L, Patterson KE, Smith JD, Paeppe B, Nickerson DA, Dea J, Dong S, Gonzalez LE, Mandell JD, Mane SM, Murtha MT, Sullivan CA, Walker MF, Waqar Z, Wei L, Willsey AJ, Yamrom B, Lee Y, Grabowska E, Dalkic E, Wang Z, Marks S, Andrews P, Leotta A, Kendall J, Hakker I, Rosenbaum J, Ma B, Rodgers L, Troge J, Narzisi G, Yoon S, Schatz MC, Ye K, McCombie WR, Shendure J, Eichler EE, State MW, Wigler M. The contribution of de novo coding mutations to autism spectrum disorder. *Nature* 2014;515:216–21.
- Sawyer JM, Harrell JR, Shemer G, Sullivan-Brown J, Roh-Johnson M, Goldstein B. Apical constriction: a cell shape change that can drive morphogenesis. *Dev Biol* 2010;341:5–19.
- Tariq M, Belmont JW, Lalani S, Smolarek T, Ware SM. SHROOM3 is a novel candidate for heterotaxy identified by whole exome sequencing. *Genome Biol* 2011;12:R91.
- Hildebrand JD, Soriano P. Shroom, a PDZ domain-containing actin-binding protein, is required for neural tube morphogenesis in mice. *Cell* 1999;99:485–97.
- Haigo SL, Hildebrand JD, Harland RM, Wallingford JB. Shroom induces apical constriction and is required for hinge-point formation during neural tube closure. *Curr Biol CB* 2003;13:2125–37.
- Samocha KE, Robinson EB, Sanders SJ, Stevens C, Sabo A, McGrath LM, Kosmicki JA, Rehnström K, Mallick S, Kirby A, Wall DP, MacArthur DG, Gabriel SB, DePristo M, Purcell SM, Palotie A, Boerwinkle E, Buxbaum JD, Cook EH, Gibbs RA, Schellenberg GD, Sutcliffe JS, Devlin B, Roeder K, Neale BM, Daly MJ. A framework for the interpretation of de novo mutation in human disease. *Nat Genet* 2014;46:944–50.
- Greene NDE, Massa V, Copp AJ. Understanding the causes and prevention of neural tube defects: Insights from the splotch mouse model. *Birt Defects Res A Clin Mol Teratol* 2009;85:322–30.
- Hockenhull EL, Carette MJ, Metcalfe K, Donnai D, Read AP, Tassabehji M. A complete physical contig and partial transcript map of the Williams syndrome critical region. *Genomics* 1999;58:138–45.
- Hegele A, Kamburov A, Grossmann A, Sourlis C, Wowro S, Weimann M, Will CL, Pena V, Lührmann R, Stelzl U. Dynamic protein-protein interaction wiring of the human spliceosome. *Mol Cell* 2012;45:567–80.
- Soto-Rifo R, Ohlmann T. The role of the DEAD-box RNA helicase DDX3 in mRNA metabolism. *Wiley Interdiscip Rev RNA* 2013;4:369–85.
- Gustavsson P, Copp AJ, Greene NDE. Grainyhead genes and mammalian neural tube closure. *Birt Defects Res A Clin Mol Teratol* 2008;82:728–35.
- Peyrard-Janvid M, Leslie EJ, Kousa YA, Smith TL, Dunnwald M, Magnusson M, Lentz BA, Unneberg P, Fransson I, Koillinen HK, Rautio J, Pegelow M, Karsten A, Basel-Vanagaite L, Gordon W, Andersen B, Svensson T, Murray JC, Cornell RA, Kere J, Schutte BC. Dominant mutations in GRHL3 cause Van der Woude Syndrome and disrupt oral periderm development. *Am J Hum Genet* 2014;94:23–32.
- Uetani N, Bertozzi K, Chagnon MJ, Hendriks W, Tremblay ML, Bouchard M. Maturation of ureter-bladder connection in mice is controlled by LAR family receptor protein tyrosine phosphatases. *J Clin Invest* 2009;119:924–35.
- Kumar RA. Two-hit wonder: a novel genetic model to explain variable expressivity in severe pediatric phenotypes. *Clin Genet* 2010;78:517–9.