OPEN ACCESS

ORIGINAL ARTICLE

Genetic spectrum of Saudi Arabian patients with antenatal cystic kidney disease and ciliopathy phenotypes using a targeted renal gene panel

Mohamed H Al-Hamed,¹ Wesam Kurdi,² Nada Alsahan,² Zainab Alabdullah,³ Rania Abudraz,² Maha Tulbah,² Maha Alnemer,² Rubina Khan,² Haya Al-Jurayb,¹ Ahmed Alahmed,¹ Asma I Tahir,¹ Dania Khalil,¹ Noel Edwards,⁴ Basma Al Abdulaziz,⁵ Faisal S Binhumaid,¹ Salma Majid,¹ Tariq Faquih,⁵ Mohamed El-Kalioby,⁵ Mohamed Abouelhoda,^{1,5} Nada Altassan,^{1,5} Dorota Monies,^{1,5} Brian Meyer,^{1,5} John A Sayer,⁴ Mamdouh Albaqumi^{1,6}

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/jmedgenet-2015-103469>).

For numbered affiliations see end of article.

Correspondence to

Dr John A Sayer, Institute of Genetic Medicine, International Centre for Life, Newcastle University, Newcastle upon Tyne NE1 3BZ, UK; John.sayer@ncl.ac.uk

Received 21 August 2015

Revised 1 January 2016

Accepted 4 January 2016

Published Online First

9 February 2016

Open Access
Scan to access more
free content

CrossMark

To cite: Al-Hamed MH, Kurdi W, Alsahan N, et al. *J Med Genet* 2016;**53**:338–347.

ABSTRACT

Background Inherited cystic kidney disorders are a common cause of end-stage renal disease. Over 50 ciliopathy genes, which encode proteins that influence the structure and function of the primary cilia, are implicated in cystic kidney disease.

Methods To define the phenotype and genotype of cystic kidney disease in fetuses and neonates, we correlated antenatal ultrasound examination and postnatal renal ultrasound examination with targeted exon sequencing, using a renal gene panel. A cohort of 44 families in whom antenatal renal ultrasound scanning findings in affected cases included bilateral cystic kidney disease, echogenic kidneys or enlarged kidneys was investigated.

Results In this cohort, disease phenotypes were severe with 36 cases of stillbirth or perinatal death. Extra renal malformations, including encephalocele, polydactyly and heart malformations, consistent with ciliopathy phenotypes, were frequently detected. Renal gene panel testing identified causative mutations in 21 out of 34 families (62%), where patient and parental DNA was available. In the remaining 10 families, where only parental DNA was available, 7 inferred causative mutations were found. Together, mutations were found in 12 different genes with a total of 13 novel pathogenic variants, including an inferred novel variant in *NEK8*. Mutations in *CC2D2A* were the most common cause of an antenatal cystic kidney disease and a suspected ciliopathy in our cohort.

Conclusions In families with ciliopathy phenotypes, mutational analysis using a targeted renal gene panel allows a rapid molecular diagnosis and provides important information for patients, parents and their physicians.

INTRODUCTION

The formation of cysts in kidney is a disease phenotype common to many inherited human diseases.¹ Kidney cysts are fluid-filled epithelial lined structures arising from dilation in any part of the nephron or collecting duct. Cystic kidney disorders are a common cause of end-stage renal disease (ESRD). It is estimated that the prevalence of cystic

kidney disease is 4.81% in the Arabian Gulf countries.²

Ciliopathy syndromes are inherited syndromes that are frequently associated with cystic kidneys and to date, mutations in over 50 genes have been identified.³ These include autosomal-dominant polycystic kidney disease (ADPKD), autosomal-recessive polycystic kidney disease (ARPKD), various forms of nephronophthisis (NPHP), Joubert syndrome (JBTS), Meckel–Gruber syndrome (MKS), Bardet–Biedl syndrome (BBS) and many others.⁴ ADPKD is common and accounts for approximately 5–10% of the ESRD cases worldwide.⁵ Mutations in two genes, *PKD1* (85% of patients with ADPKD) and *PKD2* (15% of patients with ADPKD) underlie ADPKD.⁶ One to two per cent of patients with ADPKD may present as neonates with cystic kidneys.⁷ Biallelic mutations/variants in *PKD1* and *PKD2* have been described to give a severe neonatal onset of cystic kidney disease.^{8,9}

ARPKD is a rarer condition affecting 1 in every 20 000 live births.¹⁰ It may be diagnosed in utero or prenatally by sonography showing bilateral large echogenic kidneys, and oligohydramnios in the most severe cases. Mutations in the polycystic kidney and hepatic disease 1 (*PKHD1*) gene are responsible for ARPKD, the severity of which depends on the type of mutations.¹¹ The *PKHD1* gene is located on chromosome 6p21 and encodes a fibrocystin protein that localises to the primary cilium of renal epithelial cells. There is a high risk of fetal presentation and neonatal death if the fetus carries two truncating mutations.¹²

Inherited ciliopathies may also cause multisystem pathology, which may be severe and result in early death for many patients. Aside from cystic kidney disease, other common clinical features of ciliopathies include hepatobiliary disease, laterality defects, polydactyly, agenesis of corpus callosum, retinal degeneration and occipital encephalocele.¹³ Ciliopathies with prominent renal phenotypes include NPHP, JBTS and MKS.

NPHP is an autosomal-recessive disorder responsible for 6–10% of ESRD in children.¹⁴ NPHP is

characterised by cysts that are typically restricted to the corticomedullary junction region of the kidney, and the kidney size is normal or reduced.¹⁵ The disease is genetically heterogeneous. Mutations in over 20 different recessive genes (including *NPHP1–NPHP19*, *AHI1* and *XPNPEP3*) have been identified in about 50% of NPHP patients.¹⁶ Infantile NPHP is a disease that progresses to ESRD usually before the age of 2 years and is characterised by cortical microcysts associated with tubulointerstitial lesions. Classically, it is linked to *NPHP2/INVS* gene encoding inversin, but patients carrying *NPHP3* mutations may also develop the infantile phenotype frequently associated with liver involvement.¹⁷

JBTS is neurodevelopmental disorder characterised by cerebellar vermis aplasia (CVA), a significant malformation of the cerebellum that is linked to ataxia and may be seen on brain MRI as ‘molar tooth sign’.¹⁸ JBTS follows an autosomal-recessive inheritance pattern and there are currently over 26 known causative genes.^{19–21} JBTS may be associated with cystic renal disease in a subset of cases.

MKS is a prenatally lethal autosomal-recessive condition characterised by occipital encephalocele, bilateral renal cystic dysplasia, hepatic ductal proliferation, fibrosis, cysts and polydactyly.²² Patients with MKS invariably die from respiratory and/or renal failure. Genetic heterogeneity of MKS has been established with now 13 reported genes involved.^{23–25}

For many of these ciliopathy syndromes, significant phenotypic variability has been observed even between members of the same family, making clinical diagnosis, prediction of clinical progression and genetic counselling a challenge.

Antenatal screening using ultrasound scanning (USS) is a means by which cystic kidney disease can be readily detected. Serial ultrasound evaluation starting from 11 weeks of gestation onwards can be used as a screening modality.²⁶ Abnormal findings that point towards a renal ciliopathy include increased size of kidneys, a bright echotexture (hyperechogenicity) and a loss of the normal corticomedullary differentiation. Perinatal ultrasound appearance of kidneys can look similar in fetuses with ARPKD, perinatal-onset ADPKD, MKS and some forms of NPHP. In addition to renal anomalies, prenatal ultrasound can detect other features of ciliopathies such as encephalocele, polydactyly, situs inversus, agenesis of the corpus callosum, Dandy–Walker malformation, fibrosis of the liver and structural heart defects.²²

In this study, we have combined antenatal ultrasound examination of the fetus and targeted molecular genetic ‘panel testing’ for inherited renal disorders to characterise a cohort of Saudi Arabian patients who presented antenatally with features of an inherited renal ciliopathy.

MATERIALS AND METHODS

Study cohort

The cohort consists of 44 Saudi Arabian families where there was evidence of antenatal USS anomalies of the kidney, which included cystic kidney disease, enlarged kidneys and echogenic kidneys. Additional antenatal USS findings including central nervous system (CNS) anomalies (encephalocele, CVA, ventriculomegaly), cardiac defects (congenital heart malformation, pericardial effusion) and skeletal defects (narrow thorax, polydactyly) were documented. Clinical phenotypes postnatally were also reviewed, including postnatal renal USS. For molecular genetic investigations, the cohort was divided into two groups: Group A, where DNA was available from the affected fetuses and their parents ($n=34$ families) and Group B, where DNA was available from both parents but not the affected child

($n=10$ families) (table 1). Following informed consent, DNA was extracted from available chorionic villus sampling, amniotic fluid, placental blood or peripheral blood cells using the Gentra Systems PUREGENE DNA Isolation kit (Qiagen, Valencia, California, USA). Ethical and study permissions were approved by the Research Advisory Council of King Faisal Specialist Hospital, Riyadh, Saudi Arabia (RAC#2050 045). We confirm that all the diagnostic genetic work was performed in Saudi Arabia with full ethical approval. The UK centre acted in an advisory and strategic manner to direct the study.

Antenatal USS examination

Prenatal anatomy USS examination was performed at the Obstetrics and Gynecology Department, King Faisal Specialist Hospital and Research Centre, between weeks 18 and 22 of pregnancy. For cases with a known family history of cystic kidney disease/ciliopathy serial, antenatal USS examinations started between 12 and 22 weeks. For new referrals and unknown family history, antenatal USS started at the first visit. Fetal anatomy was reported as either normal or abnormal with explanations for features that includes cranium, cerebral ventricles, posterior fossa, face, spine, chest, cardiac four-chamber view, cardiac outflow tracts, heart axis, cardiac situs, stomach, bowel, kidneys, bladder, abdominal cord insertion, number of cord vessels, upper extremities and presence of hands, and lower extremities and presence of feet. Published reference values for renal length and volume²⁷ and for renal volumes based on three-dimensional ultrasound²⁸ were used. Fetal death was defined as an intrauterine death greater than 10 weeks of gestation. A perinatal death is defined as a death within 7 days of birth, and an infant death is defined as a death within 1 year of birth.

Maternal cell contamination and molecular karyotyping

In all fetal DNA samples, maternal cell contamination was excluded by using the AmpFLSTR Identifier PCR Amplification Kit as described by the manufacturer (Applied Biosystems, Life Technologies, Paisley, UK). Where available fetal DNA was used for molecular karyotyping (Affymetrix CytoScan HD Array Kit, Santa Clara, California, USA) to exclude chromosomal aneuploidy and to determine regions of homozygosity in the affected patient.

Targeted renal genes panel and next generation sequencing

A customised 90 renal genes panel that includes ciliopathy genes (including 3 polycystic kidney disease genes, 10 NPHP genes, 9 JBTS genes and 11 MKS genes) as well as and other inherited renal disorders (see online supplementary table S1) was prepared using Life Technologies proprietary AmpliSeq multiplexing assay. This panel has previously undergone validation for its analytical sensitivity and specificity using 107 renal patients and had 89% base reads on target with a read depth (base coverage) of 840 after alignment and a of 98% coverage of all genes.²⁹ All samples were prepared within the Saudi Human Genome Project Laboratories and loaded onto a Proton I chip, and sequencing was performed on an Ion Proton system (Ion Torrent—Life Technologies) as recommended by the manufacturer and as previously described.²⁹

NGS analysis pipeline

The analysis pipeline for processing the next generation sequencing (NGS) reads went through several steps. Reads were examined for quality and parts of reads with low-quality value were trimmed out. The reads were then aligned to the human reference genome GRCh37/Hg19 with the Torrent Mapping

Table 1 Clinical and molecular findings in cohort of antenatal cystic kidney disease and ciliopathy phenotypes

A or B	Family	Consanguinity	Outcome	Renal phenotype	Oligohydramnios/ anhydramnios	Encephalocele	Other CNS abnormalities	Skeletal/ growth malformations	Other defects	Number of other affected fetus/siblings	Segregation and unaffected sib	Gene	Mutation	Remarks and ExAC MAF
A	FT-3	Yes	Fetal death	Cystic		Yes		Polydactyly		0	m,f (1× unaffected sib-het)	<i>B9D1</i>	Homo c.508_510delCTC p.L170del	Novel
A	FT-1	Yes	Fetal death	Enlarged, echogenic with cysts	Yes	Yes		Polydactyly	Cystic hygroma	2	m,f	<i>CC2D2A</i>	Homo c.3084delG p.R1028Rfs*3	Reported ³⁰ (MAF=0.00002548)
A	FT-6	Not known	Fetal death	Cystic	Yes		CVA, dilated cisterna magna, corpus callosum agenesis			0	m,f (1× unaffected sib-het)	<i>CC2D2A</i>	Homo c.3364C>T p. P1122S	Reported ³¹
A	FT-8	Yes	Fetal death	Enlarged, echogenic with cysts	Yes	Yes	Corpus callosum agenesis and holoprosencephaly			1	m,f	<i>CC2D2A</i>	Homo c.4531T>C p. W1511R	Reported ³²
A	FT-14	Yes	Fetal death	Cystic		Yes		Spina bifida		2	m,f	<i>CC2D2A</i>	Homo c.4531T>C p. W1511R	Reported ³²
A	FT-15	Yes	Fetal death	Cystic		Yes		Intrauterine growth restriction		2	m,f	<i>CC2D2A</i>	Homo c.3084delG p.R1028Rfs*3	Reported ³⁰ (MAF=0.00002548)
A	FT-21	Yes	Fetal death	Cystic		Yes		Clubfoot		0	m,f	<i>CC2D2A</i>	Homo c.3084delG p.R1028Rfs*3	Reported ³⁰ (MAF=0.00002548)
A	FT-26	Yes	Fetal death	Cystic		Yes			Ascites	2	m	<i>CC2D2A</i>	Homo c.4437 +1G>A	Novel
A	FT-7	Not known	Alive at 6 m	Cystic			CVA			0	m,f	<i>CEP290</i>	Homo c.5668G>T p. G1890*	Reported ³³ (MAF=0.0001432)
A	FT-9	Yes	Perinatal death	Enlarged echogenic	Yes		Ventriculomegaly			1	m,f (unaffected sib –het, unaffected sib –wt)	<i>INVS</i>	Homo c.1760delA p.Q587Rfs*2	Novel
A	FT-27	Yes	Fetal death	Cystic	Yes	Yes		Clubfoot		1	m	<i>MKS1</i>	Homo c.417 +1G>A	Novel
A	FT-5	Yes	Fetal death	Cystic	Yes	Yes				3	m,f	<i>MKS1</i>	Homo c.1066C>T p.Q356*	Novel
A	FT-13	Yes	Fetal death	Cystic		Yes				3	m	<i>MKS1</i>	Homo c.1066C>T p.Q356*	Novel
A	FT-31	Yes	Infant death (8 mo)	Cystic					Congenital heart malformation, lung hypoplasia	0	m,f	<i>PKHD1</i>	Homo c.4870C>T p. R1624W	Reported ³⁰ (MAF=0.0001812)
A	FT-33	Yes	Alive at 12 mo	Cystic					Hepatic cysts	0	m,f	<i>PKHD1</i>	Homo c.4870C>T p. R1624W	Reported ³⁰ (MAF=0.0001812)

Continued

Table 1 Continued

A or B	Family	Consanguinity	Outcome	Renal phenotype	Oligohydramnios/ anhydramnios	Encephalocele	Other CNS abnormalities	Skeletal/ growth malformations	Other defects	Number of other affected fetuses/ siblings	Segregation and unaffected sib	Gene	Mutation	Remarks and ExAC MAF
A	FT-34	Yes	Alive at 14 mo	Cystic						0	m,f	<i>PKHD1</i>	Homo c.4870C>T p. R1624W	Reported ³⁰ (MAF=0.0001812)
A	FT-19	Yes	Fetal death	Cystic	Yes					3	m,f (1× unaffected sib-het)	<i>RPGRIP1L</i>	Het c.640G>A p. V214I Het c.685G>A p. A229T	V214I Reported ³⁴ (V214I, MAF=0.0005292)
A	FT-20	Not known	Perinatal death	Enlarged echogenic	Yes	Yes		Micrognathia		0	m,f	<i>TCTN2</i>	Homo c.1852C>T p.Q618*	Novel
A	FT-10	Yes	Fetal death	Enlarged, echogenic with cysts	Yes			Narrow thorax, dolichocephaly		0	m,f	<i>TMEM67</i>	Homo c.457T>G p.C153G	Novel
A	FT-22	Yes	Fetal death	Enlarged cystic	Yes		CVA, hydrocephalus		Congenital heart malformation, pericardial effusion	2	m,f	<i>TMEM67</i>	Homo c.1413-2A>G	Novel
A	FT-18	Yes	Perinatal death	Cystic	Yes	Yes	Corpus callosum agenesis	Clubfoot	Hepatic cysts	1	m,f	<i>TMEM231</i>	Homo c.751G>A p.V251I	Reported ³²
A	FT-23	Yes	Fetal death	Increased echogenicity	Yes		CVA, dilated cisterna magna, Dandy–Walker malformation		Pericardial effusion	1			Unsolved	
A	FT-28	Yes	Fetal death	Increased echogenicity	Yes		Dandy–Walker malformation	Polydactyly		2			Unsolved	
A	FT-4	Yes	Fetal death	Cystic	Yes		CVA, dilated cisterna magna	Narrow thorax	Pericardial effusion	1			Unsolved	
A	FT-12	Yes	Fetal death	Increased echogenicity	Yes			Dolichocephaly		2			Unsolved	
A	FT-11	Yes	Perinatal death	Cystic	Yes					0			Unsolved	
A	FT-17	No	Alive at 36 mo	Cystic			CVA, dilated cisterna magna			0			Unsolved	
A	FT-24	Yes	Fetal death	Cystic	Yes			Narrow thorax		0			Unsolved	
A	FT-25	Yes	Fetal death	Cystic	Yes					0			Unsolved	
A	FT-29	No	Fetal death	Cystic	Yes	Yes				0			Unsolved	
A	FT-30	Yes	Fetal death	Cystic	Yes		Ventriculomegaly			1			Unsolved	
A	FT-16	Not known	Fetal death	Enlarged kidneys	Yes					0			Unsolved	

Continued

Table 1 Continued

A or B	Family	Consanguinity	Outcome	Renal phenotype	Oligohydramnios/ anhydramnios	Encephalocele	Other CNS abnormalities	Skeletal/ growth malformations	Other defects	Number of other affected fetus/ siblings	Segregation and unaffected sib	Gene	Mutation	Remarks and ExAC MAF
A	FT-2	Yes	Fetal death	Cystic	Yes		CVA, dilated cisterna magna			3			Unsolved	
A	FT-32	Yes	Alive at 20 mo	Cystic	Yes					0			Unsolved	
B	FT-35	Yes	Fetal death	Cystic			CVA, dilated cisterna magna			3	m,f	<i>CC2D2A</i>	Presumed homo c.3084delG p. R1028Rfs*3	Reported ³⁰ (MAF=0.00002548)
B	FT-40	Yes	Fetal death	Cystic						1	m,f	<i>CEP290</i>	Presumed homo c.3777_3778delAG p.R1259Sfs*16	Novel
B	FT-43	Yes	Fetal death	Cystic		Yes				2	m,f	<i>MKS1</i>	Presumed homo c.1066C>T p. Q356*	Novel
B	FT-36	Yes	Fetal death	Cystic	Yes		CVA, dilated cisterna magna	Bilateral bowed femurs		0	m,f	<i>NEK8</i>	Presumed homo c.1401G>A p. W467*	Novel (MAF=0.00008237)
B	FT-41	Yes	Fetal death	Cystic					Congenital heart malformation	1	m,f	<i>NPHP3</i>	Presumed homo c.2694-1_-2delAG	Novel (MAF=0.0003553)
B	FT-45	Yes	Fetal death	Cystic	Yes				Hepatomegaly	1	m,f	<i>PKHD1</i>	Presumed homo c.3539G>A p. G1180E	Novel
B	FT-42	Yes	Fetal death	Cystic						2	m,f	<i>TCTN2</i>	Presumed homo c.252_253delTG	Novel
B	FT-37	Yes	Fetal death	Cystic						1			Unsolved	
B	FT-38	Yes	Fetal death	Cystic						0			Unsolved	
B	FT-44	Yes	Fetal death	Cystic						1			Unsolved	

Novel mutations are in bold.

A: samples where DNA from affected and parent(s) was available. B: Samples where maternal and paternal DNA was available and mutation is presumed (with a 25% chance) to be causative.

CVA, cerebellar vermis aplasia; CNS, central nervous system; f, father; het, heterozygous; homo, homozygous; m, mother; MAF, minor allele frequency; mo, month; sib, sibling.

Alignment Program (TMAP) Aligner software. Once the reads were aligned, the variants were called using the Torrent Variant Caller (TVC) program. The TMAP and the TVC programs are distributed as part of the Torrent Suite (<https://github.com/iontorrent/TS>) package. The resulting variant files were stored in variant call format (VCF) files. The VCF file generated for each sample was processed through an annotation pipeline against databases such as OMIM, GenBank, dbSNP 1000 genome project, Human Gene Mutation Database, and a local database (SGP737) of 550 patients containing Arab-specific variants. Variants with a minor allele frequency (MAF) >1% were discounted.

In addition to allele frequency, annotation provides pathogenicity scores, homozygosity/heterozygosity, read quality scores and other parameters used to identify candidate causative variants. All NGS and targeted sequencing and bioinformatics analysis were performed at the Saudi Human Genome Project Laboratories at KFSHRC and KACST.

For predicting the damaging effect of the reported mutation, four in silico prediction tools were used: PolyPhen-2 (<http://genetics.bwh.harvard.edu/pph/>), Provean (<http://provean.jcvi.org/index.php>), MutationTaster (<http://www.mutationtaster.org/>) and Human Splicing Finder (<http://www.umd.be/HSF/#>). Reported allele frequency of all putative pathogenic variants was determined using the ExAC database (<http://exac.broadinstitute.org/>), and evolutionary conservation was determined from sequence alignments using MutationTaster and UCSC (<https://genome.ucsc.edu>).

Sanger sequencing

Direct sequencing of PCR amplicons was carried out to confirm positive gene panel results. PCR was performed using Qiagen (Manchester, UK) master mix kit. Oligonucleotide primers for PCR amplification of targeted genomic DNA were designed using Primer3 software (<http://frodo.wi.mit.edu/>) and synthesised by Metabion International AG (Munich, Germany). Primer sequences are available on request. Following treatment with the Agencourt AMPure PCR purification system (Agencourt Bioscience, Beverly, Massachusetts, USA), products were sequenced using BigDye Terminator Cycle Sequencing kit (PE Applied Biosystems, Massachusetts, USA) and run on an ABI 3730xl capillary sequencer. Sequences were analysed using Mutation Surveyor software V3.24 (SoftGenetics LLC, State College, Pennsylvania, USA).

RESULTS AND DISCUSSION

A cohort of 44 families were analysed, where 38 (86%) were known to be consanguineous and 26 (59%) had more than one affected fetus. The antenatal renal USS findings included either bilateral cystic kidney disease, echogenic kidneys or enlarged kidneys in all cases (table 1). Antenatal USS also detected extra-renal malformations at a high rate (figure 1): 25 (57%) had oligohydramnios or anhydramnios, 14 (32%) had encephalocele and 9 (20%) had CVA. Other anomalies included limb defects including polydactyly and structural cardiac defects. The phenotype of this cohort was extremely severe with 38 (86%) cases dying as stillborn infants or perinatally. Only six cases survived the perinatal period (table 1).

Where patient DNA was available (group A, n=34), none of the cases had evidence of chromosomal aneuploidy (data not shown) and therefore malformation syndromes associated with renal cysts, such as trisomy 13 (Patau), trisomy 18 (Edward) and trisomy 21 (Down), were excluded.

Using the renal gene panel in this cohort, 96.98% coverage of target genes was achieved, with an average base coverage of

Figure 1 Prenatal ultrasound images of affected fetuses. (A and B) FT-8: transverse view of the fetal abdomen at 33 weeks of gestation (A), showing enlarged echogenic kidneys. Transverse view of the fetal head at 33 weeks of gestation (B), showing a cystic mass arising from the occipital area of the fetal head representing an encephalocele. (Genotype: *CC2D2A* homozygous mutation.) (C) FT-10: a transverse view of the fetal abdomen at 23 weeks of gestation, showing enlarged echogenic kidneys with cystic areas. (Genotype: *TMEM67* homozygous mutation.) (D) FT-20: a transverse view of the fetal kidneys at 33 weeks, showing enlarged echogenic kidneys. (Genotype: *TCTN2* homozygous mutation.) (E and F) FT-1: a transverse view of the fetal kidneys at 31 weeks of gestation (E), showing enlarged echogenic cystic kidneys. A transverse view of the fetal head at 31 weeks of gestation (F), showing a cystic mass arising from the fetal occiput, which represents an encephalocele. (Genotype: *CC2D2A* homozygous mutation.) (G) FT-9: a transverse view of the fetal abdomen at 27 weeks of gestation showing bilateral enlarged echogenic kidneys. (Genotype: *INVS* homozygous mutation.) (H) FT-13: a sagittal view of a fetus at 12 weeks of gestation showing a mass arising from the posterior aspect of the fetal head, which represents an encephalocele. (Genotype: *MKS1* homozygous mutation.) (I) FT-22: a transverse view of the fetal abdomen at 18 weeks, showing enlarged kidneys with cystic changes. (Genotype: *TMEM67* homozygous mutation.) (J) FT-21: a transverse view of the fetal head at 16 weeks of gestation, showing a mass arising from the posterior aspect of the fetal head, which represents an encephalocele. (Genotype: *CC2D2A* homozygous allele.)

>500. Twenty-one patients from 34 families in cohort A had a molecular genetic diagnosis (62% mutation detection rate). Seven families from the 10 families in cohort B had an inferred molecular genetic diagnosis by the finding of identical pathogenic alleles in both parents who were known to be consanguineous. The chances of the affected child inheriting both these alleles would be 25%. The identified rare alleles identified in these parental samples are listed in online supplementary table S2. In each of the families, there is only a single rare heterozygous change that was identified and confirmed using Sanger sequencing in both parents allowing a genetic diagnosis to be inferred (see online supplementary figure S1). All mutations identified were confirmed and segregation analysis was performed (including screening unaffected siblings) using Sanger sequencing. Mutations in genes *B9D1*, *CC2D2A*, *CEP290*, *INVS*, *MKS1*, *NEK8*, *PKHD1*, *RPGRIP1L*, *TCTN2*, *TMEM67* and *TMEM231* were identified (table 1) with a total of 13 novel variants detected in this study (table 2 and figure 2). Pathogenic rare (<1% MAF) sequence variants were not detected in the other renal panel genes, in particular digenic or oligogenic changes in renal ciliopathy genes were not seen. A common *RPGRIP1L* missense variant in its heterozygous state was identified as a third allele in two cases and is discussed below. Consistent with the lethal phenotypes seen in this cohort, mutations in *B9D1*, *CC2D2A*, *CEP290*, *MKS1*, *RPGRIP1L*, *TCTN2*, *TMEM67* and *TMEM231* are all known to cause a MKS phenotype. Mutations in *INVS* and *NEK8* have been reported in severe neonatal forms of NPHP with numerous extrarenal features.³⁵ Phenotypes in some of the patients

with mutations with *PKHD1* mutations were comparatively less severe, accounting for three of the cases which survived beyond the perinatal period.

In group A (34 families), homozygous mutations were detected in 20 families with just one family with compound heterozygous mutations (FT-19), in keeping with the known high rates of consanguinity. In group B (10 families), homozygous mutations were inferred by finding identical heterozygous variants in both parents in seven cases, consistent with the known parental consanguinity. These mutations were presumed to be found in their homozygous state in the affected patient. Unfortunately, direct sequencing of patient DNA or any unaffected siblings was not available in these cases. All mutations detected were either previously reported (and known to be pathogenic) or novel and predicted to be pathogenic by using in silico scores (table 2). Novel mutations were all homozygous (or inferred to be homozygous from parental samples) and included predicted missense, frameshift, nonsense and splicing defects. The types of mutations detected in this cohort seem to correlate closely with the phenotypes observed. Most mutations detected were truncating, frameshift and splice site mutations. These mutations were often lethal, causing fetal death or perinatal death. In this study, six missense mutations resulted in fetal or perinatal death.

Mutations in *CC2D2A* gene were the most common cause of antenatally detected cystic kidney disease in our cohort, accounting for eight cases. All patients with *CC2D2A* mutations had severe CNS abnormalities; six had evidence of an encephalocoele indicative of a MKS phenotype and two had evidence of

Table 2 In silico analysis of novel mutations

Gene	Mutation	Reference sequence	Mutation type	Provean	PolyPhen-2	Mutation Taster	Human Splicing Finder	ExAC database	Evolutionary conservation
<i>B9D1</i>	c.508_510delCTC (p.L170del)	NM_015681	Indel	Deleterious (−7.568)	N/A	Disease causing (0.989)		Absent	<i>Caenorhabditis elegans</i>
<i>CC2D2A</i>	c.4437+1G>A	NM_001080522	Splice site	N/A	N/A	N/A	Donor site broken	Absent	<i>Perkinsus marinus</i>
<i>CEP290</i>	c.3777_3778delAG (p.R1259Sfs*16)	NM_025114	Deletion, frameshift	N/A	N/A	Disease causing (1.000)		Absent	<i>Danio rerio</i>
<i>INVS</i>	c.1760delA (p.Q587Rfs*2)	NM_014425	Deletion, frameshift	N/A	N/A	Disease causing (1.000)		Absent	<i>Xenopus tropicalis</i>
<i>MKS1</i>	c.417+1G>A	NM_017777	Splice site	N/A	N/A	N/A	Donor site broken	Absent	<i>D. rerio</i>
<i>MKS1</i>	c.1066C>T (p.Q356*)	NM_017777	Nonsense	N/A	N/A	Disease causing (1.000)		Absent	<i>D. rerio</i>
<i>NEK8</i>	c.1401G>A (p.W467*)	NM_178170	Nonsense	N/A	N/A	Disease causing (1.000)		1 het. allele reported (MAF=0.000008237)	<i>X. tropicalis</i>
<i>NPHP3</i>	c.2694-1_2delAG	NM_153240	Deletion, frameshift	N/A	N/A	N/A	Acceptor site broken	43 het. alleles reported (MAF=0.0003553)	<i>D. rerio</i>
<i>PKHD1</i>	c.3539G>A (p.G1180E)	NM_138694	Missense	Deleterious (−6.296)	Probably damaging (0.999)	Disease causing (0.761)		Absent	<i>Mus musculus</i>
<i>TCTN2</i>	c.252_253delTG (p.V85Dfs*24)	NM_024809	Deletion, frameshift	N/A	N/A	Disease causing (1.000)		Absent	<i>X. tropicalis</i>
<i>TCTN2</i>	c.1852C>T (p.Q618*)	NM_024809	Nonsense	N/A	N/A	Disease causing (1.000)		Absent	<i>X. tropicalis</i>
<i>TMEM67</i>	c.457T>G (p.C153G)	NM_153704	Missense	Deleterious (−7.289)	Probably damaging (1.000)	Disease causing (0.999)		Absent	<i>C. elegans</i>
<i>TMEM67</i>	c.1413-2A>G	NM_153704	Splice site	N/A	N/A	N/A	Acceptor site broken	Absent	<i>D. rerio</i>

Evolutionary conservation at the protein level for non-synonymous changes was analysed by comparing the wild-type amino acid in the human with other orthologues in lower species. The lowest species where exact conservation of amino acid was preserved is shown. Het, heterozygous; MAF, minor allele frequency; N/A, not applicable.

Figure 2 Novel mutations identified in cohort with antenatal cystic kidney disease and ciliopathy phenotypes. (A) Three nonsense (B), four splice site (C), three frameshift (D), two missense and (E) one deletion novel mutations (boxed) were detected homozygously in patients or heterozygously in both parents with ciliopathy phenotypes. (Just one parental chromatogram is shown but a comparison of maternal and paternal chromatograms is shown in online supplementary figure S1). Family number (FT) is shown as well as mutation and predicted translational changes. Healthy control sequence is shown alongside. Intron–exon boundaries are marked with a vertical dashed line.

CVA with a dilated cysterna magna, in keeping with a JBTS phenotype. Typical MKS phenotypes were seen in three patients with *MKS1* mutations, which included two novel changes (table 2). The novel nonsense *MKS1* mutation (p.Q356*) was present in two families (FT-5 and FT-13), suggesting that these families were related. In family FT-18, the *TMEM231* mutation (p.V251I) led to perinatal death in two fetuses. The mutation is predicted to cause a splicing defect (table 2), due to its position as the last nucleotide in exon 4 of the *TMEM231*, although it appears to be a missense change.

PKHD1 gene mutations were found in four families. One case (FT-31) had associated lung hypoplasia and cardiac malformations and another (FT-33) had evidence of intrahepatic cysts (table 1). It has been reported that truncating mutations in *PKHD1* gene may be lethal.¹² In this study, mutations detected in *PKHD1* gene were homozygous missense mutations rather than truncating mutations, one of which was novel (c.3539G>A; p.G1180E) and led to a perinatal death in the proband and a sibling (FT-45). Three families shared the c.4870C>T (p.R1624W) mutation, which has been reported previously (also in its homozygous state) in a Saudi Arabian patient, with a ‘later-onset’ ARPKD phenotype.³⁰

Despite known consanguinity, compound heterozygous variants in *RPGRIP1L* were identified in family FT-19, with four affected siblings. The first variant c.640G>A (V214I) (rs139067427) is rare with a reported allele frequency of 0.05% and is predicted to be pathogenic (table 2). The second *RPGRIP1L* variant c.685G>A (p.A229T) (rs61747071) was identified in this family following Sanger sequencing of exon 6 of *RPGRIP1L*, as MAF filtering via our data pipeline had excluded this relatively common variant (MAF of 3.7%). The pathogenicity of this variant has been previously explored,³⁴

and the variant has been shown to compromise the interaction of *RPGRIP1L* with *RPGR*. To determine the frequency of this allele in our patient cohort, Sanger sequencing of *RPGRIP1L* exon 6 was performed. The rs61747071 variant was also present heterozygously in affected patients from FT-8 (with a homozygous *CC2D2A* missense mutation) and FT-10 (with a homozygous *TMEM67* missense mutation). The additional pathogenicity of this allele in these patients is unknown.

Compound heterozygous mutations in *B9D1* have previously been associated with MKS,³⁶ but this gene remains a rare cause of renal ciliopathies. The fetus in this case presented with posterior encephalocele and bilaterally enlarged multicystic dysplastic kidneys and bilateral clubfeet (but not polydactyly). An additional disease allele in *CEP290* identified in the fetus may have modified the phenotype.³⁶ The *B9D1* protein has structural similarities to *MKS1* and similar severe phenotypes would be predicted. More recently, mutations in *B9D1* have been described in two unrelated patients (aged 7 and 9 years) and with JBTS and a neurological limited phenotype, suggesting a wider phenotypic spectrum.³⁷

Mutations in *NEK8* are also a rare cause of a renal ciliopathy. Previously, homozygous mutations in *NEK8* have been described in a Kurdish child with kidney microcysts and likely NPHP, reaching ESRD at 14 years of age (c.1273C>T, p.H425Y)³⁸ and in three stillborn fetuses with enlarged cystic kidneys and cystic changes in the liver and pancreas (c.1795C>T, p.R599*).³⁵ Some fetuses had additional features including heterotaxy, truncus arteriosus and other structural heart defects, hypoplastic lungs and skeletal anomalies (bowed femurs). Here, we identified a single stillborn fetus (FT-36) with a novel nonsense change in *NEK8* (c.1401G>A, p.W467*) who had cystic kidneys, oligohydramnios, CVA and bilateral bowing of the femurs. This

nonsense mutation is predicted to disrupt the highly conserved regulator of chromatin condensation 1 (RCC1) domain and is in proximity to the murine *ick* mutation (p.G448V).³⁹

Antenatal presentations of cystic kidney disease are often associated with severe phenotypes and poor outcomes. These can include early presentation of ADPKD, or more commonly in consanguineous families, a presentation of an autosomal-recessive renal ciliopathy disease, as we have seen in this cohort. Extrarenal manifestations on the antenatal USS such as encephalocele and CVA may suggest MKS or JBTS phenotypes, respectively. Other features such as polydactyly and thoracic cage abnormalities may point towards other ciliopathies such as BBS⁴⁰ or skeletal dysplasias such as Jeune syndrome.⁴¹

Screening for ciliopathy genes in the diagnostic setting, especially in the perinatal period, is challenging. While whole exome sequencing (WES) is one possible approach, targeted gene panel exome sequencing may be preferable in diagnostic laboratories for specificity, deliverability and low cost. A disease-specific gene panel approach avoids the common difficulty of reporting secondary genetic findings that often occurs following WES. However, any predesigned NGS gene panel will be limited to known genes directed towards specific phenotypes and will not allow for recently discovered ciliopathy genes to be screened. Our gene panel contained 90 genes and included the 3 known polycystic kidney disease genes (*PKD1*, *PKD2* and *PKHD1*) and 11 of the 12 known MKS genes (see online supplementary table S1). However, it included only 10 of the 21 genes known to cause NPHP and 9 of the 26 JBTS genes. Therefore, our panel was biased towards (and very effective at) diagnosing MKS in this cohort with very severe disease phenotypes, but the precise molecular genetic diagnosis remained unknown in others. Indeed, in the 14 cases whom had antenatal USS evidence of an encephalocele suggestive of a MKS phenotype, all except 2 had a molecular genetic diagnosis. A recent study confirmed the strong cystic kidney disease phenotype in MKS patients, where cystic kidneys were found in (97.7%) of MKS cases.²²

To improve diagnostic yield, unsolved samples via our panel gene testing could be subjected to WES, especially in cases where there are DNA samples available from more than one affected in each family. However, this approach is more costly and can be more time consuming, when compared with a targeted panel approach. We hope to develop an updated renal gene panel in the near future, as the NGS sequencing platform we have developed will allow for additional genes (and their amplicons) to be analysed. In this study cohort of antenatal cases, the mutation detection rate was higher than reported by others^{36–42} who used PCR exon sequencing alone. More recently, a combination of WES and targeted resequencing of a ciliopathy gene panel was successfully used in a cohort of patients with Jeune asphyxiating thoracic dystrophy.⁴³

In summary, using a cohort of patients with antenatal evidence of kidney disease and associated ciliopathy syndromes, we have performed targeted genetic panel testing using patient and/or parental DNA samples to reveal the molecular genetic diagnosis in 64% of patients. Our high detection rate of homozygous disease-causing alleles reflects a high underlying rate of consanguinity. We would predict a reduction in diagnostic yield in less consanguineous populations. The genetic spectrum remains wide and certainly we have not identified a reason to narrow our diagnostic panel, rather it should be expanded to capture more recently reported genetic causes of developmental renal disease. It is interesting to note that *CC2D2A* mutations were the commonest cause of an antenatal ciliopathy in our cohort, but the genetic heterogeneity of inherited cystic kidney

disease is also borne out by our study. In our population, renal gene panel testing provided diagnostic information that was valuable to clinicians, genetic counsellors and families. A molecular genetic diagnosis provides an accurate diagnosis, which is hugely valuable when there are such severe phenotypes affecting one or more family members and can be used to predict recurrence rates and allow planning, including preimplantation genetic diagnosis for future pregnancies.

Author affiliations

- ¹Genetics Department, King Faisal Specialist Hospital and Research Centre, Riyadh, Saudi Arabia
- ²Obstetrics and Gynecology Department, King Faisal Specialist Hospital and Research Centre, Riyadh, Saudi Arabia
- ³Obstetrics & Gynecology Department, Prince Sultan Military Medical City, Riyadh, Saudi Arabia
- ⁴Institute of Genetic Medicine, International Centre for Life, Newcastle University, Newcastle upon Tyne, UK
- ⁵Saudi Human Genome Project, King Abdulaziz City for Science and Technology (KACST), Riyadh, Saudi Arabia
- ⁶Medicine Department, King Faisal Specialist Hospital and Research Centre, Riyadh, Saudi Arabia

Acknowledgements We thank all the families for participating in the study. We also thank all team members at Saudi Human Genome project for help in conducting the study. In addition, we thank sequencing and genotyping core facilities at Genetics Department at the Research Centre, KFSH&RC for doing sequencing and molecular karyotyping. Special thanks for Ms Ghadah A Al Dakheel for preparing figures for publication. JAS acknowledges support from the Northern Counties Kidney Research Fund. All NGS library building, sequencing and bioinformatics analyses were performed by the Saudi Human Genome Project at King Abdulaziz City for Science and Technology (KACST) and at KFSHRC.

Contributors MHA-H, JAS and BM conceived of the study and participated in its design and coordination, and drafted and revised the manuscript. BAA, FSB, DK, DM and NA performed the NGS sequencing. AA and NE performed in silico modelling. HA-J, AA, AIT, DK, NE, BAA and FSB carried out all technical aspects of molecular diagnosis and helped with in silico modelling. WK, NA-S, ZA, RA, MT, MA and RK participated in the clinical diagnosis of the cases. SM analysed molecular karyotyping. TF, ME-K and MA carried out bioinformatics analysis. DM and NA-T helped conceive the study and drafted the manuscript. All authors read and approved the final manuscript.

Competing interests None declared.

Ethics approval Research Advisory Council of King Faisal Specialist Hospital, Riyadh, Saudi Arabia (RAC#2050 045).

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

- 1 Zhang Q, Taulman PD, Yoder BK. Cystic kidney diseases: all roads lead to the cilium. *Physiology (Bethesda)* 2004;19:225–30.
- 2 Hassanien AA, Al-Shaikh F, Vamos EP, Yadegarfar G, Majeed A. Epidemiology of end-stage renal disease in the countries of the Gulf Cooperation Council: a systematic review. *JRSM Short Rep* 2012;3:38.
- 3 Arts HH, Kooers NV. Current insights into renal ciliopathies: what can genetics teach us? *Pediatr Nephrol* 2013;28:863–74.
- 4 Hildebrandt F. Genetic kidney diseases. *Lancet* 2010;375:1287–95.
- 5 Gabow PA. Autosomal dominant polycystic kidney disease. *Am J Kidney Dis* 1993;22:511–12.
- 6 Igarashi P, Somlo S. Genetics and pathogenesis of polycystic kidney disease. *J Am Soc Nephrol* 2002;13:2384–98.
- 7 Zerres K, Rudnik-Schöneborn S, Deget F. Childhood onset autosomal dominant polycystic kidney disease in sibs: clinical picture and recurrence risk. German Working Group on Paediatric Nephrology (Arbeitsgemeinschaft für Pädiatrische Nephrologie). *J Med Genet* 1993;30:583–8.
- 8 Vujic M, Heyer CM, Ars E, Hopp K, Markoff A, Orndal C, Rudenhe B, Nasr SH, Torres VE, Torra R, Bogdanova N, Harris PC. Incompletely penetrant PKD1 alleles mimic the renal manifestations of ARPKD. *J Am Soc Nephrol* 2010;21:1097–102.

- 9 Losekoot M, Ruivenkamp CA, Tholens AP, Grimbergen JE, Vijfhuizen L, Vermeer S, Dijkman HB, Cornelissen EA, Bongers EM, Peters DJ. Neonatal onset autosomal dominant polycystic kidney disease (ADPKD) in a patient homozygous for a PKD2 missense mutation due to uniparental disomy. *J Med Genet* 2012;49:37–40.
- 10 Torres VE, Harris PC. Mechanisms of disease: autosomal dominant and recessive polycystic kidney diseases. *Nat Clin Pract Nephrol* 2006;2:40–55; quiz 55.
- 11 Denamur E, Delezoide AL, Alberti C, Bourillon A, Gubler MC, Bouvier R, Pascaud O, Elion J, Grandchamp B, Michel-Calemard L, Missy P, Zaccaria I, Le Nagard H, Gerard B, Loirat C, Barbet J, Beaufriere AM, Berchel C, Bessieres B, Boudjemaa S, Buenerd A, Carles D, Clemenson A, Dechelotte P, Devisme L, Dijoud F, Esperandieu O, Fallet C, Gonzales M, Hillion Y, Jacob B, Joubert M, Kermanach P, Lallemand A, Laquerriere A, Laurent N, Liprandi A, Loeuillet L, Loget P, Martinovic J, Menez F, Nancy F, Roux JJ, Rouleau-Dubois C, Sinico M, Tantau J, Wann AR. Genotype–phenotype correlations in fetuses and neonates with autosomal recessive polycystic kidney disease. *Kidney Int* 2010;77:350–8.
- 12 Bergmann C, Senderek J, Windelen E, Kupper F, Middeldorf I, Schneider F, Dornia C, Rudnik-Schoneborn S, Konrad M, Schmitt CP, Seeman T, Neuhaus TJ, Vester U, Kirfel J, Buttner R, Zerres K. Clinical consequences of PKHD1 mutations in 164 patients with autosomal-recessive polycystic kidney disease (ARPKD). *Kidney Int* 2005;67:829–48.
- 13 Sharma N, Berbari NF, Yoder BK. Ciliary dysfunction in developmental abnormalities and diseases. *Curr Top Dev Biol* 2008;85:371–427.
- 14 Hildebrandt F, Benzing T, Katsanis N. Ciliopathies. *N Engl J Med* 2011;364:1533–43.
- 15 Simms RJ, Eley L, Sayer JA. Nephronophthisis. *Eur J Hum Genet* 2009;17:406–16.
- 16 Wolf MT. Nephronophthisis and related syndromes. *Curr Opin Pediatr* 2015;27:201–11.
- 17 Tory K, Rousset-Rouviere C, Gubler MC, Moriniere V, Pawtowski A, Becker C, Guyot C, Gié S, Frishberg Y, Nivet H, Deschênes G, Cochat P, Gagnadoux MF, Saunier S, Antignac C, Salomon R. Mutations of NPHP2 and NPHP3 in infantile nephronophthisis. *Kidney Int* 2009;75:839–47.
- 18 Valente EM, Salpietro DC, Brancati F, Bertini E, Galluccio T, Tortorella G, Briuglia S, Dallapiccola B. Description, nomenclature, and mapping of a novel cerebello-renal syndrome with the molar tooth malformation. *Am J Hum Genet* 2003;73:663–70.
- 19 Kroes HY, Monroe GR, van der Zwaag B, Duran KJ, de Kovel CG, van Roosmalen MJ, Harakalova M, Nijman IJ, Kloosterman WP, Giles RH, Knoers NV, van Haaften G. Joubert syndrome: genotyping a Northern European patient cohort. *Eur J Hum Genet* 2016;24:214–20.
- 20 Srour M, Hamdan FF, McKnight D, Davis E, Mandel H, Schwartzentruber J, Martin B, Patry L, Nassif C, Dionne-Laporte A, Ospina LH, Lemyre E, Massicotte C, Laframboise R, Maranda B, Labuda D, Decarie JC, Rypens F, Goldsher D, Fallet-Bianco C, Soucy JF, Laberge AM, Maffei C, Boycott K, Brais B, Boucher RM, Rouleau GA, Katsanis N, Majewski J, Elpeleg O, Kukulich MK, Shalev S, Michaud JL. Joubert syndrome in French Canadians and identification of mutations in CEP104. *Am J Hum Genet* 2015;97:744–53.
- 21 Lambacher NJ, Bruel AL, van Dam TJ, Szymanska K, Slaats GG, Kuhns S, McManus GJ, Kennedy JE, Gaff K, Wu KM, van der Lee R, Burglen L, Doumard R, Riviere JB, Faviere L, Attie-Bitach T, Saunier S, Curd A, Peckham M, Giles RH, Johnson CA, Huynen MA, Thauvin-Robinet C, Blacque OE. TMEM107 recruits ciliopathy proteins to subdomains of the ciliary transition zone and causes Joubert syndrome. *Nat Cell Biol* 2016;18:122–31.
- 22 Barisic I, Boban L, Loane M, Garne E, Wellesley D, Calzolari E, Dolk H, Addor MC, Bergman JE, Braz P, Draper ES, Haeusler M, Khoshnood B, Klungsoyr K, Pierini A, Queisser-Luft A, Rankin J, Rissmann A, Verellen-Dumoulin C. Meckel–Gruber Syndrome: a population-based study on prevalence, prenatal diagnosis, clinical features, and survival in Europe. *Eur J Hum Genet* 2015;23:746–52.
- 23 Szymanska K, Hartill VL, Johnson CA. Unraveling the genetics of Joubert and Meckel–Gruber syndromes. *J Pediatr Genet* 2014;3:65–78.
- 24 Filges I, Nosova E, Bruder E, Tercanli S, Townsend K, Gibson WT, Rothlisberger B, Heinemann K, Hall JG, Gregory-Evans CV, Wasserman WW, Miny P, Friedman JM. Exome sequencing identifies mutations in KIF14 as a novel cause of an autosomal recessive lethal fetal ciliopathy phenotype. *Clin Genet* 2014;86:220–8.
- 25 Shaheen R, Almoisheer A, Faqeh E, Babay Z, Monies D, Tassan N, Abouelhoda M, Kurdi W, Al Mardawi E, Khalil MM, Seidahmed MZ, Alnemer M, Alsahan N, Sogaty S, Alhashem A, Singh A, Goyal M, Kapoor S, Alomar R, Ibrahim N, Alkuraya FS. Identification of a novel MKS locus defined by TMEM107 mutation. *Hum Mol Genet* 2015;24:5211–18.
- 26 Dias T, Sairam S, Kumarasiri S. Ultrasound diagnosis of fetal renal abnormalities. *Best Pract Res Clin Obstet Gynaecol* 2014;28:403–15.
- 27 Gloor JM, Breckle RJ, Gehrkling WC, Rosenquist RG, Mulholland TA, Bergstralh EJ, Ramin KM, Ogburn PL, Jr. Fetal renal growth evaluated by prenatal ultrasound examination. *Mayo Clin Proc* 1997;72:124–9.
- 28 Yu C, Chang C, Chang F, Ko H, Chen H. Fetal renal volume in normal gestation: a three-dimensional ultrasound study. *Ultrasound Med Biol* 2000;26:1253–6.
- 29 Saudi Mendeliome Group. Comprehensive gene panels provide advantages over clinical exome sequencing for Mendelian diseases. *Genome Biol* 2015;16:134.
- 30 Onuchic LF, Furu L, Nagasawa Y, Hou X, Eggemann T, Ren Z, Bergmann C, Senderek J, Esquivel E, Zeltner R, Rudnik-Schoneborn S, Mrug M, Sweeney W, Avner ED, Zerres K, Guay-Woodford LM, Somlo S, Germino GG. PKHD1, the polycystic kidney and hepatic disease 1 gene, encodes a novel large protein containing multiple immunoglobulin-like plexin-transcription-factor domains and parallel beta-helix 1 repeats. *Am J Hum Genet* 2002;70:1305–17.
- 31 Gorden NT, Arts HH, Parisi MA, Coene KL, Letteboer SJ, van Beersum SE, Mans DA, Hikida A, Eckert M, Knutzen D, Alswaid AF, Ozyurek H, Dibooglu S, Otto EA, Liu Y, Davis EE, Hutter CM, Bammler TK, Farin FM, Dorschner M, Topcu M, Zackai EH, Rosenthal P, Owens KN, Katsanis N, Vincent JB, Hildebrandt F, Rubel EW, Raible DW, Knoers NV, Chance PF, Roepman R, Moens CB, Glass IA, Doherty D. CC2D2A is mutated in Joubert syndrome and interacts with the ciliopathy-associated basal body protein CEP290. *Am J Hum Genet* 2008;83:559–71.
- 32 Shaheen R, Faqeh E, Alshammari MJ, Swaid A, Al-Gazali L, Mardawi E, Ansari S, Sogaty S, Seidahmed MZ, AlMotairi MI, Farra C, Kurdi W, Al-Rasheed S, Alkuraya FS. Genomic analysis of Meckel–Gruber syndrome in Arabs reveals marked genetic heterogeneity and novel candidate genes. *Eur J Hum Genet* 2013;21:762–8.
- 33 Sayer JA, Otto EA, O'Toole JF, Nurnberg G, Kennedy MA, Becker C, Hennies HC, Helou J, Attanasio M, Fausett BV, Utsch B, Khanna H, Liu Y, Drummond I, Kawakami I, Kusakabe T, Tsuda M, Ma L, Lee H, Larson RG, Allen SJ, Wilkinson CJ, Nigg EA, Shou C, Lillo C, Williams DS, Hoppe B, Kemper MJ, Neuhaus T, Parisi MA, Glass IA, Petry M, Kispiert A, Gloy J, Ganner A, Walz G, Zhu X, Goldman D, Nurnberg P, Swaroop A, Leroux MR, Hildebrandt F. The centrosomal protein nephrocystin-6 is mutated in Joubert syndrome and activates transcription factor ATF4. *Nat Genet* 2006;38:674–81.
- 34 Khanna H, Davis EE, Murga-Zamalloa CA, Estrada-Cuzcano A, Lopez I, den Hollander AI, Zonneveld MN, Othman MI, Waseem N, Chakarova CF, Maubaret C, Diaz-Font A, MacDonald I, Muzny DM, Wheeler DA, Morgan M, Lewis LR, Logan CV, Tan PL, Beer MA, Inglehearn CF, Lewis RA, Jacobson SG, Bergmann C, Beales PL, Attie-Bitach T, Johnson CA, Otto EA, Bhattacharya SS, Hildebrandt F, Gibbs RA, Koenekeop RK, Swaroop A, Katsanis N. A common allele in RPGRIP1L is a modifier of retinal degeneration in ciliopathies. *Nat Genet* 2009;41:739–45.
- 35 Frank V, Habbig S, Bartram MP, Eisenberger T, Veenstra-Knol HE, Decker C, Boersma RA, Gobel H, Nurnberg G, Griessmann A, Franke M, Borgal L, Kohli P, Volker LA, Dotsch J, Nurnberg P, Benzing T, Bolz HJ, Johnson C, Gerkes EH, Schermer B, Bergmann C. Mutations in NEK8 link multiple organ dysplasia with altered Hippo signalling and increased c-MYC expression. *Hum Mol Genet* 2013;22:2177–85.
- 36 Hopp K, Heyer CM, Hommerding CJ, Henke SA, Sundsbak JL, Patel S, Patel P, Consugar MB, Czarnecki PG, Gliem TJ, Torres VE, Rossetti S, Harris PC. B9D1 is revealed as a novel Meckel syndrome (MKS) gene by targeted exon-enriched next-generation sequencing and deletion analysis. *Hum Mol Genet* 2011;20:2524–34.
- 37 Romani M, Micalizzi A, Kraoua I, Dotti MT, Cavallin M, Sztrihla L, Ruta R, Mancini F, Mazza T, Castellana S, Hanene B, Carluccio MA, Darra F, Mate A, Zimmermann A, Gouider-Khouja N, Valente EM. Mutations in B9D1 and MKS1 cause mild Joubert syndrome: expanding the genetic overlap with the lethal ciliopathy Meckel syndrome. *Orphanet J Rare Dis* 2014;9:72.
- 38 Otto EA, Trapp ML, Schultheiss UT, Helou J, Quarmby LM, Hildebrandt F. NEK8 mutations affect ciliary and centrosomal localization and may cause nephronophthisis. *J Am Soc Nephrol* 2008;19:587–92.
- 39 Liu S, Lu W, Obara T, Kuida S, Lehoczy J, Dewar K, Drummond IA, Beier DR. A defect in a novel Nek-family kinase causes cystic kidney disease in the mouse and in zebrafish. *Development* 2002;129:5839–46.
- 40 Ashkinadze E, Rosen T, Brooks SS, Katsanis N, Davis EE. Combining fetal sonography with genetic and allele pathogenicity studies to secure a neonatal diagnosis of Bardet–Biedl syndrome. *Clin Genet* 2013;83:553–9.
- 41 Schramm T, Gloning KP, Minderer S, Daumer-Haas C, Hortnagel K, Nerlich A, Tschek B. Prenatal sonographic diagnosis of skeletal dysplasias. *Ultrasound Obstet Gynecol* 2009;34:160–70.
- 42 Tallila J, Salonen R, Kohlschmidt N, Peltonen L, Kestila M. Mutation spectrum of Meckel syndrome genes: one group of syndromes or several distinct groups? *Hum Mutat* 2009;30:E813–30.
- 43 Schmidts M, Frank V, Eisenberger T, Al Turki S, Bizet AA, Antony D, Rix S, Decker C, Bachmann N, Bald M, Vinke T, Toenshoff B, Di Donato N, Neuhaus T, Hartley JL, Maher ER, Bogdanovic R, Peco-Antic A, Mache C, Hurles ME, Joksic I, Guc-Scekic M, Dobricic J, Brankovic-Magic M, Bolz HJ, Pazour GJ, Beales PL, Scambler PJ, Saunier S, Mitchison HM, Bergmann C. Combined NGS approaches identify mutations in the intraflagellar transport gene IFT140 in skeletal ciliopathies with early progressive kidney Disease. *Hum Mutat* 2013;34:714–24.